



Macintosh Videolaryngoscope for intubation in the operating room: a comparative quality improvement project

Alexandre Laplace-Builhé

► To cite this version:

Alexandre Laplace-Builhé. Macintosh Videolaryngoscope for intubation in the operating room: a comparative quality improvement project. Human health and pathology. 2020. dumas-03139754

HAL Id: dumas-03139754

<https://dumas.ccsd.cnrs.fr/dumas-03139754>

Submitted on 12 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

UNIVERSITE DE MONTPELLIER
FACULTE DE MEDECINE MONTPELLIER-NÎMES

THESE

Pour obtenir le titre de

DOCTEUR EN MEDECINE

Présentée et soutenue publiquement

Par

Alexandre LAPLACE-BUILHÉ

Le 06 mars 2020

Titre :

**"Quel Vidéolaryngoscope Macintosh utiliser au bloc opératoire ?
Une évaluation prospective multicentrique"**

Directrice de thèse : Dr Audrey DE JONG

JURY :

Monsieur le Professeur Samir JABER, Président

Monsieur le Professeur Pascal COLSON, Assesseur

Monsieur le Professeur Gérald CHANQUES, Assesseur

Madame le Docteur Audrey DE JONG, Assesseur

UNIVERSITE DE MONTPELLIER
FACULTE DE MEDECINE MONTPELLIER-NÎMES

THESE

Pour obtenir le titre de

DOCTEUR EN MEDECINE

Présentée et soutenue publiquement

Par

Alexandre LAPLACE-BUILHÉ

Le 06 mars 2020

Titre :

**"Quel Vidéolaryngoscope Macintosh utiliser au bloc opératoire ?
Une évaluation prospective multicentrique"**

Directrice de thèse : Dr Audrey DE JONG

JURY :

Monsieur le Professeur Samir JABER, Président

Monsieur le Professeur Pascal COLSON, Assesseur

Monsieur le Professeur Gérald CHANQUES, Assesseur

Madame le Docteur Audrey DE JONG, Assesseur



ANNEE UNIVERSITAIRE 2019 - 2020

PERSONNEL ENSEIGNANT

Professeurs Honoriaires

ALLIEU Yves	CALLIS Albert	JAFFIOL Claude	NAV RATIL Henri
ALRIC Robert	CANAUD Bernard	JANBON Charles	OTHONIEL Jacques
ARNAUD Bernard	CHAPTAL Paul-André	JANBON François	PAGES Michel
ASTRUC Jacques	CIURANA Albert-Jean	JARRY Daniel	PEGURET Claude
AUSSILLOUX Charles	CLOT Jacques	JOURDAN Jacques	PELISSIER Jacques
AVEROUS Michel	COSTA Pierre	LAFFARGUE François	POUGET Régis
AYRAL Guy	D'ATHIS Françoise	LALLEMAND Jean Gabriel	PUJOL Henri
BAILLAT Xavier	DEMAILLE Jacques	LAMARQUE Jean-Louis	RABISCHONG Pierre
BALDET Pierre	DESCOMPS Bernard	LAPEYRIE Henri	RAMUZ Michel
BALDY-MOULINIER Michel	DIMEGLIO Alain	LE QUELLEC Alain	RIEU Daniel
BALMES Jean-Louis	DUBOIS Jean Bernard	LESBROS Daniel	ROCHEFORT Henri
BALMES Pierre	DUJOLS Pierre	LOPEZ François Michel	ROUANET DE VIGNE LAVIT Jean Pierre
BANSARD Nicole	DUMAS Robert	LORIOT Jean	SAINT AUBERT Bernard
BAYLET René	DUMAZER Romain	LOUBATIERES Marie Madeleine	SANCHO-GARNIER Hélène
BILLIARD Michel	ECHENNE Bernard	MAGNAN DE BORNIER Bernard	SANY Jacques
BLARD Jean-Marie	FABRE Serge	MARY Henri	SEGNARBIEUX François
BLAYAC Jean Pierre	FREREBEAU Philippe	MATHIEU-DAUDE Pierre	SENAC Jean-Paul
BLOTMAN Francis	GALIFER René Benoît	MEYNADIER Jean	SERRE Arlette
BONNEL François	GODLEWSKI Guilhem	MICHEL François-Bernard	SOLASSOL Claude
BOURGEOIS Jean-Marie	GRASSET Daniel	MION Charles	THEVENET André
BRUEL Jean Michel	GUILHOU Jean-Jacques	MION Henri	VIDAL Jacques
BUREAU Jean-Paul	HERTAULT Jean	MIRO Luis	VISIER Jean Pierre
BRUNEL Michel	HUMEAU Claude	NAVARRO Maurice	



Professeurs Emérites

ARTUS Jean-Claude	MARES Pierre
BLANC François	MAUDELONDE Thierry
BOULENGER Jean-Philippe	MAURY Michèle
BOURREL Gérard	MILLAT Bertrand
BRINGER Jacques	MONNIER Louis
CLAUSTRES Mireille	MOURAD Georges
DAURES Jean-Pierre	PREFAUT Christian
DAUZAT Michel	PUJOL Rémy
DAVY Jean-Marc	RIBSTEIN Jean
DEDET Jean-Pierre	SCHVED Jean-François
ELEDJAM Jean-Jacques	SULTAN Charles
GROLLEAU RAOUX Robe	TOUCHON Jacques
GUERRIER Bernard	UZIEL Alain
GUILLOT Bernard	VOISIN Michel
LANDAIS Paul	ZANCA Michel



PU-PH de classe exceptionnelle

ALBAT Bernard	Chirurgie thoracique et cardiovasculaire
ALRIC Pierre	Chirurgie vasculaire ; médecine vasculaire (option chirurgie vasculaire)
BACCINO Eric	Médecine légale et droit de la santé
BASTIEN Patrick	Parasitologie et mycologie
BLAIN Hubert	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
BONAFE Alain	Radiologie et imagerie médicale
CAPDEVILA Xavier	Anesthésiologie-réanimation et médecine péri-opératoire
CHAMMAS Michel	Chirurgie orthopédique et traumatologique
COLSON Pascal	Anesthésiologie-réanimation et médecine péri-opératoire
COMBE Bernard	Rhumatologie
COTTALORDA Jérôme	Chirurgie infantile
COUBES Philippe	Neurochirurgie
COURTET Philippe	Psychiatrie d'adultes ; addictologie
CRAMPETTE Louis	Oto-rhino-laryngologie
CRISTOL Jean Paul	Biochimie et biologie moléculaire
DE LA COUSSAYE Jean Emmanuel	Médecine d'urgence
DE WAZIERES Benoît	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
DELAPORTE Eric	Maladies infectieuses ; maladies tropicales
DEMOLY Pascal	Pneumologie ; addictologie
DOMERGUE Jacques	Chirurgie viscérale et digestive
DUFFAU Hugues	Neurochirurgie
ELIAOU Jean François	Immunologie
FABRE Jean Michel	Chirurgie viscérale et digestive
FRAPIER Jean-Marc	Chirurgie thoracique et cardiovasculaire
HAMAMAH Samir	Biologie et Médecine du développement et de la reproduction ; gynécologie médicale



Gynécologie-obstétrique ; gynécologie médicale

Médecine physique et de réadaptation

Anesthésiologie-réanimation et médecine péri-opératoire
Médecine interne ; gérontologie et biologie du vieillissement,
médecine générale, addictologie

Médecine intensive-réanimation

Thérapeutique ; médecine d'urgence ; addictologie

Biophysique et médecine nucléaire

Neurologie

Gastroentérologie ; hépatologie ; addictologie

Anesthésiologie-réanimation et médecine péri-opératoire

Biochimie et Biologie moléculaire

Chirurgie thoracique et cardiovasculaire

Physiologie

Cardiologie

Oto-rhino-laryngologie

Pédiatrie

Gastroentérologie ; hépatologie ; addictologie

Biologie cellulaire

Chirurgie vasculaire ; médecine vasculaire (option médecine
vasculaire)

Endocrinologie, diabète et maladies métaboliques ; gynécologie
médicale

Maladies infectieuses, maladies tropicales

Anesthésiologie-réanimation et médecine péri-opératoire

Cancérologie ; radiothérapie

Maladies infectieuses ; maladies tropicales

Radiologie et imagerie médicale

Bactériologie-virologie ; hygiène hospitalière

Cancérologie ; radiothérapie



PU-PH de 1^{re} classe

AGUILAR MARTINEZ Patricia	Hématologie ; transfusion
ASSENAT Éric	Gastroentérologie ; hépatologie ; addictologie
AVIGNON Antoine	Nutrition
AZRIA David	Cancérologie ; radiothérapie
BAGHDADLI Amaria	Pédopsychiatrie ; addictologie
BEREGI Jean-Paul	Radiologie et imagerie médicale
BLANC Pierre	Gastroentérologie ; hépatologie ; addictologie
BORIE Frédéric	Chirurgie viscérale et digestive
BOULOT Pierre	Gynécologie-obstétrique ; gynécologie médicale
CAMBONIE Gilles	Pédiatrie
CAMU William	Neurologie
CANOVAS François	Anatomie
CAPTIER Guillaume	Anatomie
CARTRON Guillaume	Hématologie ; transfusion
CAYLA Guillaume	Cardiologie
CHANQUES Gérald	Anesthésiologie-réanimation et médecine péri-opératoire
CORBEAU Pierre	Immunologie
COSTES Valérie	Anatomie et cytologie pathologiques
COULET Bertrand	Chirurgie orthopédique et traumatologique
CYTEVAL Catherine	Radiologie et imagerie médicale
DADURE Christophe	Anesthésiologie-réanimation et médecine péri-opératoire
DAUVILLIERS Yves	Physiologie
DE TAYRAC Renaud	Gynécologie-obstétrique, gynécologie médicale
DE VOS John	Histologie, embryologie et cytogénétique
DEMARIA Roland	Chirurgie thoracique et cardio-vasculaire



DEREURE Olivier	Dermatologie - vénérérologie
DROUPY Stéphane	Urologie
DUCROS Anne	Neurologie
DUPEYRON Arnaud	Médecine physique et de réadaptation
FESLER Pierre	Médecine interne ; gérontologie et biologie du vieillissement, médecine générale, addictologie
GARREL Renaud	Oto-rhino-laryngologie
GENEVIEVE David	Génétique
HAYOT Maurice	Physiologie
KLOUCHE Kada	Médecine intensive-réanimation
KOENIG Michel	Génétique
LAFFONT Isabelle	Médecine physique et de réadaptation
LAVABRE-BERTRAND Thierry	Histologie, embryologie et cytogénétique
LAVIGNE Jean-Philippe	Bactériologie-virologie ; hygiène hospitalière
LE MOING Vincent	Maladies infectieuses ; maladies tropicales
LECLERCQ Florence	Cardiologie
LEHMANN Sylvain	Biochimie et biologie moléculaire
MARIANO-GOULART Denis	Biophysique et médecine nucléaire
MATECKI Stéfan	Physiologie
MEUNIER Laurent	Dermato-vénérérologie
MOREL Jacques	Rhumatologie
NAVARRO Francis	Chirurgie viscérale et digestive
NOCCA David	Chirurgie viscérale et digestive
PETIT Pierre	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PERNEY Pascal	Médecine interne ; gérontologie et biologie du vieillissement, médecine générale, addictologie
PRUDHOMME Michel	Anatomie
PUJOL Jean Louis	Pneumologie ; addictologie
PURPER-OUAKIL Diane	Pédopsychiatrie ; addictologie



PU-PH de 2ème classe

BOURDIN Arnaud	Pneumologie ; addictologie
CANAUD Ludovic	Chirurgie vasculaire ; médecine vasculaire (option chirurgie vasculaire)
CAPDEVIELLE Delphine	Psychiatrie d'Adultes ; addictologie
CLARET Pierre-Géraud	Médecine d'urgence
COLOMBO Pierre-Emmanuel	Cancérologie ; radiothérapie
COSTALAT Vincent	Radiologie et imagerie médicale
CUVILLON Philippe	Anesthésiologie-réanimation et médecine péri-opératoire
DAIEN Vincent	Ophtalmologie
DORANDEU Anne	Médecine légale et droit de la santé
FAILLIE Jean-Luc	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
FUCHS Florent	Gynécologie-obstétrique ; gynécologie médicale
GABELLE DELOUSTAL Audrey	Neurologie
GAUJOUX Viala Cécile	Rhumatologie
GODREUIL Sylvain	Bactériologie-virologie ; hygiène hospitalière
GUILLAUME Sébastien	Psychiatrie d'adultes ; addictologie
GUILPAIN Philippe	Médecine Interne, gériatrie et biologie du vieillissement; addictologie
GUIU Boris	Radiologie et imagerie médicale
HERLIN Christian	Chirurgie plastique, reconstructrice et esthétique, brûlologie
HOUEDÉ Nadine	Cancérologie ; radiothérapie
JACOT William	Cancérologie ; Radiothérapie
JUNG Boris	Médecine intensive-réanimation
KALFA Nicolas	Chirurgie infantile



KOUYOUNDJIAN Pascal	Chirurgie orthopédique et traumatologique
LACHAUD Laurence	Parasitologie et mycologie
LALLEMAND Benjamin	Oto-rhino-laryngologie
LE QUINTREC DONNETTE Moglie	Néphrologie
LETOUZEY Vincent	Gynécologie-obstétrique ; gynécologie médicale
LONJON Nicolas	Neurochirurgie
LOPEZ CASTROMAN Jorge	Psychiatrie d'Adultes ; addictologie
LUKAS Cédric	Rhumatologie
MAURY Philippe	Chirurgie orthopédique et traumatologique
MILLET Ingrid	Radiologie et imagerie médicale
MORANNE Olivier	Néphrologie
MURA Thibault	Biostatistiques, informatique médicale et technologies de la communication
NAGOT Nicolas	Biostatistiques, informatique médicale et technologies de la communication
PANARO Fabrizio	Chirurgie viscérale et digestive
PARIS Françoise	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PASQUIE Jean-Luc	Cardiologie
PELLESTOR Franck	Histologie, embryologie et cytogénétique
PEREZ MARTIN Antonia	Chirurgie vasculaire ; médecine vasculaire (option médecine vasculaire)
POUDEROUX Philippe	Gastroentérologie ; hépatologie ; addictologie
RIGAU Valérie	Anatomie et cytologie pathologiques
RIVIER François	Pédiatrie
ROGER Pascal	Anatomie et cytologie pathologiques
ROSSI Jean François	Hématologie ; transfusion
ROUBILLE François	Cardiologie
SEBBANE Mustapha	Médecine d'urgence
SIRVENT Nicolas	Pédiatrie
SOLASSOL Jérôme	Biologie cellulaire

STOEBNER Pierre

Dermato-vénérérologie

SULTAN Ariane

Nutrition

THOUVENOT Éric

Neurologie

THURET Rodolphe

Urologie

VENAIL Frédéric

Oto-rhino-laryngologie

VILLAIN Max

Ophtalmologie

VINCENT Denis

Médecine interne ; gériatrie et biologie du vieillissement,
médecine générale, addictologie

VINCENT Thierry

Immunologie

WOJTUSCISZYN Anne

Endocrinologie-diabétologie-nutrition



PROFESSEURS DES UNIVERSITES

1^{re} classe :

COLINGE Jacques (Cancérologie, Signalisation cellulaire et systèmes complexes)

2^{ème} classe :

LAOUDJ CHENIVESSE Dalila (Biochimie et biologie moléculaire)

VISIER Laurent (Sociologie, démographie)

PROFESSEURS DES UNIVERSITES - Médecine générale

1^{re} classe :

LAMBERT Philippe

2^{ème} classe :

AMOUYAL Michel



PROFESSEURS ASSOCIES - Médecine Générale

CLARY Bernard

DAVID Michel

GARCIA Marc

PROFESSEURS ASSOCIES - Médecine

BESSIS Didier (Dermato-vénérérologie)

MEUNIER Isabelle (Ophtalmologie)

MULLER Laurent (Anesthésiologie-réanimation et médecine péri-opératoire)

PERRIGAULT Pierre-François (Anesthésiologie-réanimation et médecine péri-opératoire)

QUANTIN Xavier (Pneumologie)

ROUBERTIE Agathe (Pédiatrie)

VIEL Eric (Soins palliatifs et traitement de la douleur)

Maîtres de Conférences des Universités - Praticiens Hospitaliers

MCU-PH Hors classe

BADIOU Stéphanie	Biochimie et biologie moléculaire
BOULLE Nathalie	Biologie cellulaire
CACHEUX-RATABOUL Valère	Génétique
CARRIERE Christian	Bactériologie-virologie ; hygiène hospitalière
CHARACHON Sylvie	Bactériologie-virologie ; hygiène hospitalière
FABBRO-PERAY Pascale	Epidémiologie, économie de la santé et prévention
GIANSILY-BLAIZOT Muriel	Hématologie ; transfusion
HILLAIRE-BUYS Dominique	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PUJOL Joseph	Anatomie
RICHARD Bruno	Médecine palliative
RISPAIL Philippe	Parasitologie et mycologie
SEGONDY Michel	Bactériologie-virologie ; hygiène hospitalière

10 / 13



MCU-PH de 1^{re} classe

BERTRAND Martin	Anatomie
BOUDOUSQ Vincent	Biophysique et médecine nucléaire
BOURGIER Céline	Cancérologie ; Radiothérapie
BRET Caroline	Hématologie biologique
COSSEE Mireille	Génétique
GIRARDET-BESSIS Anne	Biochimie et biologie moléculaire
LAVIGNE Géraldine	Hématologie ; transfusion
LESAGE François-Xavier	Médecine et Santé au Travail
MATHIEU Olivier	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
MENJOT de CHAMPFLEUR Nicolas	Radiologie et imagerie médicale
MOUZAT Kévin	Biochimie et biologie moléculaire
OLIE Emilie	Psychiatrie d'adultes ; addictologie
PANABIERES Catherine	Biologie cellulaire
PHILIBERT Pascal	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
RAVEL Christophe	Parasitologie et mycologie
SCHUSTER-BECK Iris	Physiologie
STERKERS Yvon	Parasitologie et mycologie
THEVENIN-RENE Céline	Immunologie
TUAILLON Edouard	Bactériologie-virologie ; hygiène hospitalière

MCU-PH de 2^{ème} classe

CHIRIAC Anca	Immunologie
DE JONG Audrey	Anesthésiologie-réanimation et médecine péri-opératoire
DU THANH Aurélie	Dermato-vénéréologie
GOUZI Farès	Physiologie

HERRERO Astrid

Chirurgie viscérale et digestive

JEZIORSKI Éric

Pédiatrie

KUSTER Nils

Biochimie et biologie moléculaire

MAKINSON Alain

Maladies infectieuses, Maladies tropicales

PANTEL Alix

Bactérologie-virologie ; hygiène hospitalière

PERS Yves-Marie

Thérapeutique; addictologie

ROUBILLE Camille

Médecine interne ; gériatrie et biologie du vieillissement ;
addictologie

SZABLEWSKY

Anatomie et cytologie pathologiques



Maîtres de Conférences des Universités - Médecine Générale

MCU-MG de 1^{re} classe

COSTA David

MCU-MG de 2^{ème} classe

FOLCO-LOGNOS Béatrice

OUDE ENGBERINK Agnès

Maîtres de Conférences associés - Médecine Générale

LOPEZ Antonio

MILLION Elodie

PAVAGEAU Sylvain

REBOUL Marie-Catherine

SERAYET Philippe



Praticiens Hospitaliers Universitaires

BARATEAU Lucie	Physiologie
BASTIDE Sophie	Epidémiologie, économie de la santé et prévention
DAIEN Claire	Rhumatologie
GATINOIS Vincent	Histologie, embryologie et cytogénétique
GOULABCHAND Radjiv	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
LATTUCA Benoit	Cardiologie
MIOT Stéphanie	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PINETON DE CHAMBRUN Guillaume	Gastroentérologie ; hépatologie ; addictologie
SOUCHÉ François-Régis	Chirurgie viscérale et digestive

REMERCIEMENTS AUX ENSEIGNANTS

À Monsieur le Professeur Samir JABER

Je vous remercie de m'avoir fait l'honneur d'accepter de présider le jury de cette thèse.

Je vous remercie de m'avoir inculqué la rigueur nécessaire en Anesthésie-Réanimation lors de mes premiers semestres passés chez vous. Veuillez trouver ici le témoignage de mon profond respect. En parallèle merci pour la leçon de baby-foot reçue à l'occasion des soirées du DAR B.

À Monsieur le Professeur Pascal COLSON

Je vous remercie de me faire l'honneur de juger ce travail, de participer à mon jury de thèse et bien sûr merci de m'accueillir pour les deux prochaines années dans votre service. Merci pour votre confiance et votre soutien, soyez assuré de ma gratitude et de mon plus grand respect.

À Monsieur le Professeur Gérald CHANQUES

Merci de me faire l'honneur d'être dans mon jury de thèse, travailler avec toi depuis la fin de l'externat a été un réel plaisir ! Je pense qu'une grande partie de mon choix d'intégrer la famille Anesthésie-Réanimation est lié à notre rencontre. Sois assuré de mon dévouement.

À Madame le Docteur Audrey DE JONG

Je te remercie d'avoir accepté d'être ma directrice de thèse, grâce à toi j'ai pu participer à toutes les étapes de ce projet dès sa conception ! Merci pour ton écoute permanente, ta disponibilité, ta patience ainsi que de ton aide malgré les nombreuses responsabilités qui t'incombent.

AUTRES REMERCIEMENTS

À ma mère : Merci pour toute l'aide et le soutien reçus depuis le début de l'aventure Médecine notamment les nombreuses séances de Reiki ou les petits déjeuneurs au lit en PCEM1 !

À mon père : Malgré la distance tu as toujours été là pour moi... Grâce à toi je peux partager ma passion de l'histoire, merci de m'avoir appris également la persévérance !

À mon beau-père : Grâce à toi **Claude** j'ai pu découvrir l'univers fascinant de la musique, merci pour ta générosité et ton appui depuis notre rencontre

À ma chérie : Te rencontrer, **Anouchka**, à bouleversé ma vie ! Maintenant que nous sommes enfin ensemble, je découvre le bonheur. Plein de belles aventures nous attendent... Je t'aime !

À mes grands parents : **Papy Philippe**, j'aurais aimé que tu sois là pour cette soutenance malheureusement la Nouvelle-Calédonie est bien trop loin...merci pour ta générosité !

Mimouche, tu as toujours été ma seule et vraie grand-mère ! **Papy Jean**, tu n'es plus là pour ce jour important, malgré ton souhait que je fasse plutôt pharmacien, j'espère te rendre fier.

À mon frère : **Benjamin**, Frerot ! Malgré nos nombreuses ressemblances et divergences, on a toujours pu compter l'un sur l'autre !

À Robert : un véritable grand-père !

À mes beaux-parents : merci pour votre accueil chaleureux

À mes cousines : **Charlotte**, **Laura**, on a été très longtemps comme des frères et sœurs, j'espère qu'on se reverra d'avantage à l'avenir, merci de m'avoir suivi en médecine !

Fabian : On se connaît depuis si longtemps ! On a traversé toutes épreuves depuis la 6^{ème} ensemble en passant par les randonnées au fin fond de la Chine, l'ascension mythique du Mt Fuji jusqu'au concours de l'internat où tu es parti vers le côté obscur de la chirurgie... Merci pour ta fidélité toute ces années !

Hosam : Toi aussi tu fais partie des irréductibles du temps du Lycée Joffre, merci pour ta constante bonne humeur ! Petite dédicace au Kebab du Polygone sous l'escalator et nos concours de Harissa...

Bertrand : Tu as toujours été notre exemple à suivre depuis le lycée que ce soit du point de vu sportif que du point de vu intellectuel ! Merci pour la tenue chaude de rechange que tu avais prise au sommet du Fuji, sans elle, j'aurais pu y rester...

Pierre : Alors que tu étais le benjamin du groupe, notre petit frère, tu es parti beaucoup trop tôt... Je n'oublierais jamais nos sessions de jeu de Go, notre passion commune pour le Japon !

PE : Même si toi aussi tu deviendras chirurgien, je suis très fier de pouvoir te compter parmi mes proches amis ! Merci d'avance pour tous les avis Ortho à venir !

Guillaume : Une grande dédicace à nos fameuses Colles-Repas-Apéros de D4 !

À mes co-internes :

Jennifer : on a survécu ensemble à notre premier semestre biterrois, tu as été une vraie confidente toutes ces années d'internat !

Romain : à toutes nos heures passées au baby-foot de Lapeyronie ! Malheureusement je dois avouer que tu l'emportes...

Benjamin B. : merci pour ta grande générosité et nos soirées jeux de sociétés en commun avec

Elodie B. et Robin ! J'aurais vraiment aimé être ton co-chef au DAR D !

Philippe H. : grâce à toi et **Nicolas B.**, je sais maintenant changer une roue sur la bande d'arrêt d'urgence ! Merci pour la superbe ambiance que tu as mise dans tous nos stages en commun ! J'attends toujours ta recette secrète de l'Houmous...

Aurélien : j'ai hâte de commencer avec toi notre post-internat au DAR D !

Vivien : merci pour tous tes conseils avec **Julie B.** lorsque j'étais à un tournant de ma vie ! À ta bonne humeur permanente et à ton sens de la répartie inégalé !

Romaric : un véritable grand frère au temps du DAR B, puis un formidable moniteur de canoë !

À tous mes autres co-internes, notamment la Team Aréanimatrice du DAR C, **Luke, Léa** !

À tous mes chefs :

Yvan P. : merci pour ton aide à l'élaboration et la réalisation de ce travail !

Benjamin M. : pour ta musique de qualité au bloc Ortho !

Philippe B. : je me souviendrais toujours de ta proposition de m'héberger lorsque j'ai été en difficulté

Marion L. : pour ta gentillesse inégalée ! Tu es un vrai modèle à suivre

Philippe P. : à nos échanges cinéphiles !

Philippe C. : à tous nos jeudis « Thoraciques» à Nîmes...

À tous les autres chefs : Jacob, Norddine, Philippe G., Marine, Meriem, José, Fabien, Maxime V., Olivier C., Olivier M., Jordy, Nathalie, Marie, Christine, Jacques, Dominique, Alexandre M, Christopher, Maxime D., Kevin C., Chrystelle, Olivier R., Christophe D., Lucie, Maud, Jérémie, Christian P. et tous ceux que j'oublie, un grand merci pour tout !

Aux chirurgiens, nos indispensables partenaires !

Aux équipes d'infirmiers anesthésistes, aux infirmiers en réanimation et aux aide-soignant !

Table des matières

INTRODUCTION.....	21
MATERIALS AND METHODS	23
A. <i>Study Design</i>	23
B. <i>Population</i>	24
C. <i>Interventions</i>	24
D. <i>Data collection</i>	26
E. <i>Statistical analysis</i>	28
RESULTS.....	30
A. <i>Primary outcome</i>	30
B. <i>Secondary and exploratory outcomes</i>	30
DISCUSSION	32
CONCLUSION	36
REFERENCES	37
FIGURE LEGENDS	41
FIGURE 1. Methodology of the quality improvement project	43
FIGURE 2. Flow chart of the study	44
FIGURE 3. Main outcome (need of a hyperangulated blade) for each Macintosh videolaryngoscope after multivariable analysis.....	45
FIGURE 4. Subjective evaluation for each Macintosh videolaryngoscope	46
TABLE 1. Patients; airway and operator characteristics in the four Macintosh videolaryngoscopes groups	47
TABLE 2. Outcomes according to step I, II and III in the four Macintosh videolaryngoscope groups.....	48
SUPPLEMENTAL MATERIAL	50
SUPPLEMENTAL TABLE 1. Drugs characteristics in the four Macintosh videolaryngoscopes groups	51
SUPPLEMENTAL TABLE 2. Glottis visualization and alternative devices used in patients who reached step 3 in the four Macintosh videolaryngoscopes groups	52
SUPPLEMENTAL TABLE 3. Complications, subjective appreciation and quality of image in the four Macintosh videolaryngoscope groups	54

Macintosh Videolaryngoscope for intubation in the operating room: a comparative quality improvement project

Glossary:

VL = videolaryngoscopy

SpO₂ = peripheral oxygen saturation

CL = Cormack-Lehane

NRS = Numeric Rate Scale

SD = Standard deviation

CI = Confidence Interval

BMI = Body Mass index

INTRODUCTION

Standard Macintosh direct-laryngoscopy, which requires a direct line of sight to align airway axes, remains the first airway management device for most anesthesiologists.(1,2)

Recently the role of videolaryngoscopy (VL) in anticipated(3–5) and un-anticipated(6–9) difficult intubation has been widely recognized. High-profile national guidelines(6,7) have stated that all anesthesiologists should be trained to use and have immediate access to a VL. Following large implementation, difficult and failed intubation rates(10) by skilled providers declined significantly.(11,12) Furthermore, teaching with videolaryngoscopes can improve intubation skills in medical students.(9,13)

New devices, including APATM[®], C-MAC-S[®] (single-use), C-MAC-S-PM[®] (pocket and single-use) and McGrath Mac[®], are called “Macintosh” VLs.(5,14–17) They can be used both as direct and indirect-laryngoscopes. Two types of blades are available: first a Macintosh-style blade which integrates video capability and can be used to perform both indirect and direct-

laryngoscopy.(18) On the other hand a hyperangulated blade is available in order to further improve glottic visualization(19,20) for difficult laryngoscopy, which can be used only in indirect-laryngoscopy.(19)

To our knowledge, very limited data are available regarding the use of Macintosh-VLs as the first-intention devices for all intubation procedures performed in the operating room, no matter how difficult.

We performed a pragmatic study included in a quality improvement project aiming to compare the glottic visualization of different Macintosh-VLs used as the primary intubation device. The main hypothesis was that one of the Macintosh-VLs might show less frequent need to use the hyperangulated blade. The secondary hypotheses were that the Macintosh-VLs might differ regarding successful first attempt intubation, number of intubation attempts, glottic visualization in direct and indirect-laryngoscopy, adverse-effects, subjective assessment of difficulty of intubation, user-friendliness and image quality.

MATERIALS AND METHODS

A. Study Design

We conducted a quality improvement project from May 2017 to September 2017 in the 4 anesthesia departments of Montpellier Teaching Hospital, aiming at implementing Macintosh-VL for all intubations procedures in the operating room.

This study was part of an institutional assessment of airway management which evaluated single-use Macintosh-VLs. We obtained approval from the local scientific (Comité d'Organisation et de Gestion de l'Anesthésie Réanimation) and ethics committee (Institutional Review Board, Comité Locale d'Ethique Recherche, agreement number: 2017_CLER-MPT_11-04) of Montpellier University Hospital(21). Requirement for written informed consent was waived by the Institutional Review Board.

The four Macintosh-videolaryngoscopes assessed were loaned to the anesthesia departments for the duration of the 5 month trial. Representatives of each company providing devices for the trial afforded training in the use of the devices and support throughout the trial. The companies (including Medtronic) were otherwise not involved in this clinical assessment. The clinical evaluations were performed without funding or involvement in the decision to perform the trial, in the analysis of the results (performed by an independent statistician, NM), or in the subsequent decision-making. The departments of anesthesia had received free or at cost equipment for research and evaluation from all the airway companies providing single-use Macintosh VL at the time where the study was performed.

B. Population

Inclusion criteria

Patients aged ≥ 18 years that required oro-tracheal intubation during general anesthesia were included.

Exclusion criteria

Patients were not included if they met one or more of the following criteria: pregnancy or breast feeding, rapid sequence intubation, intubation in case of cardiopulmonary arrest, emergency airway management (severe hypoxemia or severe collapse before intubation), inter-incisor gap<2.2cm.

C. Interventions

Four devices were used including the McGrath Mac[®] (Medtronic Covidien, USA), the C-MAC-S[®] (Karl Storz, Germany), the C-MAC-S Pocket Monitor[®] (PM) (Karl Storz, Germany) and the APATM[®] (Advanced Airway Management Health Care, United Kingdom).

All the Macintosh-VLs were single-use and had both Macintosh-style blades and hyperangulated blades.

Before the beginning of the quality improvement project, Macintosh-VL were not available and only Glidescope[®] (Verathon, USA) VL was available and routinely used in case of difficult intubation.(1)

The anesthesiologists had no experience with the respective devices prior to participating in the study. In order to choose one Macintosh-VL for the institution, each of the four anesthesia departments assessed each of the four devices during one month in a randomized order: May (one period), June (one period), July- August together (due to the reduced number of patients undergoing anesthesia and surgery during the summer, one period) and September (one period).

One device was available for each anesthesia department (multiple patients could not be enrolled

simultaneously in the same anesthesia department, i.e. not consecutive included patients), and the anesthesiologists were encouraged to use the device whenever available.

All patients requiring tracheal intubation and without exclusion criteria could be included in the study. All the anesthesia providers at the participating institutions took part in the study.

All patients were monitored as usual with electrocardiogram, peripheral oxygen saturation (SpO_2) and arterial blood pressure (non-invasive or invasive as appropriate).

The anesthesia technique, including choice of drugs and order of administration,(22) size of blades and use of adjuvant airway device (ex, stylet, bougie), was at the discretion of the anesthesia provider providing patient care. In case of use of neuromuscular blockers, cisatracurium was universally used and muscle relaxation was assessed using a TOF monitoring. For each Macintosh-VL, three consecutive steps were performed in the same order and are summarized in Figure 1:

Step I: After preoxygenation for 3 minutes according to the institution protocol followed by anesthesia induction, the first step was to rate the glottic visualization, using the Cormack-Lehane (CL) scoring system,(23) using the Macintosh-VLs in direct-laryngoscopy with the Macintosh-style blade.

Step II: The second step was to rate the glottic visualization with the Macintosh-VLs in indirect-laryngoscopy (using the Macintosh-style blade); If the CL grade was \leq II, the operator made an intubation attempt, if the CL grade was \geq III, the operator moved to the third step without attempting intubation. Head repositioning or Backward Upward Rightward Pressure/ External Laryngeal Pressure was permitted at this step. Intubation using a bougie was not permitted with a CL III/IV view. The need to go to the third step was the primary outcome variable.

Step III: The third step was performed only in case of CL grade \geq III (III or IV) or if the intubation attempt at the second step failed. This step was mandatorily performed by a senior

anesthesiologist (defined as a physician, attending anesthesiologist). The Macintosh-style blade was switched to the hyperangulated blade. X blade for the McGrath Mac[©], D Blade for the C-MAC-S[©] and the C-MAC-S Pocket Monitor[©], Difficult Airway Blade (DAB) for the APA™[©]), using the same device. After rating the new glottic visualization (fourth rating) using the hyperangulated blade, the operator made an intubation attempt.

An intubation attempt was defined as the insertion of the laryngoscope blade into the mouth of the patient and the attempt to insert an endotracheal tube. At our institution usual practice is to stop the laryngoscopy and to resort to bag-mask ventilation in case of peripheral oxygen saturation dropping below 90%. The choice of an alternative technique after two failed intubation attempts was recommended by the study protocol and was in accordance with clinical standard.(7,8,24) The alternative technique was left at the discretion of the physician.

D. Data collection

The perioperative data collected by the operator consisted of:

- Demographic information (height, weight, age)
- Airway evaluation (Mallampati class, mouth opening, thyromental distance, neck range of motion, inter-incisor gap, medical history of difficult intubation)
- Type and amount of drugs administered during induction
- Operator status: physician, resident or anesthetic nurse
- Assessment of the glottic visualization using the CL scoring system
- Number of intubation attempt(s)
- Need to change to a different intubation device, and use of adjuvant airway devices (ex, stylet, Bougie). The decision to use a stylet for intubation at Step II was made by the operator before performing the first attempt.
- Vital signs (SpO₂, heart rate, blood pressure)

- Degree of ease or difficulty of tracheal intubation based on the Numeric Rate Scale (NRS) (0=easy to 10=difficult)
- Degree of user-friendliness (0= not user friendly to 10= totally user friendly) and quality of the image (0=bad to 10=perfect) based on NRS

Primary outcome

The primary outcome was the need to move to step III (using a hyperangulated blade because of failure to intubate at step II or CL grade III/IV at step III).

Secondary outcomes

- Glottic visualization assessed by CL score using the Macintosh-VL in direct-laryngoscopy with the Macintosh-style blade, the Macintosh-VL in indirect-laryngoscopy with the Macintosh-style blade and if step III was reached the Macintosh-VL in indirect-laryngoscopy with the hyperangulated blade
- Cormack I and II scores with the Macintosh-VL in indirect-laryngoscopy with the Macintosh-style blade (step II)
- Number of laryngoscopy attempts
- Adverse effects (hypoxemia defined by SpO₂<90%, bradycardia defined by heart rate <50 bpm, hypotension defined by systolic arterial pressure <90mmHg)
- Degree of ease or difficulty of tracheal intubation based on the NRS (0=easy to 10=difficult)
- Degree of user-friendliness (0= not user friendly to 10= totally user friendly) and quality of the image (0=bad to 10=perfect) based on NRS. No free text was accompanying this assessment, only an NRS was provided to the assessor.

Exploratory outcomes

- Proportion of successful attempts at tracheal intubation with the Macintosh-style blade in indirect-laryngoscopy (step II)

- Proportion of successful attempts at tracheal intubation with the hyperangulated blade (Step III)
- Failures/crossovers to other rescue techniques

E. Statistical analysis

Quantitative variables were expressed as means \pm standard deviation (SD) and qualitative variables were expressed as numbers (percentage). Comparison of means for unpaired data was performed using analysis of variance and Tukey's test for multiple comparisons. Comparison of percentages was performed using a Chi square test with Bonferroni correction for 6 comparisons (a P value <0.008 was considered significant). Confidence intervals (CI) were computed for associations of interest (99.2% CI after adjustment). For paired repeated data (Differences in CL score between steps), comparison was performed using a linear mixed effect model, for each VL group, including the subject as a random effect. Regarding outcomes recorded in step III, conditional nature of step III makes comparisons here well confounded by the primary outcome results and results of this conditional phase will be descriptive.

For the main outcome (use of a hyperangulated blade), a mixed effect multivariable logistic regression model was performed to assess the relation between the main outcome and the VL group and adjust for confounding. All variables with p-value less than 0.20 in the univariate analysis (as presented in Table 1), age (defined a priori) and body mass index (BMI, defined a priori) were entered into the model. The center was taken into account as a random effect. Pre-specified interactions between BMI and group, and age and group were tested.

All P values were two-tailed and exact. A P value <0.008 was considered statistically significant after Bonferroni correction for 6 comparisons. Statistical analysis was performed using SAS software version 9.4 (SAS Institute; Cary, NC).

Expecting a difference of use of the hyperangulated blade of 20% between two of the four Macintosh-VLs (from 30% to 10%),(25) we calculated the sample size as 97 per device, using a chi square test, with α level of 0.008 (taking into account multiple comparisons) and a power of 0.80. We decided to include 125 patients per device (total of 500 patients) to compensate for dropouts and missing data, and calculated that four months would allow to include at least the required number of subjects.

RESULTS

From May 2017 to September 2017, 589 patients were included (Figure 2). The demographic variables, operator characteristics and the choice of drugs did not differ between the four Macintosh-VLs (Table 1, Supplemental Table 1). Among airway characteristics, thyromental distance < 6 cm was significantly more frequent in the C-MAC-S-PM[®] and APATM[®] than in the McGrath Mac[®] and the C-MAC-S[®], whereas the presence of at least one predictive factor of difficult intubation did not significantly differ between Macintosh-VLs (Table 1).

A. Primary outcome

The hyperangulated blade was used in 22/180 patients (12%) for the McGrath Mac[®], in 29/139 patients (21%) for the C-MAC-S-PM[®], in 39/132 patients (30%) for the C-MAC-S[®] and in 35/138 patients (25%) for the APATM[®] (Table 2, Supplemental Figure 1). By multivariable analysis, the following variables were entered in the model: age, BMI, center, thyromental distance < 6 cm, no cervical spine extension. After adjustment for BMI, the McGrath Mac[®] was associated with less frequent use of hyperangulated blade than C-MAC-S[®] (OR, 99.2% CI of 0.34 (0.16, 0.77), p=0.0005), or APATM[®] (OR, 99.2% CI of 0.42 (0.19, 0.93), p=0.004), but not C-MAC-S-PM[®] (OR, 99.2% CI of 0.53 (0.23, 1.2), p=0.04, Figure 3).

B. Secondary and exploratory outcomes

Step I

In direct-laryngoscopy using the Macintosh-VLs (Macintosh-style blade), the CL scores were significantly lower with the McGrath Mac[®] than with the other Macintosh-VLs.

Step II

The glottic visualization with the Macintosh-VL was significantly better using indirect-laryngoscopy than using direct-laryngoscopy for all Macintosh-VLs (Table 2).

The number of Cormack I and II grades was significantly higher for the McGrath Mac[®] than for the C-MAC-S[®] and the APATM[®] (Table 2, Figure 4). The rates of successful intubation at step II were the following: n = 158/166 (95.2% (90.8 - 99.6)) for the McGrath Mac[®], n = 93/99 (93.9% (87.6 - 1.00) for the C-MAC-S[®], n = 110/121 (90.9% (84.0 - 97.8) for the C-MAC-S-PM[®], n = 103/113 (91.2% (84.1 - 98.2) for the APATM[®] (Table 2, Figure 4). Thirty-seven percent (37%) of intubation attempts were performed using a stylet (Table 2).

Step III

The rates of successful intubation at step III are presented in Table 2 and Supplemental Figure 1. Glottic visualization and alternative devices used in patients who reached step III in the four Macintosh-VLs are presented in Supplemental Table 2.

For all steps

Overall, the number of intubation attempts was significantly lower using the McGrath Mac[®] than the C-MAC-S[®] or the C-MAC-S-PM[®] VLs (Table 2). Intubation was successful in more than 90% of attempts, whatever the device used.

No serious adverse events occurred during the assessment period, and the rate of complications did not differ between Macintosh-VLs (Supplemental Table 3).

The subjective appreciation of difficulty of intubation and user friendliness of devices respectively showed lower and higher NRS scores for the McGrath Mac[®] device compared to the other devices (Supplemental Table 3 and Figure 4). The subjective assessment of image quality showed higher NRS scores for the C-MAC-S[®] and C-MAC-S-PM[®] compared to the APATM[®] or McGrath Mac[®] (Supplemental Table 3).

DISCUSSION

In this quality improvement project aiming to select one Macintosh-VL, the McGrath Mac[®] showed better performances than others Macintosh-VLs, reduced need to resort to a hyperangulated blade and showed greater user friendliness than other devices. However, C-MAC-S[®] and C-MAC-S-PM[®] showed better image quality than APATM[®] or McGrath Mac[®].

To our knowledge, this is the first time that the four single-use Macintosh-VLs on the market are assessed in a pragmatic human study. Laryngeal exposure was better with the McGrath Mac[®] though the image quality was not as good as the other devices (Figure 3), and the number of overall attempts were reduced with the McGrath Mac[®]. The McGrath Mac[®] was the only device which had a lower CI limit of first-attempt success rate higher than 90%, main criterion in a recent study by Kleine-Brueggeney et al.(25). Previous studies performed by Kleine-Brueggeney et al.(25)²⁵ in simulated difficult airway and by Shin et al.(26) in manikin which compared the McGrath Mac[®] and the reusable C-MAC[®] found no significant differences between the devices in terms of successful intubation rate and glottic visualization. One explanation to these apparent discrepancies are that, as required by French law, all the Macintosh-VLs devices used in the current study had single-use blade, whereas the reusable C-MAC[®] was assessed in the two other studies.(25,26) The blade used in the reusable C-MAC[®] differs from the blade used in the C-MAC S[®] and the C-MAC S[®] PM, being less thick in the reusable C-MAC[®]. Moreover, Shin's study(26) was a manikin study so the results cannot be fully extrapolated to human subjects, and Kleine et al.(25) excluded all subjects with known or predicted difficult airway ($BMI > 35 \text{ kg.m}^{-2}$, Mallampati score>III, thyromental distance>3.5cm, known difficult mask ventilation/laryngoscopy and planned or previous history of awake tracheal intubation). It is also worth noting that whereas the main aim of our pragmatic study was to assess the Macintosh-style blade in indirect-laryngoscopy, Kleine et al.(25) focused on the hyperangulated blade dedicated to difficult airway.

Despite its relatively poor image quality (Supplemental Table 3), the McGrath Mac[®] was chosen by most operators as the most “user friendly” device providing the easiest intubation (Figure 4). These results are in line with the more frequent use of the McGrath Mac[®] device than others (n = 180 compared to n=132, 139 and 138 for other devices). Even if one device was available for each anesthesia department, and the anesthesiologists strongly encouraged to use the device as often as possible, it is likely that the more the Macintosh-VL was considered effective, the more it was used in this quality improvement project. As the operators were not experienced with the McGrath Mac, it is unlikely that the more frequent use of the McGrath MAC results from greater familiarity. By contrast the C-MAC-S[®] delivered the best quality image but was not deemed the most user friendly device. As in previous studies,(26) image quality does not seem to affect the proportion of successful attempts. However, the teaching quality of the VL, making it possible for a supervisor to observe the trainee’s actions during direct-laryngoscopy and give advice,(27) may be more strongly associated with the quality of image and warrants further studies. Moreover, the difficulties assessed by the user-friendliness NRS may have resulted from insufficient familiarity with or training on the devices used.

Finally, it is worth noting that higher BMI was significantly associated with more use of a hyperangulated blade. These results are consistent with the findings of recent studies showing higher odds of difficult intubation in case of higher BMI.(1,28)

The aim of this quality improvement project was in line with recent guidelines.(6,29) All anesthesiologists should be trained to use, and have immediate access to, a videolaryngoscope. Proficiency with videolaryngoscope intubation is unlikely to be achieved if there are several different devices across one hospital.(30–32) Developing expertise requires frequent rather than exceptional use,(2) in operating room(6) as in ICU(29,33,34) or emergency settings.(35) Cook et al.(18) reported the conversion to VL as a routine first-choice option throughout a hospital’s anesthetic and ICU practice. Consistent with this study, the results observed in the current study

suggest that a single device to be used for easy and difficult intubation can be successfully implemented in operating room. One could suggest that these devices could even replace the standard-Macintosh laryngoscope in operating rooms.(18) The use of a single effective device for easy and difficult intubation may decrease the number of intubation attempts and by extension the complications related to the number of intubation attempts.(36,37) An abundant literature shows that delayed, difficult, or failed intubation is a major risk factor of harm and death in the patient with difficult airway.(16,38) However, although the McGrath MAC may have outperformed the other Macintosh-VL devices, none performed well enough to be advocated as a "universal" method, with a failure rate of 125/589 (21%) for the proposed universal VL. An alternative strategy would be to retain Macintosh laryngoscope and resort to a hyperangulated VL as a routine or rescue technique. There is evidence, though no prospective study, that the latter strategy is highly effective.(36) Indeed, regular use of a hyperangulated blade would increase familiarity, providing improved laryngeal visualization and fewer conversions to alternative techniques.

There are some limitations to discuss. First this is a non-blinded non-randomized quality improvement project, which weakens the conclusions. Providers had a choice as to whom to include in the study, so the trial was not purely randomized, which is a major confounding factor. However, the order of testing each Macintosh-VL in each center was randomly selected and the data were prospectively collected. Second, operator experience with the use of VL varied from new users to experienced users. However, as the groups did not differ regarding operator characteristics, it is unlikely that the comparison between groups could have been biased. As the name of the operator was not systematically recorded, we were not able to assess the improvement of their performance. Third, concerns have been raised, about the use of the Cormack–Lehane grading system during videolaryngoscopy. Although many studies have demonstrated that videolaryngoscopy improves the view, this does not always translate into easier tracheal intubation. Alternatives like the intubation difficulty scale(39) could have been

used to capture indices of difficulty common to both direct and indirect laryngoscopes. Fourth, since step III was a conditional step depending on the results of step II, results for step III should be interpreted descriptively and with caution. Fifth, the study was powered for a very large effect (10% versus 30% with the primary outcome). Power was moderate/low for smaller differences that would still be clinically important for the binary outcomes. Negative results on binary outcomes cannot be taken as definitive. Sixth, even after further adjusting for confounding as recommended, there is a potential for residual confounding bias which is a main concern in this study. The patients positioning was not standardized, whereas positioning impacts laryngoscopy (and VL) view. The use of head repositioning and Backward Upward Rightward Pressure/ External Laryngeal Pressure have not been recorded and may have resulted in improved laryngeal exposure with a reduced need to convert to a hyperangulated blade. Thyromental distance < 6 cm appeared more frequently in the patients in whom the CMAC- S was used, introducing the possibility of bias. However, the thyromental distance < 6 cm was not significant in the final multivariable model. Moreover, the presence of at least one factor for difficult intubation did not differ between groups. Seventh, the CL score does not reflect the ease or difficulty of tracheal intubation. However, the CL score provides a reasonable description of the quality of laryngoscopy (for both direct and indirect methods) but does not equate with the ease of intubation (by either direct or indirect methods).

CONCLUSION

In conclusion, the glottic visualization in direct and indirect-laryngoscopy with the Macintosh-style blade was significantly improved with the McGrath Mac[®] compared to other Macintosh-VLs, leading to a decreased resort to the hyperangulated blade and overall number of intubation attempts. Among VLs, MacGrath-Mac may be the preferred VL and eventually become the standard VL.

REFERENCES

1. De Jong A, Molinari N, Pouzeratte Y, Verzilli D, Chanques G, Jung B, Futier E, Perrigault PF, Colson P, Capdevila X, Jaber S. Difficult intubation in obese patients: incidence, risk factors, and complications in the operating theatre and in intensive care units. *Br J Anaesth.* 2015;114:297–306.
2. Cook TM, Kelly FE. A national survey of videolaryngoscopy in the United Kingdom. *Br J Anaesth.* 2017;118:593–600.
3. Holmes MG, Dagal A, Feinstein BA, Joffe AM. Airway Management Practice in Adults With an Unstable Cervical Spine: The Harborview Medical Center Experience. *Anesth Analg.* 2018;127:450–4.
4. Nam K, Lee Y, Park HP, Chung J, Yoon HK, Kim TK. Cervical Spine Motion During Tracheal Intubation Using an Optiscope Versus the McGrath Videolaryngoscope in Patients With Simulated Cervical Immobilization: A Prospective Randomized Crossover Study. *Anesth Analg.* 2019;129:1666–72.
5. Pieters BM, Theunissen M, van Zundert AA. Macintosh Blade Videolaryngoscopy Combined With Rigid Bonfils Intubation Endoscope Offers a Suitable Alternative for Patients With Difficult Airways. *Anesth Analg.* 2018;126:988–94.
6. Frerk C, Mitchell VS, McNarry AF, Mendonca C, Bhagrath R, Patel A, O’Sullivan EP, Woodall NM, Ahmad I. Difficult Airway Society 2015 guidelines for management of unanticipated difficult intubation in adults. *Br J Anaesth.* 2015;115:827–48.
7. Apfelbaum JL, Hagberg CA, Caplan RA, Blitt CD, Connis RT, Nickinovich DG, Benumof JL, Berry FA, Bode RH, Cheney FW, Guidry OF, Ovassapian A. Practice Guidelines for Management of the Difficult Airway: An Updated Report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology.* 2013;118:251–70.
8. Langeron O, Bourgain JL, Francon D, Amour J, Baillard C, Bouroche G, Chollet Rivier M, Lenfant F, Plaud B, Schoettker P, Fletcher D, Velly L, Nouette-Gaulain K. Difficult intubation and extubation in adult anaesthesia. *Anaesth Crit Care Pain Med.* 2018;37:639–51.
9. Herbstreich F, Fassbender P, Haberl H, Kehren C, Peters J. Learning endotracheal intubation using a novel videolaryngoscope improves intubation skills of medical students. *Anesth Analg.* 2011;113:586–90.
10. Meitzen SE, Benumof JL. Video Laryngoscopy: Positives, Negatives, and Defining the Difficult Intubation. *Anesth Analg.* 2019;128:399–401.
11. Schroeder RA, Pollard R, Dhakal I, Cooter M, Aronson S, Grichnik K, Buhrman W, Kertai MD, Mathew JP, Stafford-Smith M. Temporal Trends in Difficult and Failed Tracheal Intubation in a Regional Community Anesthetic Practice. *Anesthesiology.* 2018;128:502–10.
12. Cook F, Lobo D, Martin M, Imbert N, Grati H, Daami N, Cherait C, Saidi NE, Abbay K, Jaubert J, Younsi K, Bensaid S, Ait-Mamar B, Slavov V, Mounier R, Goater P, Bloc S,

- Catineau J, Abdelhafidh K, Haouache H, Dhonneur G. Prospective validation of a new airway management algorithm and predictive features of intubation difficulty. *Br J Anaesth.* 2019;122:245-54.
13. Sainsbury JE, Telgarsky B, Parotto M, Niazi A, Wong DT, Cooper RM. The effect of verbal and video feedback on learning direct laryngoscopy among novice laryngoscopists: a randomized pilot study. *Can J Anaesth.* 2017;64:252–9.
 14. De Jong A, Clavieras N, Conseil M, Coisel Y, Moury PH, Pouzeratte Y, Cisse M, Belafia F, Jung B, Chanques G, Molinari N, Jaber S. Implementation of a combo videolaryngoscope for intubation in critically ill patients: a before-after comparative study. *Intensive Care Med.* 2014;39:2144–52.
 15. De Jong A, Molinari N, Conseil M, Coisel Y, Pouzeratte Y, Belafia F, Jung B, Chanques G, Jaber S. Video laryngoscopy versus direct laryngoscopy for orotracheal intubation in the intensive care unit: a systematic review and meta-analysis. *Intensive Care Med.* 2014;40:629–39.
 16. De Jong A, Jung B, Jaber S. Intubation in the ICU: we could improve our practice. *Crit Care.* 2014;18:209.
 17. Berkow LC, Morey TE, Urdaneta F. The Technology of Video Laryngoscopy. *Anesth Analg.* 2018;126:1527-34.
 18. Cook TM, Boniface NJ, Seller C, Hughes J, Damen C, MacDonald L, Kelly FE. Universal videolaryngoscopy: a structured approach to conversion to videolaryngoscopy for all intubations in an anaesthetic and intensive care department. *Br J Anaesth.* 2018;120:173-80.
 19. Aziz MF, Abrons RO, Cattano D, Bayman EO, Swanson DE, Hagberg CA, Todd MM, Brambrink AM. First-Attempt Intubation Success of Video Laryngoscopy in Patients with Anticipated Difficult Direct Laryngoscopy: A Multicenter Randomized Controlled Trial Comparing the C-MAC D-Blade Versus the GlideScope in a Mixed Provider and Diverse Patient Population. *Anesth Analg.* 2016;122:740–50.
 20. Aziz MF, Healy D, Kheterpal S, Fu RF, Dillman D, Brambrink AM. Routine clinical practice effectiveness of the Glidescope in difficult airway management: an analysis of 2,004 Glidescope intubations, complications, and failures from two institutions. *Anesthesiology.* 2011;114:34-41.
 21. Toulouse E, Masseguin C, Lafont B, McGurk G, Harbonn A, J AR, Granier S, Dupeyron A, Bazin JE. French legal approach to clinical research. *Anaesth Crit Care Pain Med.* 2018;37:607-14.
 22. Min SH, Im H, Kim BR, Yoon S, Bahk JH, Seo JH. Randomized Trial Comparing Early and Late Administration of Rocuronium Before and After Checking Mask Ventilation in Patients With Normal Airways. *Anesth Analg.* 2019;
 23. Cormack RS, Lehane J. Difficult tracheal intubation in obstetrics. *Anaesthesia.* 1984;39:1105–11.
 24. Ruetzler K, Guzzella SE, Tscholl DW, Restin T, Cribari M, Turan A, You J, Sessler DI, Seifert B, Gaszynski T, Ganter MT, Spahn DR. Blind Intubation through Self-pressurized,

Disposable Supraglottic Airway Laryngeal Intubation Masks: An International, Multicenter, Prospective Cohort Study. *Anesthesiology*. 2017;127:307–16.

25. Kleine-Brueggemey M GR, Schoettker P, Savoldelli GL, Nabecker S, Theiler LG. Evaluation of six videolaryngoscopes in 720 patients with a simulated difficult airway: a multicentre randomized controlled trial. *Br J Anaesth*. 2016;116:670–9.
26. Shin M, Bai SJ, Lee KY, Oh E, Kim HJ. Comparing McGrath(R) MAC, C-MAC(R), and Macintosh Laryngoscopes Operated by Medical Students: A Randomized, Crossover, Manikin Study. *Biomed Res Int*. 2016;8943931.
27. Kelly FE, Cook TM, Boniface N, Hughes J, Seller C, Simpson T. Videolaryngoscopes confer benefits in human factors in addition to technical skills. *Br J Anaesth*. 2015;115:132–3.
28. Saasouh W, Laffey K, Turan A, Avitsian R, Zura A, You J, Zimmerman NM, Szarpak L, Sessler DI, Ruetzler K. Degree of obesity is not associated with more than one intubation attempt: a large centre experience. *Br J Anaesth*. 2018;120:1110–6.
29. Higgs A, McGrath BA, Goddard C, Rangasami J, Suntharalingam G, Gale R, Cook TM. Guidelines for the management of tracheal intubation in critically ill adults. *Br J Anaesth*. 2018;120:323–52.
30. McNarry AF, Patel A. The evolution of airway management - new concepts and conflicts with traditional practice. *Br J Anaesth*. 2017;119:i154-i66.
31. Cook TM, Woodall N, Frerk C. A national survey of the impact of NAP4 on airway management practice in United Kingdom hospitals: closing the safety gap in anaesthesia, intensive care and the emergency department. *Br J Anaesth*. 2016;117:182–90.
32. Lewis SR, Butler AR, Parker J, Cook TM, Schofield-Robinson OJ, Smith AF. Videolaryngoscopy versus direct laryngoscopy for adult patients requiring tracheal intubation: a Cochrane Systematic Review. *Br J Anaesth*. 2017;119:369–83.
33. Jaber S, Bellani G, Blanch L, Demoule A, Esteban A, Gattinoni L, Guérin C, Hill N, Laffey JG, Maggiore SM, Mancebo J, Mayo PH, Mosier JM, Navalese P, Quintel M, Vincent JL, Marini JJ. The intensive care medicine research agenda for airways, invasive and noninvasive mechanical ventilation. *Intensive Care Med*. 2017;43:1352–65.
34. Quintard H, l'Her E, Pottecher J, Adnet F, Constantin JM, De Jong A, Diemunsch P, Fesseau R, Freynet A, Girault C, Guitton C, Hamonic Y, Maury E, Mekontso-Dessap A, Michel F, Nolent P, Perbet S, Prat G, Roquilly A, Tazarourte K, Terzi N, Thille AW, Alves M, Gayat E, Donetti L. Intubation and extubation of the ICU patient. *Anaesth Crit Care Pain Med*. 2017;36:327–41.
35. Gellerfors M, Fevang E, Backman A, Kruger A, Mikkelsen S, Nurmi J, Rognas L, Sandstrom E, Skallsjo G, Svensen C, Gryth D, Lossius HM. Pre-hospital advanced airway management by anaesthetist and nurse anaesthetist critical care teams: a prospective observational study of 2028 pre-hospital tracheal intubations. *Br J Anaesth*. 2018;120:1103–9.
36. Aziz MF, Brambrink AM, Healy DW, Willett AW, Shanks A, Tremper T, Jameson L, Ragheb J, Biggs DA, Paganelli WC, Rao J, Epps JL, Colquhoun DA, Bakke P, Kheterpal S.

Success of Intubation Rescue Techniques after Failed Direct Laryngoscopy in Adults: A Retrospective Comparative Analysis from the Multicenter Perioperative Outcomes Group. *Anesthesiology*. 2016;125(125):656-66.

37. McNarry AF, Patel A. The evolution of airway management — new concepts and conflicts with traditional practice. *Br J Anaesth*. 2017;129:i154-i66.
38. Cook TM, Woodall N, Frerk C. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 1: anaesthesia. *Br J Anaesth*. 2011;106:617-31.
39. Adnet F, Borron SW, Racine SX, Clemessy JL, Fournier JL, Plaisance P, Lapandry C. The intubation difficulty scale (IDS): proposal and evaluation of a new score characterizing the complexity of endotracheal intubation. *Anesthesiology*. 1997;87:1290-7.

FIGURE LEGENDS

Figure 1. Methodology of the quality improvement project

Three steps were performed:

- **Step I:** After preoxygenation for 3 minutes according to the institution protocol and anesthesia induction, the first step was to rate the glottic visualization, using the Macintosh-VLs in direct-laryngoscopy with the Macintosh-style blade.
- **Step II:** The second step was to rate the glottic visualization (third rating) with the Macintosh-VLs in indirect-laryngoscopy (using the Macintosh-style blade); If the CL grade was \leq II, the operator made an intubation attempt, if the CL grade was \geq III, the operator moved to the third step without attempting intubation.
- **Step III:** The third step was performed only in case of CL grade \geq III (III or IV) or if the intubation attempt at the second step failed. This step was mandatorily performed by a senior anesthesiologist. The Macintosh-style blade was switched to the hyperangulated blade, using the same device. After rating the new glottic visualization (fourth rating) using the hyperangulated blade, the operator made an intubation attempt.

An intubation attempt was defined as the insertion of the laryngoscope blade into the mouth of the patient and the attempt to insert a tracheal tube.

The choice of an alternative technique after two intubation attempts was recommended by the study protocol and was in accordance with clinical standard.^{7,8,24} The alternative technique was left at the appreciation of the physician.

VL = videolaryngoscope

Figure 2. Flow chart of the study

Among 1114 patients assessed for eligibility, 589 patients were included, 180 in the McGrath Mac[©] group, 132 in the C-MAC-S[©] group, 139 in the C-MAC S PM[©] group and 138 in the APATM[©] group. The use of a hyperangulated blade was required in 22/180 (12%) in the McGrath Mac[©] group, 39/132 (30%) in the C-MAC-S[©] group, 29/139 (21%) in the C-MAC-S-PM[©] group and 35/138 (25%) in the APATM[©] group.

Figure 3. Main outcome (need of a hyperangulated blade) for each Macintosh videolaryngoscope after multivariable analysis

The hyperangulated blade was used in 22/180 patients (12%) for the McGrath Mac[©], in 29/139 patients (21%) for the C-MAC-S-PM[©], in 39/132 patients (30%) for the C-MAC-S[©] and in 35/138 patients (25%) for the APATM[©]. By multivariable analysis, the following variables were entered in the model: age, body mass index (BMI), center, thyromental distance < 6 cm, no cervical spine extension. After adjustment for BMI, the McGrath Mac[©] was associated with less frequent use of hyperangulated blade than C-MAC-S[©] (OR, 99.2% CI of 0.34 (0.16, 0.77), p=0.0005), or APATM[©] (OR, 99.2% CI of 0.42 (0.19, 0.93), p=0.004), but not C-MAC-S-PM[©] OR, 99.2% CI of 0.53 (0.23, 1.2), p=0.04, Figure 3).

Figure 4. Subjective evaluation for each Macintosh videolaryngoscope

The subjective appreciation of difficulty of intubation and user friendliness of devices respectively showed lower and higher numeric rating scale (NRS) scores for the McGrath Mac[©] device compared to the other devices.

* Significantly differ from McGrath Mac[©]

FIGURE 1. Methodology of the quality improvement project




FIGURE 2. Flow chart of the study




FIGURE 3. Main outcome (need of a hyperangulated blade) for each Macintosh videolaryngoscope after multivariable analysis




FIGURE 4. Subjective evaluation for each Macintosh videolaryngoscope




TABLE 1. Patients; airway and operator characteristics in the four Macintosh videolaryngoscopes groups

Characteristics	McGrath Mac [©]	C-MAC-S [©]	C-MAC-S-PM [©]	APA TM [©]	P Values	Data are summarized as mean ± SD or n (%). NA = Not Available. * C-MAC-S [©] ≠ McGrath Mac [©] . ** C-MAC S PM [©] ≠ McGrath Mac [©] . *** APA TM [©] ≠ McGrath Mac [©] .
General characteristics	(n = 180)	(n = 132)	(n = 139)	(n = 138)		
Age (years)	56 ± 18	55 ± 18	58 ± 18	56 ± 18	0.58	
Weight (kg)	74 ± 21	79 ± 19	76 ± 17	75 ± 19	0.08	
Height (cm)	168 ± 10	168 ± 10	169 ± 9	169 ± 10	0.68	
BMI (kg/m ²)	26 ± 7	28 ± 7	26 ± 6	26 ± 6	0.24	
Airway assessment						
Thyromental distance < 6cm	8/180 (4%)	5/128 (4%)	20/138 (14%)* ^{**} \$	18/138 (13%)* ^{***} \$\$	0.0008	
Inter-incisor distance < 3cm	21/180 (12%)	18/128 (14%)	21/138 (15%)	15/138 (11%)	0.67	
Limited cervical spine extension	21/180 (12%)	19/128 (15%)	19/138 (14%)	23/138 (17%)	0.63	
No cervical spine extension	7/180 (4%)	2/128 (2%)	0/138 (0%)	1/138 (1%)	0.04	
Mallampati score					0.52	
Mallampati score 1	85/177 (48%)	52/125 (22%)	73/135 (22%)	63/134 (47%)		APA TM
Mallampati score 2	45/177 (25%)	44/125 (42%)	32/135 (54%)	43/134 (32%)		© ≠ McGrath Mac [©]
Mallampati score 3	33/177 (19%)	22/125 (18%)	23/135 (17%)	21/134 (16%)		h Mac [©]
Mallampati score 4	14/177 (8%)	7/125 (6%)	7/135 (5%)	7/134 (5%)		
Previous difficult intubation	23/179 (13%)	11/125 (9%)	14/138 (10%)	9/133 (7%)		
At least one predictive criteria of difficult intubation	73/180 (41%)	54/128 (41%)	55/138 (40%)	43/138 (31%)	0.28	
Operator status						
Physician (senior anesthesiologist)	51/176 (29%)	49/129 (38%)	43/135 (32%)	47/136 (35%)	0.40	
Resident	30/176 (17%)	26/129 (20%)	33/135 (24%)	26/136 (19%)	0.44	
Anesthetist nurse	95/176 (54%)	54/129 (42%)	59/135 (44%)	63/136 (46%)	0.14	

TABLE 2. Outcomes according to step I, II and III in the four Macintosh videolaryngoscope groups

Outcomes	McGrath Mac [®] N = 180	C-MAC S [®] N = 132	C-MAC S PM [®] N = 139	APA TM [®] N = 138	P Values
CL score at step I (Videolaryngoscope, DL, Macintosh-style blade)					
1	104/180 (58%)	50/132 (38%)*	72/139 (52%)	61/138 (44%)	0.003
2	39/180 (22%)	31/132 (24%)	29/139 (21%)	37/138 (27%)	0.64
3	28/180 (15%)	40/132 (30%)	29/139 (21%)	26/138 (19%)	0.01
4	9/180 (5%)	11/132 (8%)	9/139 (6%)	14/138 (10%)	0.33
CL score 3 or 4	37/180 (20%)	51/132 (38%)*	38/139 (27%)	40/138 (29%)	0.006
Mean CL score at step I (Macintosh-style blade)	1.68 ± 0.91	2.10 ± 1.01*	1.82 ± 0.98	1.95 ± 1.02	0.001
CL score at step II (Videolaryngoscope, IL, Macintosh-style blade)					
1	146/180 (81%)	67/132 (51%)*	89/139 (64%)**	80/138 (58%)****	<0.0001
2	20/180 (11%)	32/132 (24%)*	32/139 (23%)	33/138 (24%)****	0.006
3	12/180 (7%)	27/132 (20%)*	12/139 (9%)	16/138 (12%)	0.002
4	2/180 (1%)	6/132 (5%)	6/139 (4%)	9/138 (6%)	0.09
CL score 3 or 4	14/180 (8%)	33/132 (25%)*	18/139 (13%)	25/138 (18%)****	0.0002
Mean CL score at step II	1.28 ± 0.64	1.79 ± 0.93*	1.54 ± 0.84**	1.67 ± 0.92***	<0.0001
Difference CL score Step I - Step II [¶]	0.40 ± 0.67	0.30 ± 0.68	0.28 ± 0.66	0.28 ± 0.63	0.20
Successful intubation at step II	158/166 (95%)	93/99 (94%)	110/121 (91%)	103/113 (91%)	0.43
Stylet use	56/159 (35%)	43/100 (43%)	46/104 (44%)	32/112 (29%)	0.06
Malleable stylet	30/49 (61%)	30/43 (70%)	20/37 (54%)	18/29 (62%)	0.67
Rigid stylet	18/49 (37%)	13/43 (30%)	13/37 (35%)	8/29 (28%)	0.81
Eschmann stylet	1/49 (2%)	0/43 (0%)	4/37 (11%)	3/29 (10%)	0.06

Step III rate (primary outcome)	22/180 (12%)	39/132 (30%)*	29/139 (21%)	35/138 (25%)***	0.001
CL score at step III (<i>Videolaryngoscope, IL, hyperangulated blade and senior anesthetist as operator</i>)					
1	17/22 (77%)	23/39 (59%)	17/29 (59%)	19/35 (54%)	ND
2	0/22 (0%)	12/39 (31%)	5/29 (17%)	8/35 (23%)	ND
3	4/22 (18%)	4/39 (10%)	4/29 (14%)	6/35 (17%)	ND
4	1/22 (5%)	0/39 (0%)	3/29 (10%)	2/35 (6%)	ND
CL score 3 or 4	5/22 (23%)	4/39 (10%)	7/29 (24%)	8/35 (23%)	ND
Mean CL score at step III	1.50 ± 0.96	1.51 ± 0.68	1.76 ± 1.06	1.74 ± 0.95	ND
Difference CL score Step II - Step III	1.09 ± 0.97	1.46 ± 0.85	0.93 ± 0.92	1.03 ± 0.99	ND
Stylet use	18/20 (90%)	33/39 (85%)	17/22 (77%)	6/28 (21%)	ND
Malleable stylet	13/18 (72%)	15/27 (55%)	10/16 (62%)	4/5 (80%)	ND
Rigid stylet	5/18 (28%)	8/27 (30%)	3/16 (19%)	1/5 (20%)	ND
Eschmann stylet	0/18 (0%)	4/27 (15%)	3/16 (19%)	0/5 (0%)	ND
Successful intubation at step III	19/22 (86%)	36/39 (92%)	21/29 (72%)	24/35 (69%)	ND
Successful intubation at step II and III	177/180 (98%)	129/132 (98%)	131/139 (94%)	127/138 (92%)	0.02
Number of attempts	1.14 ± 0.42	1.32 ± 0.71*	1.31 ± 0.58**	1.27 ± 0.63	0.005

Data are summarized as mean ± SD or n (%). CL = Cormack-Lehane. ND = Not Done (as the sample size for each group here is conditional on requiring step III, no statistical testing was done). DL = Direct Laryngoscopy. IL = Indirect Laryngoscopy

* The difference between Cormack-Lehane score obtained using the videolaryngoscope in direct vision and the Cormack-Lehane score obtained using videolaryngoscope in indirect vision differs significantly from 0 in the overall population and in all the videolaryngoscopes ($P<0.008$).

** C-MAC-S© ≠ McGrath Mac© *** C-MAC S PM© ≠ McGrath Mac© **** APATM© ≠ McGrath Mac© \$ C-MAC S PM© ≠ C-MAC-S© §§ APATM© ≠ C-MAC-S© £ APATM© ≠ C-MAC S PM© ($P<0.008$)

SUPPLEMENTAL MATERIAL

Macintosh Videolaryngoscopes for intubation in the operating room: A comparative quality improvement project

SUPPLEMENTAL TABLE 1. Drugs characteristics in the four Macintosh videolaryngoscopes groups

	McGrath Mac [®]	C-MAC S [®]	C-MAC S PM [®]	APA TM [®]	P Values
Drugs at induction					
Propofol	146/174 (84%)	99/128 (77%)	103/136 (76%)	108/133 (81%)	0.28
Propofol posology (mg)	304 ± 153	303 ± 136	271 ± 134	271 ± 130	0.14
Ketamine	25/174 (14%)	20/128 (16%)	23/136 (17%)	20/132 (15%)	0.94
Ketamine posology (mg)	22 ± 7	23 ± 12	19 ± 8	18 ± 7	0.17
Thiopentone	1/174 (1%)	0/128 (0%)	0/136 (0%)	0/132 (0%)	1.00
Thiopentone posology (mg)	NA	0	0	0	-
Etomidate	10/174 (6%)	3/128 (2%)	5/136 (4%)	2/132 (2%)	0.23
Etomidate posology (mg)	26 ± 9	20 ± 0	28 ± 9	20 ± 0	0.55
Sufentanil	132/174 (76%)	78/128 (61%)*	93/136 (68%)	95/132 (72%)	0.04
Sufentanil posology (mcg)	14 ± 4	15 ± 5	16 ± 7	15 ± 5	0.12
Remifentanil	2/174 (1%)	7/128 (5%)	6/136 (4%)	8/132 (6%)	0.12
Remifentanil posology (mcg)	70 ± 0	144 ± 96	67 ± 34	110 ± 17	0.41
Cisatracurium	85/174 (49%)	79/128 (62%)	59/136 (43%)	62/132 (47%)	0.02
Cisatracurium posology (mg)	13 ± 3	13 ± 3	12 ± 5	12 ± 3	0.21
Atracurium	5/174 (3%)	3/128 (2%)	5/136 (4%)	4/135 (3%)	0.95
Atracurium posology (mg)	31 ± 6	40 ± 5	35 ± 10	34 ± 11	0.66
Succinylcholine	0/174 (0%)	2/128 (2%)	0/136 (0%)	0/132 (0%)	0.05
Succinylcholine posology (mg)	0	53 ± 66	0	0	-
Rocuronium	0/174 (0%)	0/128 (0%)	1/136 (1%)	0/132 (0%)	0.95
Rocuronium posology (mg)	0	0	50 ± 0	0	-
Midazolam	0/174 (0%)	3/128 (2%)	3/136 (2%)	4/132 (3%)	0.08
Midazolam posology (mg)	0	1 ± 0	1 ± 1	8 ± 8	0.14
Propofol TIVA	24/174 (14%)	29/128 (23%)	31/136 (23%)	24/132 (18%)	0.14
Sufentanil TIVA	1/174 (1%)	15/128 (12%)*	4/136 (3%) ^{\$}	2/132 (2%) ^{\$\$}	<0.0001
Remifentanil TIVA	31/174 (18%)	23/128 (18%)	29/136 (21%)	24/132 (18%)	0.86

Data are summarized as mean ± SD or n (%). NA = Not Available. TIVA = Total intravenous anesthesia

* C-MAC ≠ MacGrath ** C-MAC pocket ≠ MacGrath *** APA ≠ MacGrath

SUPPLEMENTAL TABLE 2. Glottis visualization and alternative devices used in patients who reached step 3 in the four Macintosh videolaryngoscopes groups

	McGrath Mac [®]	C-MAC S [®]	C-MAC S PM [®]	APA™ [®]	P Values
CL score at step 2 (<i>Videolaryngoscope, Macintosh blade</i>)					
1	3/22 (14%)	1/39 (3%)	4/29 (14%)	7/35 (20%)	0.09
2	5/22 (23%)	5/39 (13%)	7/29 (24%)	3/35 (8%)	0.26
3	12/22 (54%)	27/39 (69%)	12/29 (41%)	16/35 (46%)	0.09
4	2/22 (9%)	6/39 (15%)	6/29 (21%)	9/35 (26%)	0.44
Mean CL score at step 2	2.59 ± 0.85	2.97 ± 0.63	2.69 ± 0.97	2.77 ± 1.06	0.41
Macintosh blade size					
3 (n)	27/137 (20%)	17/92 (18%)	18/93 (19%)	20/101 (20%)	0.7670
4 (n)	110/137 (80%)	75/92 (82%)	75/93 (81%)	81/101 (80%)	0.7670
CL score at step 3 of patients who were CL score 3 or 4 at step 2 (<i>Videolaryngoscope, hyperangulated blade and senior anesthetist as operator</i>)					
1	9/14 (64%)	18/33 (55%)	9/18 (50%)	11/25 (44%)	ND
2	0/14 (0%)	11/33 (33%)	2/18 (11%)	6/25 (24%)	ND
3	4/14 (29%)	4/33 (12%)	4/18 (22%)	6/25 (24%)	ND
4	1/14 (7%)	0/33 (0%)	3/18 (17%)	2/25 (8%)	ND
Mean CL score	1.79 ± 1.12	1.58 ± 0.71	2.06 ± 1.21	1.96 ± 1.02	ND
Difference CL score Step 2 vs Step 3 ^μ	1.36 ± 1.08	1.61 ± 0.83	1.28 ± 0.96	1.40 ± 0.91	ND

CL score at **step 3** of patients who were CL score 1 or 2 at **step 2** (*Videolaryngoscope, hyperangulated blade and senior anesthetist as operator*)

1	8/8 (100%)	5/6 (83%)	8/11 (73%)	8/10 (80%)	ND
2	0/8 (0%)	1/6 (17%)	3/11 (27%)	2/10 (20%)	ND
3	0/8 (0%)	0/6 (0%)	0/11 (0%)	0/10 (0%)	ND
4	0/8 (0%)	0/6 (0%)	0/11 (0%)	0/10 (0%)	ND
Mean CL score	1.00 ± 0.00	1.17 ± 0.41	1.27 ± 0.47	1.20 ± 0.42	ND
Difference CL score Step 2 - Step 3 ^{¶¶}	0.63 ± 0.52	0.67 ± 0.52	0.36 ± 0.50	0.10 ± 0.32	ND
Failure of intubation at Step III	3/22 (14%)	3/39 (8%)	8/29 (28%)	11/35 (31%)	ND
Alternatives techniques used in case of failure at Step III					
Standard Macintosh Laryngoscope		1/3 (33%)	5/8 (63%)	5/11 (45%)	-
Other VL dedicated to difficult intubation	3/3 (100%)	1/3 (33%)		6/11 (55%)	-
Same VL used in DL		1/3 (34%)			-
Straight blade			1/8 (12%)		-
Spontaneous ventilation without intubation			1/8 (12%)		-
Laryngeal mask			1/8 (13%)		-

Data are summarized as mean ± SD or n (%).

NA = Not Available. CL = Cormack-Lehane. ND = Not Done (as the sample size for each group here is conditional on requiring step III, no statistical testing was done).

[¶]The difference between Cormack-Lehane score obtained using the videolaryngoscope in indirect vision with the Macintosh blade and the Cormack-Lehane score obtained using videolaryngoscope in indirect vision with the hyperangulated blade differs significantly from 0 in the overall population and in all the videolaryngoscopes.

^{¶¶}The difference between Cormack-Lehane score obtained using the videolaryngoscope in indirect vision with the Macintosh blade and the Cormack-Lehane score obtained using videolaryngoscope in indirect vision with the hyperangulated blade did not differ significantly from 0.

* C-MAC ≠ MacGrath ** C-MAC pocket ≠ MacGrath *** APA ≠ MacGrath § C-MAC Pocket ≠ C-MAC §§ APA ≠ C-MAC

[£] APA ≠ C-MAC Pocket

SUPPLEMENTAL TABLE 3. Complications, subjective appreciation and quality of image in the four Macintosh videolaryngoscope groups

	McGrath Mac [®]	C-MAC S [®]	C-MAC S PM [®]	APA TM [®]	P Values
Complications					
Hypoxemia	6/177 (3%)	14/130 (11%)*	6/134 (5%)	8/134 (6%)	0.0452
Bradycardia	1/177 (1%)	3/130 (2%)	4/135 (3%)	5/135 (4%)	0.2799
Hypotension	6/177 (3%)	5/129 (4%)	7/134 (5%)	7/135 (5%)	0.8184
At least one	12/176 (7%)	20/129 (16%)	15/132 (11%)	18/134 (13%)	0.0953
Subjective appreciation					
Easy	159/170 (94%)	96/123 (78%)*	104/127 (82%)**	109/129 (84%)	0.0013
Difficult	9/170 (5%)	23/123 (19%)*	15/127 (12%)	10/129 (8%)	0.0016
Impossible	2/170 (1%)	4/123 (3%)	8/127 (6%)	10/129 (8%)	0.0442
NRS difficulty of intubation	1 ± 2	3 ± 3*	3 ± 3**	3 ± 3***	< 0.0001
NRS ≤ 3	144/161 (89%)	80/124 (65%)*	85/126 (67%)**	89/127 (70%)***	< 0.0001
4 ≤ NRS ≤ 6	13/161 (8%)	34/124 (27%)*	25/126 (20%)**	19/127 (15%)	0.0002
7 ≤ NRS	4/161 (2%)	10/124 (8%)	16/126 (13%)**	19/127 (15%)***	0.0011
NRS conviviality for devices	9 ± 2	7 ± 2*	7 ± 3**	6 ± 3***	< 0.0001
NRS ≤ 3	5/163 (3%)	13/123 (11%)	16/128 (13%)**	24/129 (19%)***	0.0003
4 ≤ NRS ≤ 6	12/163 (7%)	36/123 (29%)*	27/128 (21%)**	36/129 (28%)***	< 0.0001
7 ≤ NRS	146/163 (90%)	74/123 (60%)*	85/128 (66%)**	69/129 (53%)***	< 0.0001
Quality of image					
Bad	20/175 (11%)	2/126 (2%)*	5/135 (4%)	19/133 (15%)	< 0.0001
Passable	59/175 (34%)	7/126 (5%)*	15/135 (11%)**	44/133 (33%)	< 0.0001
Good	54/175 (31%)	42/126 (33%)	52/135 (39%)	52/133 (39%)	0.3665
Very good	33/175 (19%)	59/126 (47%)*	45/135 (33%)**	15/133 (11%) ^{\$\$£}	< 0.0001
Excellent	9/175 (5%)	16/126 (13%)	18/135 (13%)**	3/133 (2%) ^{\$\$£}	0.0007

Data are summarized as mean ± SD or n (%).

NA = Not Available. NRS = Numeric Rating Scale

* C-MAC ≠ MacGrath ** C-MAC pocket ≠ MacGrath *** APA ≠ MacGrath \$ C-MAC Pocket ≠ C-MAC \$\$ APA ≠ C-MAC

£ APA ≠ C-MAC Pocket

SERMENT

- En présence des Maîtres de cette école, de mes chers condisciples et devant l'effigie d'Hippocrate, je promets et je jure, au nom de l'Etre suprême, d'être fidèle aux lois de l'honneur et de la probité dans l'exercice de la médecine.
- Je donnerai mes soins gratuits à l'indigent et n'exigerai jamais un salaire au-dessus de mon travail.
- Admis (e) dans l'intérieur des maisons, mes yeux ne verront pas ce qui s'y passe, ma langue taira les secrets qui me seront confiés, et mon état ne servira pas à corrompre les mœurs, ni à favoriser le crime.
- Respectueux (se) et reconnaissant (e) envers mes Maîtres, je rendrai à leurs enfants l'instruction que j'ai reçue de leurs pères.
- Que les hommes m'accordent leur estime si je suis fidèle à mes promesses.
Que je sois couvert (e) d'opprobre et méprisé (e) de mes confrères si j'y manque.

RÉSUMÉ

Introduction : Les Vidéolaryngoscopes (VLs) “Macintosh” sont des dispositifs utilisés pour l’intubation permettant de réaliser à la fois une laryngoscopie directe et indirecte. Dans le cadre d’une évaluation prospective multicentrique au CHRU de Montpellier, nous avons comparé les performances de 4 VLs Macintosh en vision directe et indirecte.

Méthodes: Il s’agit d’un projet-qualité visant à généraliser au bloc opératoire l’utilisation des VLs Macintosh pour toutes les intubations afin d’améliorer la gestion des voies aériennes. Les 4 VLs Macintosh testés étaient : le McGrath-Mac[®], le C-MAC-S[®], le C-MAC-S-Pocket-Monitor[®] (PM) et l’APA™[®]. L’évaluation se déroulait en 3 étapes consécutives : I) une laryngoscopie directe avec le VL Macintosh, II) une laryngoscopie indirecte avec le VL Macintosh (en cas de Cormack I ou II, une tentative d’intubation avait lieu), III) Tentative d’intubation en ayant recours à la lame hyperangulée, toujours en vision indirecte, en cas de Cormack III/IV ou échec à l’étape précédente. Le critère de jugement principal était le besoin de recourir à la lame hyperangulée (étape III). Une analyse multivariée avec régression logistique a été réalisée, une valeur de p-value<0.008 était considérée comme statistiquement significative.

Résultats : De Mai à Septembre 2017, 589 patients ont été inclus. L’utilisation du McGrath-Mac[®] (22/180 (12%)) était associée à moins de recours à la lame hyperangulée que l’utilisation du C-MAC-S[®] (39/132 (30%), odds ratio (OR) = 0.34 avec un intervalle de confiance (IC) à 99.2% [0.16-0.77], p=0.005 et de l’APA[®] (35/138 (25%), OR=0.42, IC 99.2% [0.19-0.93], p=0.004, contrairement à l’utilisation du C-MAC-S-PM[®] (29/139 (21%), OR=0.53, IC99.2% [0.23-1.2], p=0.04). Globalement le nombre de tentatives d’intubation était significativement moins important en utilisant le McGrath-Mac[®] que les autres vidéolaryngoscopes. Le McGrath-Mac[®] obtenait également les meilleures appréciations subjectives en termes de facilité d’intubation et de convivialité-ergonomie. Cependant l’évaluation subjective concernant la qualité de l’image montrait des scores sur l’échelle visuelle numérique meilleurs pour le C-MAC-S[®] et le C-MAC-S-PM[®] que ceux du McGrath Mac[®] et de l’APA[®].

Conclusion: Parmi les quatre VLs Macintosh à usage unique testés, le McGrath Mac[®] améliorait de manière significative la visualisation directe et indirecte de la glotte, réduisant à la fois le recours à la lame hyperangulée et le nombre de tentatives d’intubation.

Mots clés : Voies aériennes, Vidéolaryngoscope Macintosh, Intubation, Bloc opératoire, Vidéolaryngoscopes