



**HAL**  
open science

# Does rifampin improve prosthetic staphylococcal endocarditis' prognosis? Results of a multicenter retrospective study

Audrey Le Bot

► **To cite this version:**

Audrey Le Bot. Does rifampin improve prosthetic staphylococcal endocarditis' prognosis? Results of a multicenter retrospective study. Life Sciences [q-bio]. 2020. dumas-03241294

**HAL Id: dumas-03241294**

**<https://dumas.ccsd.cnrs.fr/dumas-03241294>**

Submitted on 28 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## THESE D'EXERCICE / UNIVERSITE DE RENNES 1

*sous le sceau de l'Université Bretagne Loire*

Thèse en vue du

### DIPLÔME D'ETAT DE DOCTEUR EN MEDECINE

présentée par

**Audrey LE BOT**

Née le 02 novembre 1990 à Quimper (29)

---

**La rifampicine améliore-t-elle le pronostic des endocardites à *Staphylococcus* sp. sur prothèse ?  
Une étude rétrospective multicentrique**

***Does rifampin improve prosthetic staphylococcal endocarditis' prognosis ?  
Results of a multicenter retrospective study.***

Thèse soutenue à Rennes le 28/09/2020

devant le jury composé de :

**Pierre TATTEVIN**

PU-PH, CHU de Rennes / *Président*

**Yves LE TULZO**

PU-PH, CHU de Rennes / *Examinateur*

**David BOUTOILLE**

PU-PH, CHU de Nantes / *Examinateur*

**Séverine ANSART**

PU-PH, CHRU de Brest / *Examinatrice*

**Rozenn LE BERRE**

MCU-PH, CHRU de Brest / *Examinatrice*

**Matthieu REVEST**

PU-PH, CHU de Rennes / *Directeur de thèse*



## THESE D'EXERCICE / UNIVERSITE DE RENNES 1

*sous le sceau de l'Université Bretagne Loire*

Thèse en vue du

### DIPLÔME D'ETAT DE DOCTEUR EN MEDECINE

présentée par

**Audrey LE BOT**

Née le 02 novembre 1990 à Quimper (29)

---

**La rifampicine améliore-t-elle le pronostic des endocardites à *Staphylococcus* sp. sur prothèse ?  
Une étude rétrospective multicentrique**

***Does rifampin improve prosthetic staphylococcal endocarditis' prognosis ?  
Results of a multicenter retrospective study.***

Thèse soutenue à Rennes le 28/09/2020  
devant le jury composé de :

**Pierre TATTEVIN**  
PU-PH, CHU de Rennes / *Président*

**Yves LE TULZO**  
PU-PH, CHU de Rennes / *Examineur*

**David BOUTOILLE**  
PU-PH, CHU de Nantes / *Examineur*

**Séverine ANSART**  
PU-PH, CHRU de Brest / *Examinatrice*

**Rozenn LE BERRE**  
MCU-PH, CHRU de Brest / *Examinatrice*

**Matthieu REVEST**  
PU-PH, CHU de Rennes / *Directeur de thèse*

## LISTE DES PROFESSEURS DES UNIVERSITES

NOM	PRENOM	TITRE	CNU
ANNE-GALIBERT	Marie-Dominique	PU-PH	Biochimie et biologie moléculaire
BARDOU-JACQUET	Edouard	PU-PH	Gastroentérologie ; hépatologie ; addictologie
BELAUD-ROTUREAU	Marc-Antoine	PU-PH	Histologie ; embryologie et cytogénétique
BELLISSANT	Eric	PU-PH	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
BELOEIL	Hélène	PU-PH	Anesthésiologie-réanimation ; médecine d'urgence
BENDAVID	Claude	PU-PH	Biochimie et biologie moléculaire
BENSALAH	Karim	PU-PH	Urologie
BEUCHEE	Alain	PU-PH	Pédiatrie
BONAN	Isabelle	PU-PH	Médecine physique et de réadaptation
BONNET	Fabrice	PU-PH	Endocrinologie, diabète et maladies métaboliques ; gynécologie médicale
BOUDJEMA	Karim	PU-PH	Chirurgie générale
BOUGET	Jacques	PU Emérite	Thérapeutique-médecine d'urgence ; addictologie
BOUGUEN	Guillaume	PU-PH	Gastroentérologie ; hépatologie ; addictologie
BRASSIER	Gilles	PU-PH	Neurochirurgie
BRETAGNE	Jean-François	PU Emérite	Gastroentérologie ; hépatologie ; addictologie
BRISOT	Pierre	PU Emérite	Gastroentérologie ; hépatologie ; addictologie
CARRE	François	PU-PH	Physiologie
CATROS	Véronique	PU-PH	Biologie cellulaire
CATTOIR	Vincent	PU-PH	Bactériologie-virologie ; hygiène hospitalière
CHALES	Gérard	PU Emérite	Rhumatologie
CORBINEAU	Hervé	PU-PH	Chirurgie thoracique et cardiovasculaire
CUGGIA	Marc	PU-PH	Biostatistiques, informatique médicale et technologies de communication
DARNAULT	Pierre	PU-PH	Anatomie
DAUBERT	Jean-Claude	PU Emérite	Cardiologie
DAVID	Véronique	PU-PH	Biochimie et biologie moléculaire
DAYAN	Jacques	Pr Associé	Pédopsychiatrie ; Addictologie
DE CREVOISIER	Renaud	PU-PH	Cancérologie ; radiothérapie
DECAUX	Olivier	PU-PH	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
DESRUES	Benoît	PU-PH	Pneumologie ; addictologie
DEUGNIER	Yves	PU Emérite	Gastroentérologie ; hépatologie ; addictologie
DONAL	Erwan	PU-PH	Cardiologie
DRAPIER	Dominique	PU-PH	Psychiatrie d'adultes ; addictologie
DUPUY	Alain	PU-PH	Dermato-vénérologie
ECOFFEY	Claude	PU-PH	Anesthésiologie-réanimation ; médecine d'urgence
EDAN	Gilles	PU-PH consultant	Neurologie
FERRE	Jean-Christophe	PU-PH	Radiologie et imagerie médicale
FEST	Thierry	PU-PH	Hématologie ; transfusion
FLECHER	Erwan	PU-PH	Chirurgie thoracique et cardiovasculaire
GANDEMER	Virginie	PU-PH	Pédiatrie
GANDON	Yves	PU-PH	Radiologie et imagerie médicale
GANGNEUX	Jean-Pierre	PU-PH	Parasitologie et mycologie

GARIN	Etienne	PU-PH	Biophysique et médecine nucléaire
GAUVRIT	Jean-Yves	PU-PH	Radiologie et imagerie médicale
GODEY	Benoît	PU-PH	Oto-rhino-laryngologie
GUGGENBUHL	Pascal	PU-PH	Rhumatologie
GUILLE	François	PU-PH	Urologie
GUYADER	Dominique	PU-PH	Gastroentérologie ; hépatologie ; addictologie
HAEGELEN	Claire	PU-PH	Anatomie
HOUOT	Roch	PU-PH	Hématologie ; transfusion
HUSSON	Jean-Louis	PU Emérite	Chirurgie orthopédique et traumatologique
HUTEN	Denis	PU Emérite	Chirurgie orthopédique et traumatologique
JEGO	Patrick	PU-PH	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
JEGOUX	Franck	PU-PH	Oto-rhino-laryngologie
JOUNEAU	Stéphane	PU-PH	Pneumologie ; addictologie
KAYAL	Samer	PU-PH	Bactériologie-virologie ; hygiène hospitalière
LAMY DE LA CHAPELLE	Thierry	PU-PH	Hématologie ; transfusion
LAVIOLLE	Bruno	PU-PH	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
LAVOUE	Vincent	PU-PH	Gynécologie-obstétrique ; gynécologie médicale
LE BRETON	Hervé	PU-PH	Cardiologie
LE TULZO	Yves	PU-PH	Médecine intensive-réanimation
LECLERCQ	Christophe	PU-PH	Cardiologie
LEDERLIN	Mathieu	PU-PH	Radiologie et imagerie médicale
LEGUERRIER	Alain	PU Emérite	Chirurgie thoracique et cardiovasculaire
LEJEUNE	Florence	PU-PH	Biophysique et médecine nucléaire
LEVEQUE	Jean	PU-PH	Gynécologie-obstétrique ; gynécologie médicale
LIEVRE	Astrid	PU-PH	Gastroentérologie ; hépatologie ; addictologie
MABO	Philippe	PU-PH	Cardiologie
MAHE	Guillaume	PU-PH	Chirurgie vasculaire ; médecine vasculaire
MALLEDANT	Yannick	PU Emérite	Anesthésiologie-réanimation ; médecine d'urgence
MATHIEU-SANQUER	Romain	PU-PH	Urologie
MENER	Eric	Pr Associé	Médecine générale
MEUNIER	Bernard	PU-PH	Chirurgie digestive
MICHELET	Christian	PU Emérite	Maladies infectieuses ; maladies tropicales
MOIRAND	Romain	PU-PH	Gastroentérologie ; hépatologie ; addictologie
MORANDI	Xavier	PU-PH	Anatomie
MOREL	Vincent	Pr Associé	Médecine palliative
MOSSER	Jean	PU-PH	Biochimie et biologie moléculaire
MOURIAUX	Frédéric	PU-PH	Ophtalmologie
MYHIE	Didier	Pr Associé	Médecine générale
NAUDET	Florian	PU-PH	Thérapeutique-Médecine de la douleur ; addictologie
ODENT	Sylvie	PU-PH	Génétique
OGER	Emmanuel	PU-PH	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PARIS	Christophe	PU-PH	Médecine et santé au travail
PERDRIGER	Aleth	PU-PH	Rhumatologie
PLADYS	Patrick	PU-PH	Pédiatrie
RAVEL	Célia	PU-PH	Histologie, embryologie et cytogénétique
RENAUT	Pierric	Pr Associé	Médecine générale

REVEST	Matthieu	PU-PH	Maladies infectieuses ; maladies tropicales
RIFFAUD	Laurent	PU-PH	Neurochirurgie
RIOUX-LECLERCQ	Nathalie	PU-PH	Anatomie et cytologie pathologiques
ROBERT-GANGNEUX	Florence	PU-PH	Parasitologie et mycologie
ROPARS	Mickaël	PU-PH	Chirurgie orthopédique et traumatologique
SAINT-JAMES	Hervé	PU-PH	Biophysique et médecine nucléaire
SAULEAU	Paul	PU-PH	Physiologie
SEGUIN	Philippe	PU-PH	Anesthésiologie-réanimation ; médecine d'urgence
SEMANA	Gilbert	PU-PH	Immunologie
SIPROUDHIS	Laurent	PU-PH	Gastroentérologie ; hépatologie ; addictologie
SOMME	Dominique	PU-PH	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
SOULAT	Louis	Pr Associé	Médecine d'urgence
SULPICE	Laurent	PU-PH	Chirurgie générale
TADIE	Jean Marc	PU-PH	Médecine intensive-réanimation
TARTE	Karin	PU-PH	Immunologie
TATTEVIN	Pierre	PU-PH	Maladies infectieuses ; maladies tropicales
THIBAULT	Ronan	PU-PH	Nutrition
THIBAULT	Vincent	PU-PH	Bactériologie-virologie ; hygiène hospitalière
THOMAZEAU	Hervé	PU-PH	Chirurgie orthopédique et traumatologique
TORDJMAN	Sylvie	PU-PH	Pédopsychiatrie ; addictologie
VERHOYE	Jean-Philippe	PU-PH	Chirurgie thoracique et cardiovasculaire
VERIN	Marc	PU-PH	Neurologie
VIEL	Jean-François	PU-PH	Epidémiologie, économie de la santé et prévention
VIGNEAU	Cécile	PU-PH	Néphrologie
VIOLAS	Philippe	PU-PH	Chirurgie infantile
WATIER	Eric	PU-PH	Chirurgie plastique, reconstructrice et esthétique ; brûlologie
WODEY	Eric	PU-PH	Anesthésiologie-réanimation ; médecine d'urgence

## LISTES DES MAITRES DE CONFERENCE DES UNIVERSITES

NOM	PRENOM	TITRE	CNU
ALLORY	Emmanuel	MC Associé	Médecine générale
AME-THOMAS	Patricia	MCU-PH	Immunologie
AMIOT	Laurence	MCU-PH	Hématologie ; transfusion
ANSEMI	Amédéo	MCU-PH	Chirurgie thoracique et cardiovasculaire
BANATRE	Agnès	MC Associé	Médecine générale
BEGUE	Jean Marc	MCU-PH	Physiologie
BERTHEUIL	Nicolas	MCU-PH	Chirurgie plastique, reconstructrice et esthétique ; brûlologie
BOUSSEMART	Lise	MCU-PH	Dermato-vénéréologie
BROCHARD	Charlène	MCU-PH	Physiologie
CABILLIC	Florian	MCU-PH	Biologie cellulaire
CAUBET	Alain	MCU-PH	Médecine et santé au travail
CHHOR-QUENIART	Sidonie	MC Associé	Médecine générale
CORVOL	Aline	MCU-PH	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
DAMERON	Olivier	MCF	Informatique
DE TAYRAC	Marie	MCU-PH	Biochimie et biologie moléculaire
DEGEILH	Brigitte	MCU-PH	Parasitologie et mycologie
DROITCOURT	Catherine	MCU-PH	Dermato-vénéréologie
DUBOURG	Christèle	MCU-PH	Biochimie et biologie moléculaire
DUGAY	Frédéric	MCU-PH	Histologie, embryologie et cytogénétique
EDELINE	Julien	MCU-PH	Cancérologie ; radiothérapie
FIQUET	Laure	MC Associé	Médecine générale
GANGLOFF	Cédric	MC Associé	Médecine d'urgence
GARLANTEZEC	Ronan	MCU-PH	Epidémiologie, économie de la santé et prévention
GOUIN épouse THIBAUT	Isabelle	MCU-PH	Hématologie ; transfusion
GUILLET	Benoit	MCU-PH	Hématologie ; transfusion
JAILLARD	Sylvie	MCU-PH	Histologie, embryologie et cytogénétique
KALADJI	Adrien	MCU-PH	Chirurgie vasculaire ; médecine vasculaire
KAMMERER-JACQUET	Solène-Florence	MCU-PH	Anatomie et cytologie pathologiques
LAVENU	Audrey	MCF	Sciences physico-chimiques et ingénierie appliquée à la santé
LE GALL	François	MCU-PH	Anatomie et cytologie pathologiques
LE GALL	Vanessa	MC Associé	Médecine générale
LEMAITRE	Florian	MCU-PH	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
MARTINS	Pédro Raphaël	MCU-PH	Cardiologie
MATHIEU-SANQUER	Romain	MCU-PH	Urologie
MENARD	Cédric	MCU-PH	Immunologie
MICHEL	Laure	MCU-PH	Neurologie
MOREAU	Caroline	MCU-PH	Biochimie et biologie moléculaire
MOUSSOUNI	Fouzia	MCF	Informatique
PANGAULT	Céline	MCU-PH	Hématologie ; transfusion
RENAUT	Pierrick	MC Associé	Médecine générale
ROBERT	Gabriel	MCU-PH	Psychiatrie d'adultes ; addictologie

SCHNELL	Frédéric	MCU-PH	Physiologie
TURLIN	Bruno	MCU-PH	Anatomie et cytologie pathologiques
VERDIER épouse LORNE	Marie-Clémence	MCU-PH	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
ZIELINSKI	Agata	MCF	Philosophie



## TABLE DES MATIERES

LISTE DES PROFESSEURS DES UNIVERSITES .....	- 2 -
LISTES DES MAITRES DE CONFERENCE DES UNIVERSITES .....	- 5 -
TABLE DES MATIERES .....	- 7 -
LISTE DES FIGURES .....	- 8 -
LISTE DES TABLEAUX .....	- 8 -
INTRODUCTION (FR).....	- 9 -
ABSTRACT .....	- 10 -
INTRODUCTION.....	- 11 -
MATERIALS AND METHOD .....	- 12 -
RESULTS.....	- 14 -
DISCUSSION.....	- 16 -
CONCLUSION .....	- 19 -
PERMIS D'IMPRIMER .....	- 23 -
REFERENCES BIBLIOGRAPHIQUES.....	- 20 -
LISTE DES ABBREVIATIONS .....	- 22 -
FIGURES .....	- 23 -
TABLEAUX .....	- 24 -
ANNEXES.....	- 29 -

## LISTE DES FIGURES

Figure 1 Flow diagram of patients enrolment..... - 23 -

## LISTE DES TABLEAUX

Table 1 Characteristics of 180 cases of staphylococcal prosthetic valve endocarditis treated with, or without, rifampin..... - 24 -

Table 2 Outcomes of 180 episodes of staphylococcal prosthetic valve endocarditis treated with, or without, rifampin..... - 26 -

Table 3 Outcome of prosthetic valve endocarditis due to *Staphylococcus aureus* (n=114), and or coagulase-negative staphylococci (n=66) in patients treated with, or without, rifampin - 27 -

Table 4 Univariate and multivariate analysis of variables associated with one-year mortality . - 28 -

Supplementary Table 1 Characteristics and adverse events in 101 cases of staphylococcal prosthetic valve endocarditis treated by rifampin-based combination. .... - 29 -

Supplementary Table 2 Univariate analysis of outcomes in 149 patients with definite staphylococci prosthetic valve endocarditis ..... - 30 -

Supplementary Table 3 Outcomes of 56 episodes of methicillin-resistant staphylococcal prosthetic valve endocarditis treated with, or without, rifampin ..... - 31 -

Supplementary Table 4 Univariate analysis of outcomes in 128 patients with staphylococci prosthetic valve endocarditis without valve surgery ..... - 32 -

## INTRODUCTION (FR)

L'endocardite infectieuse est une maladie rare, d'une particulière gravité, et dont la prise en charge est régie par des recommandations régulièrement actualisées. Malgré un volume de publications sur ce thème important et une concertation internationale efficace concernant sa prise en charge, de nombreuses zones d'ombre persistent.

Parmi elles, l'intérêt de la rifampicine dans le traitement des endocardites à *Staphylococcus* sp. lorsque l'infection survient sur valve prothétique, n'est pas démontré. Cette thèse, présentée sous forme d'article est le résultat d'un travail initié depuis le début de mon internat. L'objectif était d'analyser la prescription de rifampicine dans cette indication, et son impact sur le pronostic, par une étude rétrospective menée dans 3 centres hospitaliers universitaires du grand ouest : Rennes, Nantes et Brest. Ce travail a fait l'objet d'une publication dans *Clinical Infectious Diseases* en juillet 2020.

## ABSTRACT

**Background.** International guidelines recommend rifampin-based combinations for staphylococcal prosthetic valve endocarditis (PVE). However, no robust clinical data supports this recommendation, and rifampin tolerability is an issue. We aimed to evaluate the impact of rifampin for the treatment of staphylococcal PVE.

**Methods.** An observational retrospective cohort study of all adults with staphylococcal PVE (modified Duke criteria) was conducted in three referral centers for endocarditis, during years 2000-2018. Primary outcome measurement was one-year mortality.

**Results.** We enrolled 180 patients with PVE due to *Staphylococcus aureus* (n=114, 63.3%), or coagulase-negative staphylococci (n=66, 36.7%), on bioprosthesis (n=111, 61.7%), mechanical valve (n=67, 37.2%), or both (n=2). There were 132 males (73.3%), and mean age was 70.4±12.4 years. Valvular surgery was performed in 51/180 (28.3%) cases. Despite all isolates were susceptible to rifampin, only 101 (56.1%) were treated with rifampin, for a median duration of 33.0 days, while 79 (43.9%) received no rifampin. Baseline characteristics were similar in both groups. One-year mortality was, respectively, 37.6% (38/101), and 31.6% (25/79), in patients treated with, or without, rifampin ( $P=0.62$ ). Relapse rates were 5.9% (6/101), and 8.9% (7/79),  $P=0.65$ . Patients treated with rifampin had longer hospital length-of-stay: 42.3±18.6 vs. 31.3±14.0 days ( $P<0.0001$ ). On multivariate analysis, only cerebral emboli (OR 2.95, CI95% 1.30-6.70,  $P=0.009$ ), definite endocarditis (OR 7.15, 1.47-34.77,  $P=0.018$ ), and methicillin-resistant *S. aureus* (OR 6.04, 1.34-27.26,  $P=0.019$ ), were associated with one-year mortality.

**Conclusions.** A large proportion (43.9%) of staphylococcal PVE received no rifampin. One-year survival and relapse rates were similar in patients treated with or without rifampin.

## INTRODUCTION

The profile of infective endocarditis (IE) has dramatically changed over the last decades, with the emergence of healthcare-associated IE (1,2), including prosthetic valve IE (PVE). In parallel, staphylococci became the leading cause of IE in most contemporary cohorts. The prospective cohort study of the International Collaboration on Endocarditis (ICE) found that *Staphylococcus aureus* and coagulase-negative staphylococcus (CoNS) are responsible for, respectively, 23% and 17% of PVE (3). Staphylococcal PVE are associated with high one-year mortality rates, ranging from 40% to 80% (4). Hence, optimal antibacterial treatment is of paramount importance for staphylococcal PVE.

The two major international guidelines, from America and Europe, are remarkably concordant regarding antibiotic regimen to apply in patients with staphylococcal PVE. It relies on intravenous (i.v.) anti-staphylococcal penicillin or cefazolin for methicillin-susceptible staphylococci, and i.v. glycopeptides or lipopeptides for methicillin-resistant staphylococci (5), combined with i.v. gentamicin during the first two weeks, and rifampin for the whole duration of treatment, i.e. six weeks. Minor discrepancies are found regarding rifampin administration, with a recommended daily dose from 600 to 1200 mg, oral or i.v., divided in 2 or 3 intakes per day (6,7), but its use is strongly encouraged, labelled as class 1 recommendation both for the American Heart Association (7), and the European Society of Cardiology (6). However, due to the absence of any randomized clinical trial supporting this position, this recommendation is only associated with a B level of evidence.

Rifampin is a rifamycin B derivative, which inhibits bacterial RNA polymerase by blocking the path of elongating RNA (8). It is believed to have a special role in prosthetic devices infection, due to its activity on planktonic bacteria embedded in biofilms, which would contribute to eradicate bacteria attached on foreign material, thereby reducing the risk of relapse (5,6). However, experimental and clinical data on the impact of rifampin-based combinations remain limited in the field of PVE, and rifampin use has been associated with severe adverse events, including the potential for interactions with a large number of drugs (anticoagulants, antiepileptics, immunomodulators, etc.) (9). Given the poor level of evidence supporting the systematic use of rifampin-based combination for the treatment of staphylococcal PVE, we aimed to evaluate the risks and the benefits associated with rifampin use through a retrospective multicenter study of patients managed in referral centers for IE.

## MATERIALS AND METHOD

### Design, Setting and Patients

This observational multicenter retrospective cohort study was conducted in three referral centers for IE in western France: Brest, Nantes and Rennes (population catchment area, 7 million inhabitants). In these centers, all patients with suspected IE are managed by a multidisciplinary team including members from cardiac surgery, cardiology, infectious diseases, and microbiology departments. Most patients received medical and surgical treatment in agreement with international guidelines, i.e. the American Heart Association (AHA) guidelines until 2009, the European Society of Cardiology (ESC) guidelines from 2009 to 2015, and a combination of both since they were updated in 2015.

All adult patients ( $\geq 18$  years old) with a diagnosis of staphylococcal PVE treated from January 2000 through June 2018 in Rennes, and from January 2010 through June 2018 in Nantes and Brest, were enrolled in the study. Patients were identified using the French hospital discharge database (French acronym PMSI): All medical charts with a discharge diagnosis of IE were individually screened. In one site (Nantes), a prospective cohort study of all patients with IE was initiated in January 2013. Hence, for this site, patients were enrolled through this cohort between January 2013 and July 2018. Patients were included if they were classified as definite or possible IE according to the modified Duke criteria (10), due to *S. aureus* or CoNS, with the involvement of at least one prosthetic valve. The exclusion criteria were history of congenital heart disease, aortic root replacement, and transcatheter valve replacement.

### Data collection

Data for clinical, microbiological, echocardiographic variables, as well as management of PVE and follow-up, were collected on a standardized questionnaire from medical charts. The data collected were entered into a database created specifically for the study by a single investigator (ALB).

### Ethics

The study was approved by an institutional review board, the Ethical Committee of Research in Tropical and Infectious Diseases (CER-MIT, n°2019-0703).

### Definitions

All study definitions, outcomes, and variables were determined *a priori*. The Charlson comorbidity index (11) was calculated retrospectively with the comorbidities reported at the time of admission. Early-PVE was defined as IE occurring within 60 days after prosthetic valve implantation (5). Healthcare-associated IE was defined as either nosocomial (IE developing in

a patient hospitalized for more than 48 hours), or non-nosocomial healthcare-associated infection, as defined elsewhere (3). Intracardiac devices were permanent pacemakers or cardioverter-defibrillators. Cerebral emboli were defined as acute onset of neurological symptoms attributed to stroke and confirmed on brain imaging. Early valve surgery was defined as valve surgery within 60 days after the diagnosis of endocarditis (4). All patients received appropriate i.v. antibiotics, *i.e.* cefazolin or anti-staphylococcal penicillin (cloxacillin, oxacillin), combined with gentamicin for methicillin-susceptible staphylococci, and glycopeptides (vancomycin), or lipopeptides (daptomycin), combined with gentamicin for methicillin-resistant staphylococci, at the doses recommended in the 2015 international guidelines (6,7).

### **Outcomes**

The primary endpoint was death from any cause during the one-year follow-up. Relapse was defined as a new diagnosis of IE caused by the same microorganism as the initial PVE, within six months. Hospital length of stay was defined as the time from the first positive blood culture, to discharge. Vitamin K antagonists (VKA) imbalance was defined by two consecutive International Normalized Ratio (INR)  $>4$  or  $<1.5$  more than 7 days after treatment start. Bleeding complications were defined as hemoglobin decrease  $>2$  g/dL between two measurements, or any life-threatening hemorrhage reported in the medical chart.

### **Statistical Analysis**

We compared two mutually exclusive groups of patients: i) patients who were treated with rifampin-based combination; ii) patients who received no rifampin throughout PVE treatment. Patients were allocated to the rifampin group if they received at least one dose of rifampin. Quantitative variables were expressed as mean  $\pm$  standard deviation, or as median with interquartile range [IQR], as appropriate. Qualitative variables were expressed by frequency and percentage. Continuous variables were compared using the Mann-Whitney test or the exact *t*-test, as appropriate. Categorical variables were compared using the Chi<sup>2</sup> test with Yates continuity correction. Multivariate analyses were performed using exact logistic regression. Clinically relevant factors associated to outcome with  $P<0.20$  in the univariate analysis, and variables associated with 1-year-mortality in the literature were included in the multivariate model. All tests were two-tailed, and significance was set at  $P<0.05$ . Statistical analyses were performed using PRISM (v5.0 for Windows, GraphPad Software, Inc., CA, USA) and STATA (v. 9.0 for Windows, Statacorp LLC, TX, USA).

## RESULTS

### Demographics and baseline characteristics

During the study period, 180 episodes of staphylococcal PVE were managed in the participating sites, classified as definite (n=149), or possible (n=31). Of them, 101 (56.1%) received an antibiotic regimen with rifampin, and 79 (43.9%) were treated without rifampin (Fig 1). They were 132 males (73.3%), and 48 females (26.7%), with a mean age of  $70.4 \pm 12.4$  years. PVE occurred on bioprosthesis in 111 cases (61.7%), mechanical valves in 67 (37.2%), and both in two (1.1%). Cerebral emboli were reported in 53 (29.4%) cases. The causative agent was *S. aureus* in 114 (63.3%) episodes, including 17 (14.9%) with methicillin resistance. CoNS was responsible for 66 (36.7%) episodes, including 39 (59.0%) with methicillin resistance. None of the 180 isolates was resistant to rifampin. Valvular surgery was performed in 51 (28.3%) cases, most of them within 60 days after PVE diagnosis (94%). Sixty-two patients (34.4%) received VKA during the antibacterial treatment of their PVE. Baseline demographic, clinical and microbiological features of the 2 groups were similar, except for the proportion of *S. aureus* isolates resistant to methicillin, at 21.9% (14/64) in patients treated with rifampin, vs 6.0% (3/50) in patients who received no rifampin,  $P=0.04$  (Table 1).

### Rifampin treatment

The median duration of rifampin was 33.0 days (IQR, 12.5 – 41.2) with a median dose of 1200 mg per day (IQR, 900 – 1200 mg). Median time between first positive blood culture and rifampin start was 7 days (IQR, 3 - 15 days). Rifampin had to be prematurely discontinued because of severe adverse events in 31 patients (30.7%), mainly because of liver toxicity (n=11), digestive disorders (n=4), cytopenia (n=4), VKA imbalance (n=3), renal toxicity (n=2), allergy (n=2), treatment failure (n=2), or not specified (n=3). Of the 35 patients who were treated with VKA when rifampin was introduced, 15 (42.9%) presented VKA imbalance during rifampin treatment or during the week following rifampin discontinuation, deemed to be related to drug interactions (Supplementary Table 1). Four episodes of drug interactions with drugs other than VKA were identified, with methadone (n=2), phenytoin (n=1), and mianserin (n=1). None of these were responsible for rifampin discontinuation. No emergence of resistance to rifampin was observed in the six patients who relapsed with positive blood cultures.

### Outcome

In-hospital mortality was 23.6% (42/180), and one-year mortality was 35.4% (63/180). Thirteen patients (7.3%) relapsed. Outcomes were similar in patients treated with, or without rifampin, except for hospital length-of-stay, with a mean of  $42.3 \text{ days} \pm 18.6$  in patients treated with rifampin (n=101) vs  $31.3 \text{ days} \pm 14.0$  in patients who did not receive rifampin,  $P<0.0001$  (Table 2). Results were similar when the analysis was stratified by pathogen (*S. aureus* and CoNS,



Table 3), when it was restricted to patients with definite IE according to modified Duke classification (supplementary table 2), to patients with PVE due to methicillin-resistant staphylococci (supplementary table 3), or when it was restricted to patients who underwent no valve replacement (supplementary table 4).

The only factor associated with one-year mortality on univariate analysis was cerebral emboli. On multivariate analysis, cerebral emboli (odds ratio 2.95, CI95% 1.30-6.70,  $P=0.009$ ), definite endocarditis according to modified Duke criteria (OR 7.15, 1.47-34.77,  $P=0.018$ ), and methicillin-resistant *S. aureus* (OR 6.04, 1.34-27.26,  $P=0.019$ ), were independently associated with one-year mortality (Table 4).

## DISCUSSION

The major findings of this retrospective study of staphylococcal PVE managed in 3 referral centers for IE are the following: i) in spite of international guidelines, a large proportion of patients with rifampin-susceptible staphylococcal PVE are treated without rifampin (79/180, 43.9%); ii) baseline characteristics of patients treated with, or without rifampin, were similar; iii) rifampin use was not associated with any measurable benefit, neither for survival, nor for risk of relapse; iv) patients treated with rifampin had longer hospital length-of-stay, despite similar incidence of severe adverse events; v) the three variables independently associated with one-year mortality were cerebral emboli, definite endocarditis, and methicillin-resistant *S. aureus* PVE.

These discrepancies between international guidelines, and practices, regarding rifampin use in patients with staphylococcal PVE in three referral centers for IE, may be related to the limited clinical evidence supporting these guidelines. Pioneer studies in this field were of limited sample size, and mostly enrolled *Staphylococcus epidermidis* PVE (12,13). More recent studies found no difference in valve sterilization rate in patients with *S. aureus* IE treated with, or without rifampin (14), and that the addition of rifampin after valvular replacement during the acute phase of staphylococcal IE was not associated with lower mortality, or decreased risk of relapse (15). A case-control study even suggested that rifampin use could be associated with increased mortality, and longer duration of staphylococcal bacteremia, in patients with *S. aureus* native valve IE (16). Nevertheless, beneficial effect of rifampin on another type of foreign-device infections (i.e. prosthetic joint) has been documented in a randomized trial (17). In addition, *ex vivo* studies and animal models have documented that rifampin is remarkable for its penetration, and bactericidal activity against dormant staphylococci within the biofilm (18–20). A recent post-hoc analysis of a large *S. aureus* bacteremia cohort suggested that antimicrobials combination, most of them containing rifampin, would be associated with reduced mortality and lower risk of late complications in patients with implanted foreign bodies (21).

The second reason why rifampin use is low in patients with staphylococcal PVE could be related to its association with adverse events, and drug interactions. Rifampin-related liver toxicity is not rare, mostly affects patients with underlying liver diseases (22), and may lead to treatment discontinuation in as much as 30% of patients (16,17), as in our cohort. Moreover, rifampin is a strong inducer of cytochrome P450 3A4 expression, thereby reducing plasma concentrations of several drugs (8). In our study, most clinically significant interactions were reported with VKA, but we found no increase of thrombo-embolic events, or bleeding complications. This may be due to close monitoring of these interactions in our centers experienced with rifampin use, primarily for treatment of staphylococcal prosthetic joint

infections, and tuberculosis. The third potential reason behind the reluctance to use rifampin in staphylococcal PVE rifampin could be the theoretical risk of rifampin resistance emergence, especially in case of high inoculum or inappropriate backbone antistaphylococcal regimen (13,14). In PVE, some experts recommend delaying rifampin initiation after bacteremia has been cleared, to allow penetration and activity of other antibiotics into valvular vegetations before rifampin start (7). In our study, rifampin was started with a median delay of 7 days after first positive blood culture, and no emergence of rifampin resistance was observed.

Despite similar baseline characteristics in patients treated with (n=101), or without (n=79) rifampin, rifampin use was associated neither with better survival, nor with reduced risk of relapse. This suggests that rifampin has no added value for the eradication of staphylococci from valvular prosthesis, contrarily to what experimental studies suggested. The only significant difference between patients treated with, or without rifampin, was the longer duration of hospital stay in patients treated with rifampin. This may be related to VKA imbalance, defined as INR>4 or <1.5, and documented in 42.9% of patients co-treated with VKA and rifampin in our study, as this is an indication to delay patients discharge in our practices. Drug interactions between rifampin and VKA are usually anticipated by increasing VKA dosing, under close supervision of INR. Hence, as the degree of drug interaction varies from one patient to another, the comedication may result in INR too low, or too high. This may delay patients discharge, which may be an explanation why, although we found no significant differences between patients treated with, or without, rifampin, for all outcomes criteria, the length of stay was significantly longer in patients who received rifampin ( $42.3 \pm 18.6$  days), than in patients who did not receive rifampin ( $31.3 \pm 14.0$  days,  $P<0.0001$ ). Of note, although rifampin was not associated with one-year survival, we found that definite IE according to modified Duke criteria, cerebral emboli, and methicillin-resistant PVE were all independent predictors of one-year mortality, in line with previous studies in the field (1,4,23,24), thus corroborating the relevance of our cohort study.

Our study has limitations. Firstly, as data were collected retrospectively, this study has potential bias, and confounding factors. However, the rate of missing data was very low, and data were collected by a single investigator (ALB), hence reducing heterogeneity. Secondly, although this study is, to our knowledge, the largest on staphylococcal PVE to date, it may have been underpowered to detect any beneficial effect of rifampin, especially in certain subgroups (e.g. patients with no valve replacement), due to limited sample size. This limitation is particularly relevant for the effect of rifampin on the prevention of relapse, as only 13 relapses were identified in our study. Thirdly, as this observational study was conducted in three distinct sites, over a large period (2000-2018), there may be some heterogeneity in practices. However, this study was performed in referral centers, where patients were managed by a multidisciplinary team, according to international guidelines, so that these findings may apply to most sites with

similar practices. Lastly, we have no data on rifampin exposure for the patients enrolled in this study, although wide inter-individual variability has been reported with this drug. Therapeutic drug monitoring of rifampin plasma concentrations may optimize its efficacy and tolerability, by allowing tailored rifampin dosing.

## **CONCLUSION**

In conclusion, we found that a large proportion (43.9%) of staphylococcal PVE receive no rifampin in referral centers for IE, in spite of the strong recommendation for the use of rifampin-based combinations in international guidelines. Rifampin use was associated neither with better survival, nor with lower risk of relapse. These findings advocate for a randomized controlled trial to evaluate the impact of rifampin use for the treatment of staphylococcal PVE. Meanwhile, our study suggests that patients with staphylococcal PVE may be reasonably treated without rifampin in case of contra-indication, or poor tolerability.

## REFERENCES BIBLIOGRAPHIQUES

1. Duval X, Delahaye F, Alla F, Tattevin P, Obadia J-F, Le Moing V, *et al.* Temporal trends in infective endocarditis in the context of prophylaxis guideline modifications: three successive population-based surveys. *J Am Coll Cardiol.* 29 mai 2012;59(22):1968-76.
2. Fowler VG, Miro JM, Hoen B, Cabell CH, Abrutyn E, Rubinstein E, *et al.* *Staphylococcus aureus* endocarditis: a consequence of medical progress. *JAMA.* 2005;293(24):3012-21.
3. Wang A, Athan E, Pappas PA, Fowler VG, Olaison L, Paré C, *et al.* Contemporary clinical profile and outcome of prosthetic valve endocarditis. *JAMA* 2007;297(12):1354-61.
4. Chirouze C, Cabell CH, Fowler VG, Khayat N, Olaison L, Miro JM, *et al.* Prognostic factors in 61 cases of *Staphylococcus aureus* prosthetic valve infective endocarditis from the International Collaboration on Endocarditis merged database. *Clin Infect Dis* 2004;38(9):1323-7.
5. Saeed K, Bal AM, Gould IM, David MZ, Dryden M, Giannitsioti E, *et al.* An update on *Staphylococcus aureus* infective endocarditis from the International Society of Antimicrobial Chemotherapy (ISAC). *Int J Antimicrob Agents* 2019;53(1):9-15.
6. Habib G, Lancellotti P, Antunes MJ, Bongiorni MG, Casalta J-P, Del Zotti F, *et al.* 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J* 2015;36(44):3075-128.
7. Baddour LM, Wilson WR, Bayer AS, Fowler VG, Tleyjeh IM, Rybak MJ, *et al.* Infective Endocarditis in Adults: Diagnosis, Antimicrobial Therapy, and Management of Complications: A Scientific Statement for Healthcare Professionals From the American Heart Association. *Circulation* 2015;132(15):1435-86.
8. Niemi M, Backman JT, Fromm MF, Neuvonen PJ, Kivistö KT. Pharmacokinetic interactions with rifampicin : clinical relevance. *Clin Pharmacokinet.* 2003;42(9):819-50.
9. Thwaites GE, Scarborough M, Szubert A, Nsutebu E, Tilley R, Greig J, *et al.* Adjunctive rifampicin for *Staphylococcus aureus* bacteraemia (ARREST): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet* 2018;391(10121):668-78.
10. Li JS, Sexton DJ, Mick N, Nettles R, Fowler VG, Ryan T, *et al.* Proposed modifications to the Duke criteria for the diagnosis of infective endocarditis. *Clin Infect Dis* 2000;30(4):633-8.
11. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987;40(5):373-83.
12. Karchmer AW, Archer GL, Dismukes WE. *Staphylococcus epidermidis* causing prosthetic valve endocarditis: microbiologic and clinical observations as guides to therapy. *Ann Intern Med* 1983;98(4):447-55.

13. Karchmer AW, Archer GL, Dismukes WE. Rifampin treatment of prosthetic valve endocarditis due to *Staphylococcus epidermidis*. Rev Infect Dis 1983;5 Suppl 3:S543-548.
14. Drinković D, Morris AJ, Pottumarthy S, MacCulloch D, West T. Bacteriological outcome of combination versus single-agent treatment for staphylococcal endocarditis. J Antimicrob Chemother 2003;52(5):820-5.
15. Shrestha NK, Shah SY, Wang H, Hussain ST, Pettersson GB, Nowacki AS, *et al.* Rifampin for Surgically Treated Staphylococcal Infective Endocarditis: A Propensity Score-Adjusted Cohort Study. Ann Thorac Surg 2016;101(6):2243-50.
16. Riedel DJ, Weekes E, Forrest GN. Addition of rifampin to standard therapy for treatment of native valve infective endocarditis caused by *Staphylococcus aureus*. Antimicrob Agents Chemother 2008;52(7):2463-7.
17. Zimmerli W, Widmer AF, Blatter M, Frei R, Ochsner PE. Role of rifampin for treatment of orthopedic implant-related staphylococcal infections: a randomized controlled trial. Foreign-Body Infection (FBI) Study Group. JAMA 1998;279(19):1537-41.
18. Zheng Z, Stewart PS. Penetration of rifampin through *Staphylococcus epidermidis* biofilms. Antimicrob Agents Chemother 2002;46(3):900-3.
19. Saginur R, Stdenis M, Ferris W, Aaron SD, Chan F, Lee C, *et al.* Multiple combination bactericidal testing of staphylococcal biofilms from implant-associated infections. Antimicrob Agents Chemother 2006;50(1):55-61.
20. Revest M, Jacqueline C, Boudjemaa R, Caillon J, Le Mabeque V, Breteche A, *et al.* New *in vitro* and *in vivo* models to evaluate antibiotic efficacy in *Staphylococcus aureus* prosthetic vascular graft infection. J Antimicrob Chemother 2016;71(5):1291-9.
21. Rieg S, Joost I, Weiß V, Peyerl-Hoffmann G, Schneider C, Hellmich M, *et al.* Combination antimicrobial therapy in patients with *Staphylococcus aureus* bacteraemia-a *post hoc* analysis in 964 prospectively evaluated patients. Clin Microbiol Infect 2017;3(6):406.e1-406.e8.
22. Galar A, Weil AA, Dudzinski DM, Muñoz P, Siedner MJ. Methicillin-Resistant *Staphylococcus aureus* Prosthetic Valve Endocarditis: Pathophysiology, Epidemiology, Clinical Presentation, Diagnosis, and Management. Clin Microbiol Rev 2019;32(2):e00041-18, /cmr/32/2/CMR.00041-18.atom.
23. Park LP, Chu VH, Peterson G, Skoutelis A, Lejko-Zupa T, Bouza E, *et al.* Validated Risk Score for Predicting 6-Month Mortality in Infective Endocarditis. J Am Heart Assoc.2016;5(4):e003016.
24. Fauchier L, Pericart L, Bourguignon T, Genet T, Bisson A, Bernard A, *et al.* Comparison of Outcome of Possible Versus Definite Infective Endocarditis Involving Prosthetic or Bioprosthetic Heart Valves. Am J Cardiol 2017 ;120(10) :1884-90.

## LISTE DES ABBREVIATIONS

CoNS: coagulase negative *Staphylococcus*

HLOS: Hospital length of stay

IE: Infective endocarditis

INR: International Normalized Ratio

IQR : Interquartile Range

I.V. : Intraveinuous

MRSA: Methicillin resistant *Staphylococcus aureus*

PVE: prosthetic valve endocarditis


SA: *Staphylococcus aureus*

VKA: Vitamin K antagonists



## FIGURES

Figure 1 Flow diagram of patients enrolment.



## TABLEAUX

**Table 1 Characteristics of 180 cases of staphylococcal prosthetic valve endocarditis treated with, or without, rifampin**

Variable	Total (n=180)	Rifampin- based combination (n=101)	No rifampin (n=79)	P Value
<b>Demographic features</b>				
Age, years	70.4 ± 12.4	69.0 ± 12.8	72.2 ± 11.6	.08
Gender, male	132 (73.3)	74 (73.3)	58 (73.4)	.88
Charlson comorbidity index	4.7 ± 2.3	4.7 ± 2.4	4.5 ± 2.2	.56
Intravenous drug user	5 (2.8)	4 (4.0)	1 (1.3)	.53
Cirrhosis	9 (5.0)	4 (4.0)	5 (6.3)	.70
Healthcare-associated infection	108 (60.0)	57 (56.4)	51 (64.6)	.34
<b>Clinical features</b>				
Definite endocarditis (modified Duke)	149 (82.8)	88 (87.1)	61 (77.2)	.12
Type of prosthetic valve				
Bioprosthesis	111 (61.7)	60 (59.4)	51 (64.6)	.58
Mechanical prosthesis	67 (37.2)	41 (40.6)	26 (32.9)	.37
Both	2 (1.1)	0 (0)	2 (2.5)	.37
Prosthetic valve endocarditis location				
Aortic	138 (76.7)	78 (77.2)	60 (75.9)	.98
Mitral	25 (13.9)	14 (13.9)	11 (13.9)	.84
Tricuspid	3 (1.7)	2 (2.0)	1 (1.3)	.82
Pulmonary	1 (0.6)	1 (1.0)	0 (0)	.90
Multiple	13 (7.2)	6 (5.9)	7 (8.9)	.65
Interval between prosthetic valve implantation and endocarditis, months	32 (4-104.5]	39 [8-117.5]	24 [1-98]	.16
Early prosthetic valve endocarditis*	34 (18.9)	14 (13.9)	20 (25.3)	.08
Intracardiac device	38 (21.1)	19 (18.8)	19 (24.0)	.50
Osteoarticular infection	18 (10.0)	8 (7.9)	10 (12.7)	.42
Cerebral emboli	53 (29.4)	35 (34.7)	18 (28.8)	.12
<b>Microbiological features</b>				
<i>Staphylococcus</i> species				
<i>Staphylococcus aureus</i>	114 (63.3)	64 (63.4)	50 (63.3)	.88
Coagulase-negative staphylococci	66 (36.7)	37 (36.6)	29 (36.7)	.88
Methicillin resistance				
<i>Staphylococcus aureus</i>	17 (14.9)	14 (21.9)	3 (6.0)	.04
Coagulase-negative staphylococci	39 (59.0)	24 (64.9)	15 (51.7)	.41
Rifampin resistance	0 (0)	0 (0)	0 (0)	
Duration of bacteremia, days	5.5 ± 3.0	5.6 ± 3.6	5.5 ± 3.5	.98
Positive valve culture if surgery	11 (22)	5 (15.6)	6 (33.3)	.19

**Table 1 (part 2)**

<b>Variable</b>	<b>Total (n=180)</b>	<b>Rifampin- based combination (n=101)</b>	<b>No rifampin (n=79)</b>	<b>P Value</b>
<b>Treatment</b>				
Valve surgery	51 (28.3)	34 (33.7)	17 (21.5)	.10
Early valve surgery <sup>§</sup>	48 (94)	32 (94.1)	16 (94.1)	.53
Interval between first positive blood culture and surgery, days	13 [8–20.2]	14.0 [11–19.5]	10.0 [5-17]	.23
Osteoarticular surgery	5 (27.8)	3 (37.5)	2 (20.0)	.77
Lifelong suppressive antibacterial treatment	7 (3.9)	2 (2.0)	5 (6.3)	.27
Vitamin K antagonists	62 (34.4)	35 (34.7)	27 (34.2)	.93
Heparin	83 (46.1)	52 (51.5)	31 (39.2)	.14

Quantitative variables are expressed as mean +/- standard deviation or median (IQR), qualitative variables are expressed by numbers (%); \* time between valve implantation, and endocarditis, <60 days; <sup>§</sup> Cardiac surgery within 60 days after endocarditis diagnosis

**Table 2 Outcomes of 180 episodes of staphylococcal prosthetic valve endocarditis treated with, or without, rifampin**

<b>Variable</b>	<b>Total (n=180)</b>	<b>Rifampin- based combination (n=101)</b>	<b>No rifampin (n=79)</b>	<b>Odds-Ratio (CI 95%)</b>	<b>P Value</b>
<b>Mortality</b>					
In-hospital mortality	42 (23.6)	26 (25.7)	16 (20.3)	1.4 (0.67-2.77)	.49
Six-month mortality	58 (32.6)	36 (35.6)	22 (27.8)	1.4 (0.76-2.72)	.34
One-year mortality	63 (35.4)	38 (37.6)	25 (31.6)	1.2 (0.66-2.28)	.62
<b>Relapse</b>	13 (7.3)	6 (5.9)	7 (8.9)	0.64 (0.21-2.02)	.65
Vitamin K antagonist imbalance during endocarditis	21 (33.9)	15 (42.9)	6 (22.2)	2.63 (0.85-8.11)	.15
<b>Bleeding complication</b>	23 (12.9)	13 (12.8)	10 (12.7)	1.02 (0.42-2.46)	.85
Length of stay, days	37 ± 17.6	42.3 ± 18.6	31.3 ± 14.0	-	<.0001

Quantitative variables are expressed as mean +/- standard deviation, qualitative variables are expressed by numbers (%)

**Table 3 Outcome of prosthetic valve endocarditis due to *Staphylococcus aureus* (n=114), and or coagulase-negative staphylococci (n=66) in patients treated with, or without, rifampin**

Variable	<i>Staphylococcus aureus</i> (n= 114)				Coagulase negative staphylococci (n= 66)			
	Rifampin- based (n= 64)	No rifampin (n= 50)	Odd- Ratio (CI 95%)	P Value	Rifampin- based (n= 37)	No rifampin (n= 29)	Odd- Ratio (CI 95%)	P Value
<b>Mortality</b>								
In-hospital mortality	18 (28.1)	12 (24.0)	1.24 (0.53-2.89)	.78	8 (21.6)	4 (13.8)	1.72 (0.46-6.41)	.61
Six-month mortality	26 (40.6)	16 (32.0)	1.45 (0.66-3.16)	.45	10 (27.0)	6 (20.7)	1.42 (0.45-4.50)	.76
One-year mortality	27 (42.2)	18 (36.0)	1.30 (0.61-2.78)	.63	11 (29.7)	7 (24.1)	1.33 (0.44-4.01)	.82
Relapse	4 (6.3)	4 (8.0)	0.93 (0.22-3.91)	.79	2 (5.4)	3 (10.3)	0.49 (0.08-3.18)	.78
Vitamin K antagonist imbalance	9 (39.1)	4 (22.2)	2.25 (0.56-9.05)	.41	6 (50.0)	2 (22.2)	3.5 (0.50-24.3)	.40
Bleeding complication	10 (15.6)	10 (20.0)	0.72 (0.28-1.95)	.71	3 (8.1)	0 (0)	5.99 (0.29-120.8)	.33
Length of stay, days	42.8±20.1	30.7±14.7	-	.0006	41.4±16.1	32.4±12.9	-	.02

**Table 4 Univariate and multivariate analysis of variables associated with one-year mortality**

Variable	Univariate			Multivariate		
	Dead during the 1-year follow-up (n=63)	Alive at 1-year (n=117)	Odd-Ratio (CI 95%)	P Value	Odd-Ratio (CI 95%)	P Value
Age, per one year increment	70.6 ± 13.2	70.3 ± 11.9	0.86 (0.43-1.71)	.73	0.98 (0.94-1.02)	.45
Gender, male	45 (71.4)	87 (74.4)		.80		
Charlson comorbidity index, per one point increment	5.1 ± 2.6	4.5 ± 2.1		.12	1.14 (0.91-1.44)	.24
Healthcare-associated infection	35 (56.6)	73 (64.0)	0.72 (0.39-1.37)	.41		
Definite endocarditis (modified Duke criteria)	57 (90.5)	92 (78.6)	2.38 (0.91-6.19)	.11	7.15 (1.47-34.77)	.018
Bioprostheses	38 (60.3)	75 (64.1)	0.85 (0.45-1.60)	.73		
Mechanical prosthesis	25 (39.7)	43 (36.8)	1.16 (0.62-2.19)	.76		
Aortic location	44 (69.8)	94 (80.3)	0.57 (0.28-1.15)	.16	0.79 (0.25-2.46)	.68
Mitral location	8 (12.7)	17 (14.5)	0.86 (0.36-2.11)	.91		
Interval between prior cardiac surgery and endocarditis	31 (4-119)	35 (3-103)		.87	0.90 (0.62-1.29)	.56
Intracardiac device	11 (17.5)	27 (23.1)	0.71 (0.32-1.54)	.49		
Cerebral emboli	27 (42.9)	26 (22.2)	2.62 (1.35-5.10)	.006	2.95 (1.30-6.70)	.009
<i>Staphylococcus aureus</i>	45 (71.4)	69 (59.0)	1.74 (0.90-3.36)	.14		
Methicillin-resistant <i>S. aureus</i>	9 (14.3)	8 (6.8)	2.27 (0.83-6.22)	.17	6.04 (1.34-27.26)	.019
Methicillin-resistant coagulase-negative staphylococci	13 (20.6)	26 (22.2)	0.91 (0.43-1.93)	.95		
Duration of bacteremia	5.8 ± 3.3	5.4 ± 3.7		.37	0.88 (0.24-3.24)	.85
Valve surgery	14 (22.2)	36 (30.8)	0.64 (0.32-1.31)	.30	0.60 (0.24-1.52)	.29
Vitamin K antagonist during endocarditis treatment	17 (27.0)	45 (38.5)	0.59 (0.30-1.16)	.16	0.63 (0.26-1.56)	.32
Bleeding complication	6 (9.5)	12 (10.3)	0.92 (0.33-2.59)	.92		
Rifampin treatment	38 (60.3)	63 (53.8)	1.30 (0.70-2.42)	.50	0.90 (0.38-2.11)	.81

Quantitative variables are expressed as mean +/- standard deviation or median (IQR) as appropriate, qualitative variables are expressed by numbers (%)

## ANNEXES

**Supplementary Table 1 Characteristics and adverse events in 101 cases of staphylococcal prosthetic valve endocarditis treated by rifampin-based combination.**

<b>Variable (n=101)</b>	
Duration of rifampin, days	33.0 (28.8)
Rifampicin dose, mg per day	1200 (300)
Interval between first positive blood culture and rifampin start, days	7 (12.0)
Interval between first rifampin dose and surgery, days	7 (15.3)
<b>Reasons for stopping rifampin treatment</b>	
End of planned duration of treatment	50 (49.5)
Death / withholding treatment, no. (%)	20 (19.8)
Liver toxicity	12 (11.9)
Digestive disorder	4 (4.0)
Cytopenia(s)	4 (4.0)
Vitamin K antagonist imbalance	3 (3.0)
Renal toxicity	2 (2.0)
Others	9 (8.9)
<b>Drug interactions</b>	
Concomitant vitamin K antagonist	35 (34.7)
Vitamin K antagonist imbalance during rifampin treatment or the week after rifampin stop	15 (42.9)
Other clinically significant interactions	4 (4.0)

Quantitative variables are expressed as median (IQR), qualitative variables are expressed by numbers (%)

**Supplementary Table 2 Univariate analysis of outcomes in 149 patients with definite staphylococci prosthetic valve endocarditis**

<b>Variable</b>	<b>Total (n=149)</b>	<b>Rifampin- based combination (n=88)</b>	<b>No rifampin (n=61)</b>	<b>Odd-Ratio (CI 95%)</b>	<b>P Value</b>
<b>Mortality</b>					
In-hospital mortality	39 (26.2)	25 (28.4)	14 (23.0)	1.33 (0.63-2.84)	.57
Six-month mortality	52 (34.9)	33 (37.5)	19 (31.1)	1.3 (0.66-2.65)	.53
One-year mortality	57 (38.3)	35 (39.8)	22 (36.1)	1.2 (0.60-2.30)	.77
<b>Relapse</b>	15 (31.9)	10 (38.5)	5 (23.8)	2.0 (0.56-7.18)	.45
Vitamin K antagonist imbalance during endocarditis	18 (12.1)	11 (12.5)	6 (9.8)	1.31 (0.46-3.76)	.81
Bleeding complication	11 (7.4)	5 (5.7)	7 (8.9)	0.46 (0.14-1.54)	.33
Length of stay, days	38 ± 18.0	42.0 ± 19.1	31.7 ± 14.2	-	.0002

Quantitative variables are expressed as mean +/- standard deviation, qualitative variables are expressed by numbers (%).



**Supplementary Table 3 Outcomes of 56 episodes of methicillin-resistant staphylococcal prosthetic valve endocarditis treated with, or without, rifampin**

<b>Variable</b>	<b>Rifampin-based (n= 38)</b>	<b>No rifampin (n= 18)</b>	<b>Odd-Ratio (CI 95%)</b>	<b>P Value</b>
<b>Mortality</b>				
In-hospital mortality	9 (23.7)	3 (16.7)	1.55 (0.36-6.60)	.80
One-month mortality	10 (26.3)	5 (27.8)	0.93 (0.26-3.27)	.84
Three-month mortality	12 (31.6)	5 (27.8)	1.20 (0.35-4.14)	.98
Six-month mortality	14 (36.8)	5 (27.8)	1.51 (0.45-5.16)	.71
One-year mortality	15 (39.5)	7 (38.9)	1.03 (0.32-3.24)	.80
<b>Relapse</b>	3 (7.9)	2 (11.1)	0.68 (0.10-4.52)	.68
Vitamin K antagonist imbalance	5 (41.7)	2 (22.2)	1.97 (0.83-46.59)	.16
Bleeding complication	5 (13.2)	0 (0)	6.08 (0.32-116.2)	.27
Length of stay, days	42.2 ± 16.7	34.6 ± 14.7	-	.013

Quantitative variables are expressed as mean +/- standard deviation, qualitative variables are expressed by numbers (%)

**Supplementary Table 4 Univariate analysis of outcomes in 128 patients with staphylococci prosthetic valve endocarditis without valve surgery**

<b>Variable</b>	<b>Total (n=128)</b>	<b>Rifampin- based combination (n=66)</b>	<b>No rifampin (n=62)</b>	<b>Odds-Ratio (CI 95%)</b>	<b>P Value</b>
<b>Mortality</b>					
In-hospital mortality	29 (22.7)	18 (27.3)	11 (17.7)	1.73 (0.74-4.06)	.28
One-month mortality	37 (28.9)	21 (31.8)	16 (25.8)	1.34 (0.62-2.90)	.58
Three-month mortality	39 (30.5)	23 (34.8)	16 (25.8)	1.54 (0.72-3.29)	.36
Six-month mortality	44 (34.4)	27 (40.9)	17 (27.4)	1.83 (0.87-3.85)	.16
One-year mortality	49 (38.3)	29 (43.9)	20 (32.3)	1.65 (0.80-3.39)	.24
Relapse	12 (9.4)	6 (9.0)	6 (9.7)	0.93 (0.28-3.07)	.93
Vitamin K antagonist imbalance during endocarditis	16 (12.5)	11 (16.7)	5 (8.6)	2.3 (0.74-7.00)	.23
Bleeding complication	15 (11.7)	7 (10.6)	8 (12.9)	0.80 (0.27-2.36)	.90
Length of stay, days	35.2 ± 17.1	39.7 ± 18.9	30.4 ± 13.6	-	.007

Quantitative variables are expressed as mean +/- standard deviation, qualitative variables are expressed by numbers (%)

**LE BOT, Audrey – La rifampicine améliore-t-elle le pronostic des endocardites à *Staphylococcus* sp. sur prothèse ? Une étude rétrospective multicentrique**

36 feuilles, 0 illustration, 1 graphique, 4 tableaux, 30cm – Thèse : Médecine ; Rennes 1 ; 2020 ; N°

**Introduction** Les sociétés savantes recommandent la prescription de rifampicine en association avec une antibiothérapie antistaphylococcique par pénicilline ou glycopeptides dans le traitement des endocardites infectieuses (EI) à *Staphylococcus* sp. sur prothèse valvulaire (PV). Néanmoins les données cliniques concernant son intérêt dans cette indication sont limitées.

**Matériels et méthodes.** Etude de cohorte multicentrique rétrospective conduite dans trois CHU français, concernant les adultes hospitalisés pour la prise en charge d'une EI sur PV à *Staphylococcus* sp. Le critère de jugement principal était la mortalité à un an.

**Résultats.** Cent quatre-vingts patients ont été inclus : 101 (56,1%) ont reçu un traitement comprenant de la rifampicine, et 79 (43,9%) ont été traités sans rifampicine. La mortalité à un an ne différait pas entre les événements traités avec ou sans rifampicine (37,6% vs 31,6%,  $p=0,62$ ), y compris après analyse multivariée. Il n'y avait pas non plus de différence en termes de rechutes (5,9% vs 8,9%,  $p=0,65$ ). Les épisodes traités avec de la rifampicine avaient une durée d'hospitalisation plus longue de 11 jours ( $p<0,0001$ ), mais il n'y avait pas plus d'évènements indésirables ou de déséquilibre des AVK retrouvés.

**Conclusion.** Près de 44% des épisodes ont été traités sans rifampicine. La survie à un an et les taux de rechutes étaient similaires dans les 2 groupes. Ces résultats permettent d'apporter des éléments rassurants en cas de difficultés associées à la prescription de rifampicine : son absence dans l'arsenal thérapeutique des EI sur PV ne semble pas associée à un pronostic péjoratif.

Rubrique de classement : MALADIES INFECTIEUSES

Mots-clés : Endocardite bactérienne, prothèse valvulaire cardiaque, mortalité, *Staphylococcus*, rifampicine, rechute

Mots-clés anglais MeSH Bacterial endocarditis, heart valve prosthesis, *Staphylococcus*, rifampin, mortality, relapse

Président:	Monsieur le Professeur Pierre TATTEVIN
Assesseurs:	Monsieur le Professeur Matthieu REVEST (directeur de thèse)
JURY	Monsieur le Professeur Yves LE TULZO
	Monsieur le Professeur David BOUTOILLE
	Madame le Professeure Séverine ANSART
	Madame le Docteur Rozenn LE BERRE, Maître de Conférences des Universités