



HAL
open science

Predicting symptomatic intracranial haemorrhage in acute ischemic stroke: introduction to a clinical-radiological and mri-based machine learning approach

Florent Lépilliet

► **To cite this version:**

Florent Lépilliet. Predicting symptomatic intracranial haemorrhage in acute ischemic stroke: introduction to a clinical-radiological and mri-based machine learning approach. Human health and pathology. 2021. dumas-03468513

HAL Id: dumas-03468513

<https://dumas.ccsd.cnrs.fr/dumas-03468513>

Submitted on 7 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AVERTISSEMENT

Ce document est le fruit d'un long travail approuvé par le jury de soutenance.

La propriété intellectuelle du document reste entièrement celle du ou des auteurs. Les utilisateurs doivent respecter le droit d'auteur selon la législation en vigueur, et sont soumis aux règles habituelles du bon usage, comme pour les publications sur papier : respect des travaux originaux, citation, interdiction du pillage intellectuel, etc.

Il est mis à disposition de toute personne intéressée par l'intermédiaire de [l'archive ouverte DUMAS](#) (Dépôt Universitaire de Mémoires Après Soutenance).

Si vous désirez contacter son ou ses auteurs, nous vous invitons à consulter la page de DUMAS présentant le document. Si l'auteur l'a autorisé, son adresse mail apparaîtra lorsque vous cliquerez sur le bouton « Détails » (à droite du nom).

Dans le cas contraire, vous pouvez consulter en ligne les annuaires de l'ordre des médecins, des pharmaciens et des sages-femmes.

Contact à la Bibliothèque universitaire de Médecine
Pharmacie de Grenoble :
bump-theses@univ-grenoble-alpes.fr

Année: 2021

**PREDICTING SYMPTOMATIC INTRACRANIAL HAEMORRHAGE IN ACUTE
ISCHEMIC STROKE: INTRODUCTION TO A CLINICAL-RADIOLOGICAL AND
MRI-BASED MACHINE LEARNING APPROACH**

THÈSE
PRÉSENTÉE POUR L'OBTENTION DU TITRE DE DOCTEUR EN MÉDECINE

DIPLÔME D'ÉTAT

Florent LEPILLIET

[Données à caractère personnel]

THÈSE SOUTENUE PUBLIQUEMENT À LA FACULTÉ DE MÉDECINE DE
GRENOBLE

Le : 26/02/2021

DEVANT LE JURY COMPOSÉ DE

Président du jury :

M. le Professeur Olivier DETANTE

Membres :

M. le Professeur Alexandre KRAINIK

M. le Professeur Mikaël MAZIGHI

M. Thomas CHRISTEN

M. le Docteur Jérémie PAPASSIN (directeur de thèse)

*L'UFR de Médecine de Grenoble n'entend donner aucune approbation ni improbation aux opinions émises dans les thèses ;
ces opinions sont considérées comme propres à leurs auteurs.*

Doyen de la Faculté : Pr. Patrice MORAND

Année 2020-2021

ENSEIGNANTS DE L'UFR DE MEDECINE

CORPS	NOM-PRENUM	Discipline universitaire
PU-PH	ALBALADEJO Pierre	Anesthésiologie-réanimation et médecine péri-opératoire
PU-PH	APTEL Florent	Ophthalmologie
PU-PH	ARVIEUX-BARTHELEMY Catherine	Chirurgie viscérale et digestive
PU-PH	BAILLET Athan	Rhumatologie
PU-PH	BARONE-ROCHETTE Gilles	Cardiologie
PU-PH	BAYAT Sam	Physiologie
MCF Ass.MG	BENDAMENE Farouk	Médecine Générale
PU-PH	BENHAMOU Pierre-Yves	Endocrinologie, diabète et maladies métaboliques
PU-PH	BERGER François	Biologie cellulaire
MCU-PH	BIDART-COUTTON Marie	Biologie cellulaire
PU-PH	BLAISE Sophie	Chirurgie vasculaire ; médecine vasculaire
PR Ass. Méd.	BOILLOT Bernard	
MCU-PH	BOISSET Sandrine	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	BONAZ Bruno	Gastroentérologie ; hépatologie ; addictologie
PU-PH	BONNETERRE Vincent	Médecine et santé au travail
PU-PH	BOREL Anne-Laure	Nutrition
PU-PH	BOSSON Jean-Luc	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	BOTTARI Serge	Biologie cellulaire
PR Ass.MG	BOUCHAUD Jacques	Médecine Générale
PU-PH	BOUGEROL Thierry	Psychiatrie d'adultes
PU-PH	BOUILLET Laurence	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
MCU-PH	BOUSSAT Bastien	Epidémiologie, économie de la santé et prévention
PU-PH	BOUZAT Pierre	Anesthésiologie-réanimation et médecine péri-opératoire
PU-PH émérite	BRAMBILLA Christian	Pneumologie
PU-PH émérite	BRAMBILLA Elisabeth	Anatomie et cytologie pathologiques
MCU-PH	BRENIER-PINCHART Marie Pierre	Parasitologie et mycologie
PU-PH	BRICAULT Ivan	Radiologie et imagerie médicale
PU-PH	BRICHON Pierre-Yves	Chirurgie thoracique et cardiovasculaire
MCU-PH	BRIOT Raphaël	Thérapeutique-médecine de la douleur ; Addictologie
PU-PH émérite	CAHN Jean-Yves	Hématologie
PU-PH émérite	CARPENTIER Patrick	Chirurgie vasculaire, médecine vasculaire
PR Ass.MG	CARRILLO Yannick	Médecine Générale
PU-PH	CESBRON Jean-Yves	Immunologie
PU-PH	CHABARDES Stephan	Neurochirurgie
PU-PH	CHABRE Olivier	Endocrinologie, diabète et maladies métaboliques
PU-PH	CHAFFANJON Philippe	Anatomie
MCF Ass.MG	CHAMBOREDON Benoît	Médecine Générale
PU-PH	CHARLES Julie	Dermato-vénérologie
MCF Ass.MG	CHAUVET Marion	Médecine Générale
PU-PH	CHAVANON Olivier	Chirurgie thoracique et cardiovasculaire

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH	CHIQUET Christophe	Ophtalmologie
PU-PH	CHIRICA Mircea	Chirurgie viscérale et digestive
PU-PH	CINQUIN Philippe	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	CLAVARINO Giovanna	Immunologie
PU-PH	COHEN Olivier	Histologie, embryologie et cytogénétique
PU-PH	COURVOISIER Aurélien	Chirurgie infantile
PU-PH	COUTTON Charles	Génétique
PU-PH	COUTURIER Pascal	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PU-PH	CRACOWSKI Jean-Luc	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PU-PH	DEBATY Guillaume	Médecine d'Urgence
PU-PH	DEBILLON Thierry	Pédiatrie
PU-PH	DECAENS Thomas	Gastroentérologie ; hépatologie ; addictologie
PR Ass. Méd.	DEFAYE Pascal	Cardiologie
PU-PH	DEGANO Bruno	Pneumologie ; addictologie
PU-PH	DEMATTEIS Maurice	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PU-PH émérite	DEMONGEOT Jacques	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	DERANSART Colin	Physiologie
PU-PH	DESCOTES Jean-Luc	Urologie
PU-PH	DETANTE Olivier	Neurologie
MCU-PH	DIETERICH Klaus	Génétique
MCU-PH	DOUTRELEAU Stéphane	Physiologie
MCU-PH	DUMESTRE-PERARD Chantal	Immunologie
PU-PH	EPAULARD Olivier	Maladies infectieuses ; Maladies tropicales
PU-PH	ESTEVE François	Biophysique et médecine nucléaire
MCU-PH	EYSSERIC Hélène	Médecine légale et droit de la santé
PU-PH	FAUCHERON Jean-Luc	Chirurgie viscérale et digestive
MCU-PH	FAURE Julien	Biochimie et biologie moléculaire
PU-PH	FERRETTI Gilbert	Radiologie et imagerie médicale
PU-PH	FEUERSTEIN Claude	Physiologie
PU-PH	FONTAINE Éric	Nutrition
PU-PH	FRANCOIS Patrice	Epidémiologie, économie de la santé et prévention
MCU-MG	GABOREAU Yoann	Médecine Générale
PU-PH	GARBAN Frédéric	Hématologie ; Transfusion
PU-PH	GAUDIN Philippe	Rhumatologie
PU-PH	GAVAZZI Gaétan	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PU-PH	GAY Emmanuel	Neurochirurgie
MCU-PH	GILLOIS Pierre	Biostatistiques, informatique médicale et technologies de communication
PU-PH	GIOT Jean-Philippe	Chirurgie plastique, reconstructrice et esthétique ; Brûlologie
MCU-PH	GRAND Sylvie	Radiologie et imagerie médicale
PU-PH émérite	GRIFFET Jacques	Chirurgie infantile
MCU-PH	GUZUN Rita	Nutrition
PU-PH	HAINAUT Pierre	Biochimie et biologie moléculaire
PU-PH émérite	HALIMI Serge	Nutrition
PU-PH	HENNEBICQ Sylviane	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PU-PH	HOFFMANN Pascale	Gynécologie-obstétrique ; gynécologie médicale
PU-PH émérite	HOMMEL Marc	Neurologie
PU-MG	IMBERT Patrick	Médecine Générale

CORPS	NOM-PRENUM	Discipline universitaire
PU-PH émérite	JOUK Pierre-Simon	Génétique
PU-PH	KAHANE Philippe	Physiologie
MCU-PH	KASTLER Adrian	Radiologie et imagerie médicale
PU-PH	KRAINIK Alexandre	Radiologie et imagerie médicale
PU-PH	LABARERE José	Epidémiologie, économie de la santé et prévention
MCU-PH	LABLANCHE Sandrine	Endocrinologie, diabète et maladies métaboliques
MCU-PH	LANDELLE Caroline	Bactériologie – virologie ; Hygiène hospitalière
PU-PH	LANTUEJOL Sylvie	Anatomie et cytologie pathologiques
PR Ass. Méd.	LARAMAS Mathieu	Cancérologie ; radiothérapie
MCU-PH	LARDY Bernard	Biochimie et biologie moléculaire
MCU - PH	LE GOUELLEC LE PISSART Audrey	Biochimie et biologie moléculaire
PU-PH	LECCIA Marie-Thérèse	Dermato-vénérologie
MCF Ass.MG	LEDoux Jean-Nicolas	Médecine Générale
PU-PH émérite	LETOUBLON Christian	Chirurgie viscérale et digestive
PU-PH	LEVY Patrick	Physiologie
PU-PH	LONG Jean-Alexandre	Urologie
MCU-PH	LUPO Julien	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	MAIGNAN Maxime	Médecine d'urgence
PU-PH	MAITRE Anne	Médecine et santé au travail
MCU-PH	MALLARET Marie-Reine	Bactériologie – virologie ; Hygiène hospitalière
MCU-PH	MARLU Raphaël	Hématologie ; Transfusion
PR Ass. Méd.	MATHIEU Nicolas	Gastroentérologie ; hépatologie ; addictologie
MCU-PH	MAUBON Danièle	Parasitologie et mycologie
PU-PH	MAURIN Max	Bactériologie-virologie ; Hygiène hospitalière
MCU-PH	MC LEER Anne	Histologie, embryologie et cytogénétique
MCU-PH	MONDET Julie	Histologie, embryologie et cytogénétique
PU-PH	MORAND Patrice	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	MOREAU-GAUDRY Alexandre	Biostatistiques, informatique médicale et technologies de communication
PU-PH	MORO Elena	Neurologie
PU-PH	MORO-SIBILOT Denis	Pneumologie ; addictologie
MCU-PH	MORTAMET Guillaume	Pédiatrie
PU-PH	MOUSSEAU Mireille	Cancérologie ; radiothérapie
PU-PH émérite	MOUTET François	Chirurgie plastique, reconstructrice et esthétique ; brûlologie
MCF Ass.MG	ODDOU Christel	Médecine Générale
PR Ass. Méd.	ORMEZANO Olivier	Cardiologie
MCU-PH	PACLET Marie-Hélène	Biochimie et biologie moléculaire
PU-PH	PAILHE Régis	Chirurgie orthopédique et traumatologie
PU-PH	PALOMBI Olivier	Anatomie
PU-PH	PARK Sophie	Hématologie ; Transfusion
PU-PH	PASSAGGIA Jean-Guy	Anatomie
PR Ass.MG	PAUMIER-DESBRIERES Françoise	Médecine Générale
PU-PH	PAYEN DE LA GARANDERIE Jean-François	Anesthésiologie-réanimation et médecine péri-opératoire
MCU-PH	PAYSANT François	Médecine légale et droit de la santé
MCU-PH	PELLETIER Laurent	Biologie cellulaire
PU-PH	PELLOUX Hervé	Parasitologie et mycologie
PU-PH	PEPIN Jean-Louis	Physiologie
PU-PH	PERENNOU Dominique	Médecine physique et de réadaptation

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH	PERNOD Gilles	Chirurgie vasculaire ; Médecine vasculaire
PU-PH	PIOLAT Christian	Chirurgie infantile
PU-PH	PISON Christophe	Pneumologie ; Addictologie
PU-PH	PLANTAZ Dominique	Pédiatrie
PU-PH	POIGNARD Pascal	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	POLACK Benoît	Hématologie ; Transfusion
PU-PH	POLOSAN Mircea	Psychiatrie d'adultes ; Addictologie
PU-PH	RAMBEAUD Jean-Jacques	Urologie
PU-PH	RAY Pierre	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PR Ass. Méd.	RECHE Fabian	Chirurgie viscérale et digestive
MCU-PH	RENDU John	Biochimie et biologie moléculaire
MCU-PH émérite	RIALLE Vincent	Biostatistiques, informatique médicale et technologies de communication
PU-PH	RIETHMULLER Didier	Gynécologie-obstétrique ; gynécologie médicale
PU-PH	RIGHINI Christian	Oto-rhino-laryngologie
PU-PH émérite	ROMANET Jean Paul	Ophtalmologie
PU-PH	ROSTAING Lionel	Néphrologie
PU-PH	ROUSTIT Matthieu	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
MCU-PH	ROUX-BUISSON Nathalie	Biochimie et biologie moléculaire
PR Ass.MG	ROYER DE VERICOURT Guillaume	Médecine Générale
PU-PH émérite	SARAGAGLIA Dominique	Chirurgie orthopédique et traumatologie
MCU-PH	SATRE Véronique	Génétique
PU-PH	SAUDOU Frédéric	Biologie cellulaire
PU-PH	SCHMERBER Sébastien	Oto-rhino-laryngologie
PU-PH	SCHWEBEL Carole	Médecine intensive-réanimation
PU-PH	SCOLAN Virginie	Médecine légale et droit de la santé
MCU-PH	SEIGNEURIN Amaud	Epidémiologie, économie de la santé et prévention
PU-PH émérite	STAHL Jean-Paul	Maladies infectieuses ; Maladies tropicales
PU-PH	STANKE Françoise	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
MCU-PH	STASIA Marie-José	Biochimie et biologie moléculaire
PU-PH	STURM Nathalie	Anatomie et cytologie pathologiques
PU-PH	TAMISIER Renaud	Physiologie
PU-PH	TERZI Nicolas	Médecine intensive-réanimation
PU-PH	THEVENON Julien	Génétique
MCU-PH	TOFFART Anne-Claire	Pneumologie ; Addictologie
PU-PH	TONETTI Jérôme	Chirurgie orthopédique et traumatologie
PU-PH	TOUSSAINT Bertrand	Biochimie et biologie moléculaire
PU-PH	VALMARY-DEGANO Séverine	Anatomie et cytologie pathologiques
PU-PH	VANZETTO Gérald	Cardiologie
PU-PH	VUILLEZ Jean-Philippe	Biophysique et médecine nucléaire
PU-PH	WEIL Georges	Epidémiologie, économie de la santé et prévention
PU-PH	ZAQUI Philippe	Néphrologie
PU-PH émérite	ZARSKI Jean-Pierre	Gastroentérologie ; hépatologie ; addictologie

PU-PH : Professeur des Universités - Praticiens Hospitaliers
MCU-PH : Maître de Conférences des Universités - Praticiens Hospitaliers
PU-MG : Professeur des Universités de Médecine Générale
MCU-MG : Maître de Conférences des Universités de Médecine Générale
PR Ass. Méd. : Professeur des Universités Associé de Médecine
PR Ass.MG : Professeur des Universités Associé de Médecine Générale
MCF Ass.MG : Maître de Conférences Associé de Médecine Générale

Remerciements

Aux membres du jury

Au **Professeur Olivier Detante**, merci de m'avoir fait l'honneur de présider ce jury et pour la relecture attentive de ce travail. Mais avant tout merci Olivier pour ta pédagogie unanimement reconnue, ton accessibilité si précieuse, ton investissement et ton attention portée à nos projets professionnels et personnels.

Au **Professeur Mikaël Mazighi**, je suis honoré que vous ayez accepté de faire partie de ce jury depuis la capitale. Merci pour cette belle expérience parisienne que vous m'avez permis de réaliser, votre pédagogie de qualité, votre écoute attentive et votre expérience inégalable m'ont été précieux.

Au **Professeur Alexandre Krainik**, chef du service de neuroradiologie du CHU de Grenoble. Merci de m'avoir fait l'honneur de participer à ce jury pour apporter votre expertise à ces travaux.

À **Monsieur Thomas Christen**, ingénieur-chercheur hors pair et spécialiste de l'intelligence artificielle, merci de m'avoir intégré à vos équipes et de m'avoir guidé dans la réalisation de ce projet. Merci d'avoir accepté de participer à ce jury.

Au **Docteur Jérémie Papassin**, neurologue hyperactif et papa comblé, merci de m'avoir guidé au cours de ce travail dans les méandres de la recherche médicale. Merci pour ta disponibilité et tes relectures attentives. Peut-être qu'un jour je te montrerai la convivialité du stade Bollaert comme tu m'as fait découvrir l'ambiance survoltée noir et jaune du Phare Chambérien.

À tous ceux qui ont participé à ce projet

À **Benjamin Lemasson** et **Stenzel Cackowski** sans qui ce projet n'existerait pas, merci pour votre implication et votre accessibilité.

À **Lucie Miquel** merci pour son aide précieuse au recueil des données d'imagerie.

À **Clément Brossard** merci pour son aide au maniement de Shanoir et à la conversion des données.

À mes collègues et amis

Aux médecins qui ont jalonné jusqu'ici mon parcours,

aux Drs **Delphine Lamotte** et **Sylvie Cantalloube** pour leurs notions de rééducation neurologique. Aux Drs **Katia Garambois** et **Isabelle Favre-Wiki** pour leurs enseignements en neurovasculaire et leur bienveillance. Au Dr **Mathieu Vaillant** pour sa pédagogie en douceur et ses notions de neuro-ophtalmologie. Au Dr **Olivier Casez** pour la formation en pathologies inflammatoires qu'il nous apporte, mais aussi pour ses bons mots toujours savoureux et ses recettes de bières artisanales. Au Dr **Gérard Besson** pour ses notions en maladies génétiques et métaboliques et les quelques notes de basson entendues. Au Dr **Catalina Coclitu**, aux qualités professionnelles rares malgré des goûts musicaux passables (O-Zone c'est démodé Catalina), merci pour ton aide précieuse et ta pédagogie. Aux Drs **Martial Mallaret** et **Emmeline Lagrange** pour m'avoir éclairci le chemin sinueux des neuropathies

périphériques et de l'ENMG par leurs qualités pédagogiques et leur patience. Aux **Drs Laurent Vercueil** et **Cécile Sabourdy** pour avoir réussi à rendre l'EEG ludique et pour leurs topos rapides sur tableau blanc ou leurs « instants culture générale » entre deux relectures. Au Dr **Alexa Garros** pour m'avoir initié à la neuro-pédiatrie et pour cet apéro de fin de stage au cours duquel on a découvert le vrai visage d'Aline. Au Pr **Elena Moro**, au Pr **Philippe Kahane**, aux Drs **Lorella Minotti**, **Valérie Fraix**, **Anna Castrioto** et **Mathilde Sauvée** pour leurs avis éclairés et leur bienveillance. Merci au Dr **Pauline Cuisenier** pour tout ce qu'elle m'a appris pendant cette année cumulée en neurologie vasculaire et générale, sa bienveillance et son soutien sur le plan professionnel comme personnel (promis mes articulations vont cesser d'avoir besoin d'être imagées).

Aux équipes médicales et paramédicales des services traversés jusqu'ici pour leur accueil bienveillant et complice : l'équipe de MPR neuro de Reine Hortense, l'équipe de neurovasculaire du CHU, l'équipe d'HDJ neurologique du CHU, l'équipe de neurologie générale du CHU, l'équipe des Explorations Fonctionnelles du Système Nerveux (EFSN) du CHU. À toute l'équipe de Neuroradiologie Interventionnelle de la Fondation Rothschild pour leur accueil, leur pédagogie et les bons moments passés au bloc ou en terrasse.

À mes co-internes qui m'accompagnent dans ces longues études,

Gauthier le pyromane de la piste et amateur de bonnes choses. On partage l'amour du Nord et beaucoup de séances de kiné (mais tu as quand même un ligament de retard pour être son préféré). **Loïc** mon « jumeau maléfique », partenaire de soirée, de soccer ou de montagne bref un ami en or... jusqu'à ce que vous l'affrontiez au babyfoot. **Inès** ma co-interne de toujours, merci pour ta motivation perpétuelle et pour ces fous rires quotidiens qui ont rythmé nos journées même dans la galère. **Thomas** le renard des surfaces, qui ne distille le cynisme que pour masquer son cœur d'artichaut. **Hugo** dont je me dois pour la bienséance de censurer ici le surnom, cyclotouriste aguerri, fêtard dans l'âme et la bonne humeur incarnée ! **Lucie**, la bienveillance et la classe à l'état pure, ton émerveillement perpétuel fait toujours plaisir à voir. **Pomme** la pétillante, un jour promis on arrêtera de faire des vanes sur ton prénom. **Christo(phe)** notre délégué désigné, toujours prompt à défendre la veuve et l'orphelin quand ils sont internes de neurologie. **Célia** l'auvergnate, ce fanatisme pour le poulet m'étonnera toujours. À **Giovanni** le charismatique, merci pour les bons plans professionnels et musicaux. À **Hélène**, jeune maman et jeune assistante, qui nous a mis un sacré coup de vieux ! À **Sarah** la spécialiste des rêves, n'oublie pas de sortir des escaliers de temps en temps. À **Guillaume** de retour du pays des caribous, notre Samantha Oups à nous. À **Alex**, **Sylvain** et **Annaëlle** qui assurent déjà pleinement la relève. À **Céline** qui m'a supporté pendant toute la période de confinement et avec qui j'ai perdu deux ou trois points d'acuité visuelle à m'arracher les yeux sur les tracés (« onde lente à front raide ? »). À mon co-interne d'UNV **Louis**, pour sa rigueur scientifique mais surtout son humour noir qui nous a valu tous ces fous rires.

A mes co-internes parisiens de la FOR **Erwan**, **Ihsan**, **Félix**, **Firas**, **Zakaria** pour ces bons moments de franche camaraderie dans l'euphorie comme dans la galère et mention particulière à **Thomas** mon compagnon de « province ».

À mes amis

Aux « Yolos Gorgous » qui ont égailé mon externat et continuent de le faire aujourd'hui. **Vincent**, **Louis**, **Victor**, **Robin**, **Justin**, **Pierro**, **Bertille**, **Camille**, **Sixtine**, **Agathe**, **Raphaëlle**, **Pia**, **Laura**, **Diane**. Merci à tous pour le nombre incalculable de souvenirs tissés ensemble depuis l'intégration jusqu'aux vacances en Corse ou à l'autre bout du monde ! Nos retrouvailles régulières sont toujours un grand bol d'air frais et vous m'êtes tous très chers !

À **Alex, Tristan et Quentin** le trio infernal, merci pour ces fous rires mémorables qui ne cessent jamais !
À **Etienne** devenu ingénieur germanophone (enfin presque), vivement qu'on puisse se la faire cette Oktoberfest ! À **Damien et Baptiste** les copains d'enfance, du primaire à l'adolescence merci pour ces souvenirs irremplaçables. À **Antoine** pour ces moments inoubliables aux quatre coins du globe. À **Marine**, ma binôme de galère en P1, merci pour ces tonnes de souvenirs entre les voyages scolaires, les premières soirées, les vacances aux Canebières et bien d'autres encore à écrire. À **Julie et Hélène, Jennifer et Louise** pour ces bons moments partagés en vacances, en week-ends, sur les bancs de l'école pour les unes ou jusqu'à la faculté pour les autres.

Aux « Grelous » **Alix et François, Capu et Alex, Charline et Benji, Marion et Seb, Eve et Thomas, Raphaëlle et Thibaut** merci pour ces soirées et ces week-ends riches en émotions (et en karaoké) que l'on ne se lasse pas de partager ! A **Flavie et Guillaume, Alice et Hector, Eva et Ben, Lison et Liam** merci pour toutes ces randos et week-ends en Ardèche mémorables !

À ma famille

À mes parents pour l'éducation que vous nous avez fournie et les valeurs de curiosité et d'humilité que vous continuez de nous inculquer. Merci **Maman** pour ton soutien infaillible depuis toujours, tu nous as appris à faire les choses dans le respect de l'autre et avec le sérieux nécessaire tout en gardant du recul. Merci de nous avoir fait aimer les belles lettres quand papa nous berçait de chiffres ! Merci **Papa** pour ton soutien sans faille et ta disponibilité, toujours là pour nous aider à faire les bons choix. Merci pour toutes ces heures passées sur les DM de maths mais surtout merci de m'avoir appris à ne pas trop se prendre au sérieux et à garder le sens de l'humour ! Ces quelques lignes ne suffiront pas pour vous exprimer ma gratitude.

À ma grande sœur **Émilie**, qui continue de veiller sur ses petits frères comme au premier jour. Merci d'être toujours là pour moi malgré la vie de famille et professionnelle qui t'accapare maintenant, et bravo pour les défis que tu as relevés ces dernières années !

À mon grand frère **Thibaut** qui a réussi à infiltrer le monde secret du sport automobile. On continue de se chamailler comme des enfants à chaque repas de famille mais malgré tout on s'est bien assagi ! Merci de ton écoute et de ton soutien.

À **Elodie et Julien** qui partagent la vie des sus-cités, je suis très heureux que vous ayez intégré la fratrie !
À mes filleules **Louise et Lucie** qui me rendent gaga, à mon neveu **Axel** « le président » et au petit dernier à venir.

À **mes grands-parents** Gisèle, Marie-Thérèse, René et Ernest tous partis trop tôt.

Aux **Lernould** pour m'avoir intégré à leur fratrie déjà bien fournie !

À **Cassandra** évidemment. Merci pour ton insouciance et ta douceur au quotidien. Merci de me faire toujours autant rire. Notre complémentarité m'est devenue indispensable et j'aurai toujours besoin de toi pour faire mes choix. Hâte de parcourir les fjords norvégiens à tes côtés.

Table of contents

Remerciements	6
Table of contents	9
Abbreviations	10
Résumé / Abstract :	11
Table and Figures	12
1. Introduction	13
1.1. Pathophysiology and risk factors	13
1.2. Prediction scores	14
1.3. About machine learning	15
1.4. Objectives	20
2. SUBJECTS AND METHODS	20
2.1. Study population	20
2.2. Clinical data collection	21
2.3. Imaging	21
2.4. Machine learning algorithm	23
2.5. Statistical analysis	24
3. RESULTS	25
3.1. Baseline characteristics	25
3.2. Radiological and Clinical outcomes	27
3.3. sICH characteristics: univariate and multivariate analyses	29
3.4. Symptomatic intracranial haemorrhage prediction	31
3.4.1. THRIVE score performance	31
3.4.2. DWI-ASPECTS performance	31
3.4.3. Machine Learning (ML) algorithm	32
4. Discussion	34
4.1. sICH outcomes	34
4.2. Predictive performances	35
4.3. Study limitations and future directions	37
5. Conclusion	40
References	42

Abbreviations

AF – Atrial Fibrillation

AIS - Acute Ischemic Stroke

CT - Computerized Tomography

DM – Diabetes Mellitus

HBP – High Blood Pressure

HI – Haemorrhagic Infarction

HT – Haemorrhagic Transformation

ICH – Intracranial Haemorrhage

IVT – Intravenous Thrombolysis

ML – Machine Learning

MRI - Magnetic Resonance Imaging

mRS – Modified Rankin Scale

MT – Mechanical Thrombectomy

NIHSS - National Institute of Health Stroke Scale

OTT – Onset-to-Treatment Time

PH – Parenchymal Hematoma

sICH – Symptomatic Intracranial Haemorrhage

tPA – Tissue Plasminogen Activator

Résumé :

L'infarctus cérébral (IC) est l'une des principales causes de décès et d'invalidité. Les thérapies de reperfusion, thrombolyse intraveineuse (TIV) et thrombectomie mécanique (TM), améliorent le pronostic des patients victimes d'IC mais augmentent le risque de transformation hémorragique (TH) du parenchyme cérébral infarcté. La TH symptomatique (THs) est la définition la mieux corrélée à un mauvais pronostic. Plusieurs méthodes basées sur des données cliniques et/ou radiologiques ont échoué à prédire le risque de THs en pratique clinique. Nous avons élaboré un algorithme d'apprentissage machine supervisé (ML) sur des données cliniques et d'IRM au sein d'une cohorte de 100 sujets atteints d'IC de circulation antérieure traités par TIV et/ou TM. Certains ont présenté une THs (n=28) ou une TH non symptomatique (n=27) et 45 témoins sans saignement ont été appariés sur la gravité clinique et l'âge. Nous avons comparé la performance de l'algorithme ML à celles du score clinique THRIVE (Total Health Risks in Vascular Events) et au score radiologique ASPECT (Alberta Stroke Program Early CT Score) appliqué à l'imagerie IRM (DWI-ASPECTS). L'algorithme ML a prédit la THs avec une aire sous la courbe ROC (AUC) de 0,658. Les AUC estimées du score THRIVE et du DWI-ASPECTS étaient respectivement de 0,664 et 0,634. L'algorithme ML a réussi à synthétiser les données cliniques et radiologiques pour fournir une probabilité de THs équivalente à celles des scores actuels. Des travaux complémentaires sont nécessaires pour améliorer ces performances en élargissant les effectifs, intégrer des données clinico-radiologiques plus pertinentes puis valider ces résultats sur une population hétérogène.

Mots-clés : intelligence artificielle, IRM, thrombolyse, thrombectomie, transformation hémorragique

Abstract:

Acute ischemic stroke (AIS) is a leading cause of death and disability. Reperfusion therapies using intravenous thrombolysis (IVT) and mechanical thrombectomy (MT) improve functional outcomes in patients with AIS but unfortunately increase the risk of intracranial haemorrhage (ICH) of ischemic brain tissue. Symptomatic ICH (sICH) is the most relevant definition as it is most correlated with poor outcome. Several methods, based on clinical and/or radiological data, failed to predict the risk of sICH in clinical practice. We attempted a new approach by training a supervised machine learning (ML) algorithm on clinical and MRI data within a 100 subjects cohort of patients with anterior circulation AIS treated by IVT and/or MT who underwent sICH (n=28), non-symptomatic ICH (n=27) and 45 controls with no bleeding, matched on clinical severity and age. We compared ML algorithm accuracy to the performance of the clinical Total Health Risks in Vascular Events (THRIVE) score and to the radiological Alberta Stroke Program Early CT Score applied to MR imaging (DWI-ASPECTS). ML algorithm predicted sICH with an Area Under receiver operating characteristic Curve (AUC) of 0.658. Estimated AUC of THRIVE score and DWI-ASPECTS were 0.664 and 0.634, respectively. Although it does not outperform current tools, ML algorithm was able to synthesize clinical and radiological data to provide a sICH probability. Further studies are needed to improve these performances. Enlarging dataset and integrating more relevant clinical and radiological data should be the first avenues for improvement. Validation of these results on a more heterogeneous external population is also warranted.

Keys words: machine learning, deep learning, MRI, thrombolysis, thrombectomy, symptomatic haemorrhagic transformation

Table and Figures

TABLE 1: BASELINE CHARACTERISTICS ACCORDING TO EACH SAMPLE.....	26
TABLE 2: RADIOLOGICAL AND CLINICAL OUTCOMES REGARDING SICH AND NON-SYMPTOMATIC ICH SUBJECTS.	27
TABLE 3: BASELINE CHARACTERISTICS AND OUTCOMES BETWEEN SICH SUBJECTS AND CONTROL GROUP, UNI AND MULTIVARIATE ANALYSES.....	30
TABLE 4: THRIVE SCORE AND DWI-ASPECTS SICH PREDICTION: SENSITIVITY (Se), SPECIFICITY (Sp), POSITIVE (PPV) AND NEGATIVE (NPV) PREDICTIVE VALUES ACCORDING TO RELEVANT CUTPOINTS.....	32
FIGURE 1: CONCEPTUAL PINCIPLES OF ANN. DEEP LEARNING: A PRIMER FOR RADIOLOGISTS. CHARTRAND ET AL. RADIOGRAPHICS 2017	17
FIGURE 2: PREDICTION LEARNING AND “LOSS” FUNCTION PRINCIPLE.....	18
FIGURE 3: OVERALL MACHINE LEARNING (ML) PROCESSING SCHEME	19
FIGURE 4: FLOW CHART	25
FIGURE 5: MODIFIED RANKIN SCALE (MRS) AT DISCHARGE ACCORDING TO RADIOLOGICAL FEATURES.....	28
FIGURE 6: RECEIVER OPERATING CURVES (ROC) AND AREA UNDER ROC CURVES (AUC) OF DWI-ASPECT SCORE (PURPLE), THRIVE SCORE (BEIGE), ARTIFICIAL NEURAL NETWORK (ANN) MASKED BRAIN (BLUE) AND ANN ALL BRAIN (GREEN) MODELS.	33

1. Introduction

Stroke is a leading cause of death and disability in adult. Ischemic stroke is the most common type of stroke with a global life-time risk of 18% (1). Reperfusion therapies using intravenous thrombolysis (IVT) and endovascular interventions such as mechanical thrombectomy (MT) are timely restricted but improve functional outcomes in patients with acute ischemic stroke (AIS) (2–4). Unfortunately, these therapeutics increase the risk of haemorrhagic transformation of ischemic brain tissue (4–6), also called intracranial haemorrhage (ICH). The symptomatic intracranial haemorrhage (sICH) is defined as a neurological deterioration concomitant to an intracranial haemorrhage after reperfusion therapies. Radiological definitions of ICH are classically based on the opposition between haemorrhagic infarction (HI), which corresponds to haemorrhagic petechiae within the ischaemic area, and intra-parenchymal haematoma (PH), which represents a haemorrhagic area with clear sides with or without a mass effect (7). Because of different definitions of sICH in randomized clinical trials and prospective registries, incidence of sICH is ranged between 2-7% after reperfusion therapies (8–10). Rates of sICH are significantly reduced after IVT using MRI-guide selection of patients, ranging from 2 to 3% (3,11,12). sICH is strongly associated with 3-months poor functional outcomes and mortality (13).

1.1. Pathophysiology and risk factors

ICH seems to be part of the natural history of ischemic infarction and could be worsen by reperfusion therapies. Numerous clinical factors have been identified as potential risk factors for ICH, such as age (6); high blood pressure (6) ; hyperglycemia (14) ; atrial fibrillation (15); clinical severity based on NIHSS (16) ; delay of treatment (17) and used of antithrombotic treatments (antiplatelet and anticoagulant agents) (18).

Meanwhile, some radiological data have been identified of interest to predict ICH. The presence of early ischemia signs on CT scan has been well identified as a risk factor for ICH (7,19). Large vessel occlusion (LVO) reflected by hyperdense middle cerebral artery sign on CT scan has also been suggested as independent predictive factor (20). Regarding MR-based imaging, which tends to become a gold standard for management of AIS, increasingly works are conducted on its contribution to ICH prediction. MR imaging provides a wide range of morphological information of ischemic lesion and adjacent parenchyma tissue which could be predictive of ICH such as apparent diffusion coefficient value (21), FLAIR hyperintensities (22), visible leukopathy in FLAIR (23), microbleeds burden visible on T2* sequence (24). But brain infarction volume on DWI appears as an independent and strongest risk factor of ICH compared to others factors (25).

Post-thrombolytic haemorrhagic transformation depends on multiple and interconnected pathological processes, including ischaemic lesions and coagulopathy. Quantifying the risk of sICH would be a valuable aid for clinicians to guide therapeutic decision.

1.2. Prediction scores

Several scores aimed to predict sICH mixing clinical and radiological data, which showed similar predictive values (26–28). These are rarely used in current practice because there is not enough confidence in their prediction to generally justify a deferral of thrombolysis when high-risk patients are often those whose prognosis would be bleakest without thrombolysis (29). None of these scores are based on initial MRI. The Total Health Risks in Vascular Events (THRIVE) score appears as the most practical and robust (30). The THRIVE score assigns 1 point for 60 to 79 years of age, 2 points for ≥ 80 years of age, 2 points for NIHSS score 11 to 20, 4 points for NIHSS score ≥ 21 , and 1 point each for hypertension, DM, and AF. The Alberta Stroke Programme Early CT Score (ASPECTS) is 10 points score originally based on early

ischaemic changes identified on CT during the first few hours after stroke onset, representing early cytotoxic oedema and possibly the development of irreversible injury. It was significantly correlated to functional outcome and predicts sICH with sensitivity and specificity values of respectively 0.90 and 0.62 (31). A threshold of 7 has been identified as most relevant (19). In routine practice, DWI-ASPECTS is used to identify patients with acute stroke and large DWI lesions, avoiding the need for measuring lesion volumes in the acute setting. The risk of sICH after IVT is inversely correlated to DWI-ASPECTS and a cut-off at 7 remained the most relevant (32).

1.3. About machine learning

Artificial Intelligence (AI) refers to computer systems able to perform tasks normally requiring human intelligence, such as visual perception, speech recognition, decision-making, and translation between languages. One of the most widely used method is machine learning (ML) whose goal is to build data analysis algorithms to extract characteristics from these data. This technique generally involves two phases. A “training phase” aimed at training the algorithm to perform a task (classification, segmentation) on a selected database. The task can be provided to the algorithm (supervised learning) or must be deduced by the algorithm itself (unsupervised). The second phase called "testing phase" aims at using the previously established model to process new unknown data.

The supervised machine learning method consists in learning a function that maps an input to an output based on a labelled training set of data, composed of input-output pairs. A supervised ML algorithm analyses the training data, learns from previous mistakes after repeated analysis, and produces an inferred function, which can be used for mapping new examples. The optimal scenario will allow for the algorithm to correctly determine the class labels for unseen instances. The most popular technique currently used is Artificial Neural Network (ANN). ANN is

inspired by biological neural networks: an artificial neuron receives inputs from other neurons, integrates inputs that are weighted and activates when a defined condition is met. This method is very flexible and applicable to a wide variety of characteristic data. Concerning imaging data, it consists of performing multiple statistical tests (regression) in each voxel without selection in order to select the most relevant characteristics.

Deep Learning is a recent refinement of neural networks that mimics the human brain using multiple layers of neural networks, interconnected in different ways depending on the type of network (Convolutional Neural Network, Recurrent Neural Network...). This was made possible thanks to the improvement of computer techniques, processing power and management of Big Data. The ability for such algorithms to manage various kinds of data (quantitative, qualitative, non-linear...) and the amount of data currently collected in our hospitals, explains the recent craze in the medical world and the exponential number of publications in the medical field referring to such techniques, mainly in the field of medical imaging (33–35). Having regard to stroke management, researchers recently attempt to improve stroke imaging using ML algorithm in various ways : automatically detection of infarcted area (36), segmentation (37), prediction of the final infarcted area (38), prediction of the clinical outcome (39).

Prediction of upcoming ICH occurrence using ML has recently provided encouraging results when based on clinical variables (40), CT images (41) or permeability parameters derived from PWI (42), some models can even be used to predict territory and extent of ICH, using a voxel to voxel analysis (43).

The algorithm implemented in this work is based on supervised learning using an ANN. In the brain, neurons exchange information via chemical and electrical synapses. Electrochemical signals are propagated from the synaptic area through the dendrites when a certain excitation threshold is reached, the cell then releases an activation signal through its axon toward synapses

with neighbouring neurons. Complex signals can be encoded by networks of neurons based on this paradigm; for instance, a hierarchy of neurons in the visual cortex is able to detect edges by combining signals from independent visual receptors. ANN are inspired by this biologic process. The basic unit of an ANN, the artificial neuron or node, is a simplified model that mimics this basic mechanism. The artificial neuron takes as an input a set of values representing features, each multiplied by a corresponding weight. The weighted features are summed and passed through a nonlinear activation function. In this way, an artificial neuron can be viewed as producing a decision by weighting a set of evidence (Figure 1).




Figure 1: Conceptual principles of ANN.
 Deep Learning: a primer for radiologists. Chartrand et al. Radiographics 2017

Thousands of neurons assembled together form a layer. Stacking these layers, connecting the output of one layer to the input of the following layer, form the global neural network architectures which can represent very complex nonlinear functions.

The first layer, called the input layer, represents input data such as here individual pixel intensities or clinical data materialized by either categorical or continuous data. The output layer produces target values such as a classification result. The intermediate layers are called hidden layers, since they do not directly produce visible desired outputs, but rather compute intermediate representations of the input features that are useful in the inference process. A neural network is trained by adjusting the parameters, which consist of the weights and biases of each node. Starting from a random initial configuration, the parameters are adjusted via an optimization algorithm which attempts to find a set of parameters that performs well on a training dataset. For each data sample, the performance of the network is assessed through a loss (error) function that quantitatively measures the inaccuracy of the prediction (Figure 2). Each parameter of the network is then adjusted by small increments in the direction that minimizes the loss, a process called back-propagation. This training procedure is performed multiple times for each sample in the training dataset, so the parameters approach values that maximize the model accuracy (44).




Figure 2: Prediction learning and "loss" function principle.

Convolutional Neuron Network (CNN) is a subclass of neural networks which have at least one convolution layer. Mathematically, a convolution is a mathematical operation which, for two input signals, produces a new signal which is a function of the two previous ones. A convolution unit receives its input from multiple units from the previous layer which together create a proximity. Therefore, the input units (that form a small neighbourhood) share their weights. The convolution units are especially beneficial as they reduce the number of units in the network (since they are many-to-one mappings). This means, there are fewer parameters to learn which reduces the chance of overfitting as the model would be less complex than a fully connected network. Moreover, they consider the context/shared information in the small neighbourhoods. This feature is very important in many applications such as image, video, text, and speech processing/mining as the neighbouring inputs (eg pixels, frames, words, etc). That is why this subclass of ANN is widely used in medical imaging field to extract features from the image by groups of pixels and finally simplify the image into a single set of features which can then be integrated in one or several last “fully connected” linear layers (Figure 3).




Figure 3: Overall machine learning (ML) processing scheme

A particular CNN subtype Residual Network algorithm was used here. Residual Networks (also called ResNet) are built to use “skip connections”, or “shortcuts” to jump over some layers. This kind of ANN have empirically shown to increase classification performance by simplifying the network, especially when increasing the number of layers.

1.4. Objectives

We aim to (1) evaluate a clinical-radiological MRI-based ML algorithm to predict the occurrence of a sICH and (2) compare ML algorithm accuracy to validated clinical and radiological scores performances.

2. SUBJECTS AND METHODS

2.1. Study population

Our study retrospectively analysed data from a prospective stroke registry of individual data (Northern French Alps Emergency Network, RENAUS Stroke database). We selected all AIS patients referred to Grenoble University Hospital (CHUGA) and Chambéry Hospital (CHMS) between January 2014 and December 2018 with a sICH after a reperfusion therapies which was IVT, MT or “bridging” which is a combination of IVT then MT. We matched controls with non-symptomatic ICH (1:1) and controls without ICH (1:2) on age and initial stroke severity based NIHSS score.

Symptomatic ICH was defined, according to the ECASS II definition, by an increase in the NIHSS of more than 3 points concomitant with the appearance of haemorrhagic infarction or

intra-parenchymal haematoma on the control imaging which can be either CT or MR scanner carried out within 36h, or death due to this haemorrhagic transformation.

All patients treated with IVT received ALTEPLASE with a 0.09 mg/kg bolus followed by 0.81 mg/kg infusion in one hour.

Exclusion criteria were no pre-treatment imaging available; no control imaging available; posterior circulation stroke; MT failure in patient with contraindication of IVT.

2.2. Clinical data collection

For each patient, the following data were extracted from computerized medical file and collected in a prospective registry: National Institutes of Health Stroke Scale (NIHSS) at admission (0 to 42, the severity increasing proportionally to the score); Onset-to-Treatment Time (OTT) defined as delay between stroke recognition and IVT administration or groin puncture in case of MT; medical history of high blood pressure, diabetes mellitus or atrial fibrillation; prior use of antithrombotic treatment; lateralization of the lesion; LVO as defined by intracranial carotid or first segment of middle-cerebral-artery occlusion; acute treatment carried out; modified Rankin Scale (mRS) score at discharge. Functional independence at discharge was defined as a mRS between 0 and 2.

2.3. Imaging

We only included MRI-based AIS patients. Each subject underwent a MR imaging routine protocol including diffusion-weighted imaging (DWI) b0/b1000; automatically calculated apparent diffusion coefficient (ADC) maps; Fluid Attenuated Inversion Recovery (FLAIR), flow-sensitive time-of-flight (TOF); T2*-gradient-echo. MR scanner model depends on centre

and year of acquisition: Philips Achieva 1.5T, Philips Achieva 3T, Philips Achieva-dStream 3T, SIEMENS Skyra 3T, SIEMENS – Aera 1.5T. Thus, MRI parameters: Repetition Time (TR), Echo Time (TE), slice thickness, flip angle, Inversion Time or resolution were heterogeneous.

DWI-ASPECTS on each baseline MRI was visually calculated by a neurologist according to the ASPECT study group method (31) (i.e, ASPECTS value was calculated from two standard axial DWI-MR slices, one at the level of the thalamus and basal ganglia, and one just rostral to the ganglionic structures). Radiological outcomes (HI, PH1, PH2) was collected in control imaging's radiologists report or assessed by a neurologist.

Anonymization process and data sharing

The imaging data were extracted from the central server after anonymization. Correspondence between anonymization number and patient identity was held only by the main investigators. The files were extracted into standardized compressed DICOMDIR format and then transferred to the secure and anonymized SHANOIR database (<https://project.inria.fr/shanoir/>). A standardized file containing the above-mentioned clinical data was attached to the imaging data for each patient. These data were then manipulated by the engineers and processed by the algorithm in a totally anonymous and secure way.

Imaging pre-processing

A noise reduction technique was applied. All MRI contrasts have been reframed together (patient by patient) and then reframed on a shared template. Then, brains were extracted, and images resized to get the same geometry (voxel size and section thickness).

2.4. Machine learning algorithm

A ResNet algorithm was used in this work.

The common architecture of ResNet algorithms is freely available online (https://www.tensorflow.org/api_docs/python/tf/keras/applications/resnet). It was composed of convolutional layers preceding “fully connected” linear layers (Figure 3). In order to compensate for the imbalance between the two groups thus formed (n=28 versus n=72), a different weighting has been applied to the two classes within the loss function used into the algorithm. Class weights applied to control group values and sICH group values was respectively 1.39 & 3.57. The binary-cross entropy loss function integrated inside the algorithm

can be expressed as follows:

$$H_p(q) = -\frac{1}{N} \sum_{i=1}^N y_i \cdot \log(p(y_i)) + (1 - y_i) \cdot \log(1 - p(y_i))$$

The cohort was designed into two groups according to prediction attempted: sICH group consisted of symptomatic intracerebral haemorrhage (sICH) subjects while a second was formed by merging both non-symptomatic ICH and No ICH groups to constitute one single control group.

A cross-validation (CV) approach has been used. Briefly, the dataset was randomly partitioned into 10 equal size subsamples. A single subsample is retained as the validation data, 2 subsamples are retained as the testing data and the remaining 7 subsamples are used as training data. The validation process is then repeated 10 times, with each of the 10 subsamples used exactly once as the validation data. The results (n=10) can then be averaged to produce the final estimation. The CV is an efficient way for validating a model and for assessing how it will generalize to an independent data set. These applied techniques allow to avoid bias that would otherwise occur if the performance were assessed on the same data that trained the model. The algorithm, after training, attributed to each slice a value which could be used as a threshold to

classify this slice in either group (sICH or control). In order to obtain a clinically relevant patient-level result, we averaged the probability of each slice for the same patient to obtain a single probability per patient. We then varied the probability threshold to find the one which gave the best ranking performance.

A first analysis model was considering in 2D analysis slices of interest previously selected by a neurologist by analysing DWI sequence. A first CV process was carried out under the following conditions: input data included the following five contrasts DWI b0 and b1000, ADC, FLAIR and T2*. A second CV process integrated in addition clinical data. These clinical data were age, baseline NIHSS, high blood pressure history, diabetes mellitus history, atrial fibrillation history, onset-to-treatment time, anti-thrombotic treatment in progress. As Time-Of-Flight (TOF) could not be integrated to the input data, due to a too large difference in the field-of-view acquired as compare to the other MR sequences, an additional parameter has been included in the clinical data: the presence or absence of LVO.

A second analysis model was considering overall slices using a 2D analysis slice by slice. Two CV process was carried out as described above, with and without clinical data.

2.5. Statistical analysis

Categorical data were analysed in contingency tables using the χ^2 statistic to assess significance. Continuous data were compared using a student t-test when normality was assessed by Shapiro-Wilk test, using a Wilcoxon nonparametric test when it was not. P-value < 0.05 was considered significant. Multivariate analysis was performed considering significant variables with $p < 0.2$ in univariate analysis and relevant variables. Sensitivity and specificity of DWI-ASPECTS and THRIVE score were calculated regarding relevant cut-off value.

Accuracy was calculated for ML algorithm. Receiver Operating Characteristic (ROC) curves were built and their respective Area Under ROC curve (AUC) were calculated to assess DWI-ASPECTS, THRIVE score and ML algorithm performance. SPSS version 26.0 was used for all statistical analysis.

3. RESULTS




Figure 4: Flow Chart

3.1. Baseline characteristics

A total of 106 patients satisfied inclusion criteria. Four subjects were excluded because imaging data were corrupted. Two patients were excluded due to imaging data extraction fail.

Our cohort was then composed of 100 subjects: 28 in sICH group, 27 in non-symptomatic ICH group, 45 in no ICH group (Figure 4).

Distribution of baseline demographic characteristics, comorbidities and radiological findings, acute treatment undertaken and main outcomes between patients along the 3 groups are shown

in Table 1. Regarding acute treatment undertaken, 53 patients underwent only IVT, 9 patients underwent only MT and, 38 patients underwent a bridging therapy.

	sICH n = 28	Non-symptomatic ICH n = 27	No ICH n = 45
Age, mean ± sd	75.2 ± 14.5	72.4 ± 12.1	71.04 ± 14.2
Sex, n (%)			
Female	15 (54%)	10 (37%)	22 (49%)
Male	13 (46%)	17 (63%)	23 (51%)
Right sided stroke	12 (43%)	18 (67%)	28 (62%)
Treatment, n (%)			
IVT	16 (57%)	13 (48%)	24 (53%)
MT	5 (18%)	1 (4%)	3 (7%)
Bridging	7 (25%)	13 (48%)	18 (40%)
Clinical data			
NIHSS, median [IQ]	17 [13-21.3]	16 [14-17.5]	15 [13-17]
OTT, mean ± sd	199.3 ± 46.5	171.8 ± 44.2	172.97 ± 59.2
HBP, n (%)	20 (71.4%)	12 (44.4%)	28 (59.6%)
AF, n (%)	10 (35.7%)	11 (40.7%)	12 (26.7%)
DM, n (%)	9 (32.1%)	2 (7.4%)	10 (21.3%)
Antithrombotic, n (%)	19 (67.9%)	8 (29.6%)	13 (28.9%)
Imaging data			
DWI ASPECTS, median [IQ]	7 [5.75-8]	7 [6-8]	8 [7-9]
LVO, n (%)	22 (78%)	23 (85%)	36 (80%)
Clinical and radiological outcomes			
PH1	6 (21.4%)	0	-
PH2	16 (57.2%)	3 (11.2%)	-
HI	6 (21.4%)	24 (88.8%)	-
Functional independence, n (%)	0	12 (44.4%)	28 (62.2%)
In-hospital mortality, n (%)	15 (53%)	2 (7.4%)	1 (2%)

Table 1: Baseline characteristics according to each sample – IVT: Intravenous thrombolysis ; MT : mechanical thrombectomy; NIHSS : National Institutes of Health Stroke Scale; OTT : Onset-to-Treatment Time; HBP : High Blood Pressure history; AF : Atrial Fibrillation; DM : Diabetes Mellitus; LVO : Large Vessel Occlusion; mRS : modified Rankin Scale

3.2. Radiological and Clinical outcomes

Radiological and clinical outcomes between sICH and non-symptomatic ICH subjects are summarized in Table 2. In-hospital mortality was significantly higher in sICH subjects ($p < 0.001$). On the contrary functional independence at discharge was significantly less frequent in sICH subjects ($p < 0.05$). Rate of PH 1 and 2 were significantly higher in sICH group compared to non-symptomatics ($p < 0.05$). Rate of PH 1 and 2 were significantly higher in sICH group compared to non-symptomatics ($p < 0.001$). Functional outcomes remained significantly poorer in sICH group compared to control group with a lower rate of functional independence (0% vs 44.4%, $p < 0.005$) and a higher in-hospital mortality rate (53% vs 4.1%, $p < 0.005$) as shown in Table 3. Figure 5 illustrates Rankin at discharge's distribution according to group.

Radiological and clinical outcomes	sICH	Non-symptomatic ICH	P-value
Functional independence at discharge, n (%)	0	12 (44.4%)	$P < 0.05$
In-hospital mortality, n (%)	15 (53%)	2 (7.4%)	$P < 0.001$
PH1, n (%)	6 (21.4%)	0	$P < 0.001$
PH2, n (%)	16 (57.2%)	3 (11.2%)	
HI, n (%)	6 (21.4%)	24 (88.8%)	

Table 2: Radiological and clinical outcomes regarding sICH and non-symptomatic ICH subjects.




Figure 5: Modified Rankin scale (mRS) at discharge according to radiological features

3.3. sICH characteristics: univariate and multivariate analyses

Table 3 summarizes univariate analyses between sICH and control groups and main results of the multivariate model built afterwards. Baseline demographic characteristics were similar in age and sex. Type of treatment undertaken did not differ significantly between both groups. A non-significant higher rate of left stroke can be noted in the sICH group. Stroke severity, according initial NIHSS score did not differ significantly (median NIHSS was 17 [13–21.3] in sICH group versus 15.5 [13-17] in control group, $p = 0.12$). Main comorbidities also did not differ between the two groups. There was higher rate of diabetes mellitus in sICH group without reaching significance compared control group (32.1% vs 15.3%, $p = 0.058$).

Onset-to-Treatment Time was significantly higher in sICH group (mean OTT was 199.3 ± 46.5 vs 172.3 ± 53.8 , OR = 1.01 [1.001 – 1.02], $p = 0.01$). Use of prior anti-thrombotic treatment was more frequent in sICH group (67.9% vs 29.2%, OR = 5.1 [2 – 13.2], $p < 0.005$).

Concerning baseline imaging data, DWI-ASPECTS was lower in sICH group (median DWI-ASPECTS was 7 [5.75–8] in sICH group versus 8 [6–9] in control group, OR = 0.78 [0.63 – 0.97], $p = 0.04$). LVO was found for 78% of patients in sICH and 82% in controls which did not differ significantly.

Results of multivariate model main are shown in Table 3. Briefly, only prior use of antithrombotic treatment remained a predictive factor of sICH with OR = 4.4 [1.5 – 13.2] ($p = 0.008$), as well a DWI-ASPECTS with OR = 0.73 [0.56 – 0.97], $p = 0.03$ and THRIVE score (OR = 5.14, CI 95% [1.7 – 15.6], $p = 0.004$). Baseline NIHSS, high blood pressure history, atrial fibrillation, diabetes mellitus history, OTT and large vessel occlusion did not differ significantly between both groups.

	Groups		Univariate analysis		Multivariate model	
	sICH n = 28	Controls n = 72	OR [95%]	p-value	OR [95%]	p-value
Age, mean ± sd	75.2 ± 14.5	71.5 ± 13.4	1.02 [0.98 – 1.06]	0.25	1.007 [0.95 – 1.05]	0.8
Sex, n (%)			0.69 [0.29 – 1.67]	0.41	-	-
Female	15 (54%)	32 (45%)				
Male	13 (46%)	40 (55%)				
Right sided stroke	12 (43%)	46 (64%)	2.36 [0.96 – 5.74]	0.056	-	-
Treatment, n (%)			-	0.073	-	-
IVT	16 (57%)	37 (51%)				
MT	5 (18%)	4 (6%)				
Bridging	7 (25%)	31 (43%)				
Clinical data						
NIHSS, median [IQ]	17 [13–21.3]	15.5 [13-17]	1.09 [0.98 – 1.2]	0.12	1.07 [0.94 – 1.21]	0.27
OTT, mean ± sd	199.3 ± 46.5	172.3 ± 53.8	1.01 [1.001 – 1.02]	0.01	1.009 [0.99 – 1.02]	0.1
HBP, n (%)	20 (71.4%)	38 (52.8%)	2.2 [0.9 – 5.7]	0.09	1.04 [0.26 – 4.09]	0.95
AF, n (%)	10 (35.7%)	23 (31.9%)	1.18 [0.47 – 2.96]	0.7	0.98 [0.29 – 3.21]	0.97
DM, n (%)	9 (32.1%)	11 (15.3%)	2.63 [0.9 – 7.3]	0.058	1.82 [0.52 – 6.35]	0.34
Antithrombotic, n (%)	19 (67.9%)	21 (29.2%)	5.1 [2 – 13.2]	< 0.005	4.4 [1.5 – 13.2]	0.008
THRIVE score, median [IQ]	5 [4-6]	4 [3-5]	1.54 [1.15 – 2.07]	0.004	5.14 [1.7 – 15.6]	0.004
Imaging data						
DWI ASPECTS, median [IQ]	7 [5.75–8]	8 [6–9]	0.78 [0.63 – 0.97]	0.04	0.73 [0.56 – 0.97]	0.03
LVO, n (%)	22 (78%)	59 (82%)	0.8 [0.27 – 2.39]	0.69	0.33 [0.08 – 1.36]	0.12
Clinical outcomes						
Functional independence, n (%)	0	40 (56%)	-	< 0.005	-	-
In-hospital mortality, n (%)	15 (53%)	3 (4.1%)	-	< 0.005	-	-

Table 3: Baseline characteristics and outcomes between sICH subjects and control group, uni and multivariate analyses. Abbreviations are detailed in Table 1

3.4. Symptomatic intracranial haemorrhage prediction

3.4.1. THRIVE score performance

THRIVE score was significantly higher in the sICH group (OR = 1.54, CI 95% [1.15 – 2.07], $p = 0.004$) and remained an independent predictor of sICH in multivariate analysis (OR = 5.14, CI 95% [1.7 – 15.6], $p = 0.004$).

THRIVE score performance is represented by the ROC curve in Figure 6, with AUC of 0.664 (CI 95% [0.548 – 0.781]). Table 4 indicates Sensitivity/Specificity values, Positive Predictive and Negative Predictive Values (PPV/NPV) for each cutpoint. We can notice that a threshold of 4 gave an excellent sensitivity of 92.8% but a poor specificity of 40.2% for an overall 55% accuracy. Raising the threshold to 5 improved specificity to 55.5% but at the expense of a poor sensitivity of 60.7% for an overall 57% accuracy.

3.4.2. DWI-ASPECTS performance

DWI-ASPECT score was an independent factor of sICH in multivariate analysis (OR = 0.73 CI 95% [0.56 – 0.97], $p = 0.03$). DWI-ASPECT score predicted sICH with an AUC of 0.634 (CI 95% [0.508 – 0.761]). Table 4 indicates Sensitivity/Specificity values, Positive Predictive and Negative Predictive Values (PPV/NPV) for interest cutpoints. We can notice that setting a threshold of 7 - which is the most commonly used threshold in the literature- resulted in poor sensitivity and specificity calculated at 60.7% and 61.1% respectively which corresponds to 61% accuracy. ROC curve of DWI-ASPECTS is presented in Figure 6.

	Cutpoint	Se (%)	Sp (%)	PPV (%)	NPV (%)	Accuracy (%)	AUC
THRIVE SCORE	4	92.86%	40.28%	37.68%	93.55%	55%	0.664
	5	60.71%	55.56%	34.69%	78.43%	57%	
DWI-ASPECTS	7	60.71%	61.11%	37.78%	80%	61%	0.634
	8	82.14%	36.11%	33.33%	83.87%	49%	

Table 4: THRIVE score and DWI-ASPECTS sICH prediction: Sensitivity (Se), Specificity (Sp), Positive (PPV) and Negative (NPV) Predictive values according to relevant cutpoints.

3.4.3. Machine Learning (ML) algorithm

“Resnet” neural network performances were assessed in various configurations. Enlarging number of layers which, for reminder, increases the number of parameters taken into account and therefore the complexity of the network, did not improve performances. The network was finally made up of 15 convolutional layers preceding a single fully connected layer. Various data augmentation methods were used such as Gaussian blur effect, x-axis filing flip, contrast variation via the gamma function. None of these methods resulted in a significant improvement in performance.

According to each imaging dataset model, including either overall slices (all brain model) or pre-selected slices of interest (masked brain model), best results after learning phase and on validation dataset are listed below.

The best performance was achieved using the masked brain model computing the 5 contrasts and clinical data inputs (accuracy 77%). Without clinical data accuracy was 74%.

“All brain” model with clinical data achieved similar accuracy performances with best accuracy calculated at 74% compared 72% without clinical data inputs.

ROC curve and AUC for all brain and masked brain models including clinical data are illustrated in Figure 6. AUC were estimated at 0.624 (CI 95% [0.497 - 0.752]) and 0.658 (CI 95% [0.534 - 0.783]) respectively.




Figure 6: Receiver operating curves (ROC) and Area Under ROC Curves (AUC) of DWI-ASPECT SCORE (purple), THRIVE score (beige), artificial neural network (ANN) masked brain (blue) and ANN all brain (green) models.

4. Discussion

4.1. sICH outcomes

First of all, this study highlighted, in a population of anterior circulation AIS subjects matched on clinical severity and age, a number of clinical and radiological differences in patients with sICH.

Demographic data or reperfusion strategies did not differ significantly between sICH subjects and controls. More surprisingly, cardiovascular co-morbidities and risk factors were not significantly more present in the sICH group.

Only prior use of anti-thrombotic agent appeared as a predictive factor of sICH in multivariate analysis, which was consistent with literature (18). OTT, known as risk factor of sICH (17), was significantly longer in sICH group in univariate analysis but not in multivariate model.

Regarding imaging baseline parameters, DWI-ASPECTS was significantly lower in the sICH group. This argues in favour of a higher lesions volume in the sICH group and is also consistent with data available in the literature (32). Although some authors have linked signs of proximal occlusion and especially spontaneous hyperdensity of the middle cerebral artery on CT scan (20), LVO rate was not significantly higher in sICH subjects inside our cohort. Thus, these results raises the hypothesis that infarcted tissue volume may be more relevant than hypoperfusion volume to predict sICH, although these two parameters have been related to poor outcome (45).

Functional outcomes and survival rate were poorer in sICH group compared to the reference group but also, more specifically, compared to non-symptomatic ICH patients, strengthens the importance of identifying patients at risk of sICH. From a radiological point of view, we noticed a significantly higher rate of PH within sICH subjects compared to the reference group but also compared to ICH patients. This underlines the probable link between the type of haemorrhagic

transformation and its severity, as suggested by other authors : post hoc analysis of ECASS I and II data demonstrated an increased risk of early neurological deterioration and 3-month mortality after PH2-related ICH (13,17), describing close to a 50% mortality with PH2. Future work could therefore specifically focus on the identification of PH2 predictive factors rather than sICH. The analysis of the initial imaging, including using ML methods as presented here, could be particularly relevant in this case.

4.2. Predictive performances

Secondly, this work highlighted difficulties for current tools to predict sICH.

The THRIVE score showed in our cohort moderate performance in predicting sICH. It was significantly higher in the sICH group and remained an independent predictor of symptomatic haemorrhagic transformation in multivariate analysis. Nevertheless, in detail, its performance on a ROC curve remained modest with AUC estimated 0.664. Applying a decision-making threshold of 4 allowed to obtain an excellent sensitivity of 92.8% but a poor specificity of 40.2%. These results were similar to other authors (30) and probably explain why this score is not used in routine practice.

DWI-ASPECT score also showed poor performance to predict sICH in our cohort. It was significantly higher in the sICH group and remained an independent predictor of sICH in multivariate analysis. Nevertheless, performances remained modest with AUC estimated 0.634. Sensitivity and specificity were 61% for both with a 7 points cut-off which is current threshold used in current practice. Setting a 8 points cut-off allows to get close to other authors results (32) and resulted in an interesting 82% Sensitivity but once again at the expense of a poor 33% specificity. To discuss these poor DWI-ASPECTS results, we should remember that ASPECTS was originally assessed on CT scan and so based on hypoattenuation signs (19). This hypodensity in CT scan appears more latter than DWI hyperintensity, and therefore reflects

greater substance changes in infarcted tissue and a greater progression towards necrosis. Thus, the ASPECTS on CT scan was probably more specific of a tissue at risk of haemorrhage. Moreover, DWI-ASPECT is a good estimation of lesion volume, but some limits are recurrently noted: lesion location is not considered and lesion volume could be more reliably quantified on DWI maps nowadays (46). In addition, this global estimate of lesion volume, unlike an automated analysis, does not take into account differences in the intensity of the DWI signal and ADC, even though this parameter could be an indicator of the degree of ischaemia and has been linked to the risk of sICH (21,47).

Concerning ML algorithm performances, this study showed that in a cohort of patients matched on clinical severity and age, an MRI based ML algorithm was able to discriminate subjects at risk of sICH. Best accuracy of 77% was reached with a model focused on slices of interest only including ischemic parenchyma and including clinical data. Interestingly, removing clinical data from the inputs did not significantly affect the algorithm performance (accuracy 74%). This finding seems consistent with the absence of significant differences between groups for these parameters and points to other possibilities that we will discuss below. Surprisingly, performances of “all brain” model were similar to the “masked brain” model. From a conceptual point of view, we expected that this model would not work properly in a 2D analysis, as independent slices of unaffected parenchyma to be labelled sICH or control group. This could be explained by the presence of predictive elements in unaffected parenchyma (i.e., FLAIR intensities or microbleeds burden as described above). Nevertheless, in detail, building ROC curves revealed AUC estimated at 0.624 and 0.658 which do not outperform DWI-ASPECT and THRIVE scores.

To sum up, we have attempted a new approach using a machine learning algorithm capable of synthesizing the diversity of data that seem necessary for sICH prediction. This study brings novel contributions to sICH prediction in that it was designed to develop a ML method usable in current practice by clinicians to predict sICH. As a result, we chose to use standard practice MRI data computed using different MRI equipment and we decided to implement data available before introduction of reperfusion therapies. We proved that ML algorithm was able to process the diversity of information provided to achieve sICH prediction. Nevertheless, ML algorithm's predictive performances did not outperform current tools. These poor performances and, moreover, the importance of pre-processing, which requires the selection of interesting slices by the investigator, do not make it a tool that can be routinely used to date. These modest but encouraging results open up a number of avenues for development.

4.3. Study limitations and future directions

The above-mentioned choices led to certain limitations.

First, to be as close as possible to current practice we included all treated patients whatever the treatment introduced. But this has consequently increased the heterogeneity of a cohort which is also small in size, in particular because some patients did not receive IVT.

Then, clinical data used here may not be the most relevant ones, as highlighted by the fact that they little differed in sICH subjects. This leads us to look for more relevant input clinical data. Baseline mean blood pressure and glycaemia are missing, while those factors are more strongly correlated to sICH risk than HBP and diabetes mellitus history (48,49). In order to be useful in current practice, it would also seem more relevant to consider the Onset-to-Imaging Time rather than OTT which cannot be estimated *a priori*, especially in case of MT without prior IVT.

MRI data seemed to be particularly relevant since we can already see a discriminating power of the algorithm mainly based on these imaging data. We can discuss the choice of MR

sequences as input data. In this work, the absence of perfusion sequence or at least a TOF sequence that might have provided information on collaterality can be highlighted. Perfusion sequences were not usually performed in most of stroke centres. However, permeability imaging, which consists of delayed acquisition perfusion imaging assessing late extravasation into the brain parenchyma, has been identified as predictor of sICH in that it requires the blood-brain barrier to be sufficiently disrupted to allow entry of gadolinium (42). Blood brain barrier disruption is a critical element involved in the development of haemorrhagic transformation (50) and could be identified by MRI on gadolinium enhanced-T1-weighted sequences (51). Perfusion imaging currently has no place in stroke emergency imaging. However, in this field too, certain ML techniques could bring new contributions, by reconstructing fully injected sequences after a low-dose injection or recovering permeability parameters from a single brief acquisition using fingerprinting methods (52,53). TOF sequence failed to be integrated for technical reasons because of differences in acquisition volume. However it has been suggested that the presence of a robust collateral circulation on vessel imaging is associated with a lower risk of ICH, which may be partly due not only to the smaller volume of the infarction but also to the collateral circulation, ensuring sufficient circulation for the cerebral endothelial cells so that they do not rupture as a result of prolonged ischemia (54).

Then, the small size of the cohort may be the most important limitation. Indeed, ML methods require a large amount of data to improve learning phase. Due to the small workforce, particular attention had to be paid to the risk of overfitting. Overfitting occurs when the algorithm, through repeated training, finally recognises the training dataset. We then observe excellent performance on the learning data but paradoxically poor performance on the validation set. This risk of "overlearning" increases in inverse proportion to the cohort size. This led us to limit the learning phase.

As a result, this lack of data led us to focus on a 2D analysis which had the advantage of artificially increasing cohort size. Each patient gave rise to a certain number of cuts which could be analysed independently during the training phase. However, this strategy most certainly led to the loss of a lot of relevant information to prediction. The advantage of “all brain” model was to benefit from all the information provided by the parenchyma both inside and outside the lesion, it may consider visible leukopathy in FLAIR (23,55) and microbleeds burden visible on T2* sequence which has also been correlated with sICH risk by some authors (24). But the interpretation of non-affected and non-informative parenchyma slices may have reduced its performance. On the contrary, “slices of interest” model was supposed to get around this difficulty of 2D analysis but at the expense of a loss of information. The 2D analysis also meant that neither of the two models could accurately calculate the lesion volume, although this data seems to be a key factor (25). It has been reported that patients with large lesions on DWI-MRI and Perfusion Weighted MRI have a high risk of developing sICH and a worse clinical outcome after intravenous thrombolysis (47). The Diffusion and Perfusion Imaging Evaluation for Understanding Stroke Evolution (DEFUSE) study identified a threshold of ≥ 100 -mL lesion volume in DWI sequence (45), characterizing a malignant course. Moreover, little access to the location of the lesion within the brain was allowed using this 2D analysis, which could be essential to assess the symptomatic nature of the haemorrhage.

Finally, increasing dataset seems therefore to be a first necessary condition to further works. A larger dataset would allow 3D analysis, avoiding a selection of interest slices that results in the loss of valuable information such as lesion volume and unaffected parenchyma characteristics. Enlarging dataset through a prospective work would also be an opportunity to collect more relevant clinical and radiological inputs. Moreover, validation of these results on a more heterogeneous external population is also required.

5. Conclusion

THÈSE SOUTENUE PAR : Florent LEPILLIET

TITRE:

PREDICTING SYMPTOMATIC INTRACRANIAL HAEMORRHAGE IN ACUTE ISCHEMIC STROKE: INTRODUCTION TO A CLINICAL-RADIOLOGICAL AND MRI-BASED MACHINE LEARNING APPROACH

Ischemic stroke is a leading cause of death and disability in adult. Reperfusion therapies using intravenous thrombolysis (IVT) and endovascular interventions such as mechanical thrombectomy (MT) improve functional outcomes in patients with acute ischemic stroke (AIS) but unfortunately, these therapeutics increase the risk of intracranial haemorrhage (ICH) of ischemic brain tissue. Several definitions and classifications exist to define this complication, on a clinical or radiological level. However, high evidence indicates that symptomatic ICH (sICH) is the most relevant definition as it is most correlated with poor outcome. Several methods, based on clinical and/or radiological data, failed to predict the risk of haemorrhagic transformation in clinical practice.

We attempted a new approach by training a supervised machine learning (ML) algorithm on clinical and MRI data within a 100 subjects cohort of patients with anterior circulation AIS treated by IVT and/or MT who underwent sICH (n=28), non-symptomatic ICH (n=27) and 45 controls with no bleeding, matched on clinical severity and age. We compared ML algorithm accuracy to the performance of the clinical Total Health Risks in Vascular Events (THRIVE) score and to the radiological Alberta Stroke Program Early CT Score applied to MR imaging (DWI-ASPECTS).

ML algorithm predicted sICH with an Area Under receiver operating characteristic Curve (AUC) of 0.658 (CI 95% [0.534 – 0.783]). Applied in the cohort, estimated AUC of THRIVE

score and DWI-ASPECTS were 0.664 (CI 95% [0.548 – 0.781]) and 0.634 (IC 95% [0.508 – 0.761]), respectively. Nevertheless, this work showed that this algorithm was able to synthesize all clinical and radiological data provided and integrating the variety of information provided by MR imaging to provide a probability of sICH.

Further studies are needed to improve these performances. Enlarging dataset is needed to improve learning phase, reduce overfitting risk and allow a 3D analysis to avoid data loss. More relevant clinical and radiological data could also be integrated to improve performances. Validation of these results on a more heterogeneous external population is also required.

VU ET PERMIS D'IMPRIMER
Grenoble, le : 8/02/21

LE DOYEN


Pr. Patrice MORAND
pour le conseil
et par délégation
—
Le Doyen de Médecine
Pr. Patrice MORAND

LE PRÉSIDENT DE LA THÈSE

Pr. Olivier DETANTE



References

1. Johnson CO, Nguyen M, Roth GA, Nichols E, Alam T, Abate D, et al. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology*. 1 mai 2019;18(5):439-58.
2. Hacke W, Kaste M, Bluhmki E, Brozman M, Dávalos A, Guidetti D, et al. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med*. 25 sept 2008;359(13):1317-29.
3. Wahlgren N, Ahmed N, Dávalos A, Hacke W, Millán M, Muir K, et al. Thrombolysis with alteplase 3–4.5 h after acute ischaemic stroke (SITS-ISTR): an observational study. *The Lancet*. oct 2008;372(9646):1303-9.
4. Goyal M, Menon BK, van Zwam WH, Dippel DWJ, Mitchell PJ, Demchuk AM, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *The Lancet*. avr 2016;387(10029):1723-31.
5. Emberson J, Lees KR, Lyden P, Blackwell L, Albers G, Bluhmki E, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *Lancet*. 29 nov 2014;384(9958):1929-35.
6. Larrue V, von Kummer R R, Müller A, Bluhmki E. Risk factors for severe hemorrhagic transformation in ischemic stroke patients treated with recombinant tissue plasminogen activator: a secondary analysis of the European-Australasian Acute Stroke Study (ECASS II). *Stroke*. févr 2001;32(2):438-41.
7. Hacke W, Kaste M, Fieschi C, von Kummer R, Davalos A, Meier D, et al. Randomised double-blind placebo-controlled trial of thrombolytic therapy with intravenous alteplase in acute ischaemic stroke (ECASS II). *The Lancet*. 17 oct 1998;352(9136):1245-51.
8. Seet RCS, Rabinstein AA. Symptomatic intracranial hemorrhage following intravenous thrombolysis for acute ischemic stroke: a critical review of case definitions. *Cerebrovasc Dis*. 2012;34(2):106-14.
9. Neuberger U, Kickingereeder P, Schönenberger S, Schieber S, Ringleb PA, Bendszus M, et al. Risk factors of intracranial hemorrhage after mechanical thrombectomy of anterior circulation ischemic stroke. *Neuroradiology*. avr 2019;61(4):461-9.
10. Maier B, Desilles JP, Mazighi M. Intracranial Hemorrhage After Reperfusion Therapies in Acute Ischemic Stroke Patients. *Front Neurol*. 2020;11:599908.
11. Thomalla G, Schwark C, Sobesky J, Bluhmki E, Fiebich JB, Fiehler J, et al. Outcome and Symptomatic Bleeding Complications of Intravenous Thrombolysis Within 6 Hours in MRI-Selected Stroke Patients: Comparison of a German Multicenter Study With the Pooled Data of ATLANTIS, ECASS, and NINDS tPA Trials. *Stroke*. mars 2006;37(3):852-8.

12. Thomalla G, Simonsen CZ, Boutitie F, Andersen G, Berthezene Y, Cheng B, et al. MRI-Guided Thrombolysis for Stroke with Unknown Time of Onset. *New England Journal of Medicine* [Internet]. 16 mai 2018 [cité 8 févr 2021]; Disponible sur: <https://www-nejm-org.proxy.insermbiblio.inist.fr/doi/10.1056/NEJMoa1804355>
13. Fiorelli M, Bastianello S, von Kummer R, del Zoppo GJ, Larrue V, Lesaffre E, et al. Hemorrhagic transformation within 36 hours of a cerebral infarct: relationships with early clinical deterioration and 3-month outcome in the European Cooperative Acute Stroke Study I (ECASS I) cohort. *Stroke*. nov 1999;30(11):2280-4.
14. Tanne D, Kasner SE, Demchuk AM, Koren-Morag N, Hanson S, Grond M, et al. Markers of increased risk of intracerebral hemorrhage after intravenous recombinant tissue plasminogen activator therapy for acute ischemic stroke in clinical practice: the Multicenter rt-PA Stroke Survey. *Circulation*. 9 avr 2002;105(14):1679-85.
15. Whiteley William N., Slot Karsten Bruins, Fernandes Peter, Sandercock Peter, Wardlaw Joanna. Risk Factors for Intracranial Hemorrhage in Acute Ischemic Stroke Patients Treated With Recombinant Tissue Plasminogen Activator. *Stroke*. 1 nov 2012;43(11):2904-9.
16. Kaesmacher J, Kaesmacher M, Maegerlein C, Zimmer C, Gersing AS, Wunderlich S, et al. Hemorrhagic Transformations after Thrombectomy: Risk Factors and Clinical Relevance. *Cerebrovasc Dis*. 2017;43(5-6):294-304.
17. Lees KR, Bluhmki E, von Kummer R, Brott TG, Toni D, Grotta JC, et al. Time to treatment with intravenous alteplase and outcome in stroke: an updated pooled analysis of ECASS, ATLANTIS, NINDS, and EPITHET trials. *Lancet*. 15 mai 2010;375(9727):1695-703.
18. Prabhakaran S, Rivolta J, Vieira JR, Rincon F, Stillman J, Marshall RS, et al. Symptomatic intracerebral hemorrhage among eligible warfarin-treated patients receiving intravenous tissue plasminogen activator for acute ischemic stroke. *Arch Neurol*. mai 2010;67(5):559-63.
19. Schwamm LH, Rosenthal ES, Swap CJ, Rosand J, Rordorf G, Buonanno FS, et al. Hypoattenuation on CT Angiographic Source Images Predicts Risk of Intracerebral Hemorrhage and Outcome after Intra-Arterial Reperfusion Therapy. *American Journal of Neuroradiology*. 1 août 2005;26(7):1798-803.
20. Zou M, Churilov L, He A, Campbell B, Davis SM, Yan B. Hyperdense middle cerebral artery sign is associated with increased risk of hemorrhagic transformation after intravenous thrombolysis for patients with acute ischaemic stroke. *J Clin Neurosci*. juill 2013;20(7):984-7.
21. Selim Magdy, Fink John N., Kumar Sandeep, Caplan Louis R., Horkan Clare, Chen Yi, et al. Predictors of Hemorrhagic Transformation After Intravenous Recombinant Tissue Plasminogen Activator. *Stroke*. 1 août 2002;33(8):2047-52.
22. Suh CH, Jung SC, Cho SJ, Woo D-C, Oh WY, Lee JG, et al. MRI for prediction of hemorrhagic transformation in acute ischemic stroke: a systematic review and meta-analysis. *Acta Radiol*. 18 nov 2019;0284185119887593.

23. Kongbunkiat K, Wilson D, Kasemsap N, Tiamkao S, Jichi F, Palumbo V, et al. Leukoaraiosis, intracerebral hemorrhage, and functional outcome after acute stroke thrombolysis. *Neurology*. 14 févr 2017;88(7):638-45.
24. Tsiygoulis G, Zand R, Katsanos AH, Turc G, Nolte CH, Jung S, et al. Risk of Symptomatic Intracerebral Hemorrhage After Intravenous Thrombolysis in Patients With Acute Ischemic Stroke and High Cerebral Microbleed Burden: A Meta-analysis. *JAMA Neurol*. 1 juin 2016;73(6):675-83.
25. El Nawar R, Yeung J, Labreuche J, Chadenat M-L, Duong DL, De Malherbe M, et al. MRI-Based Predictors of Hemorrhagic Transformation in Patients With Stroke Treated by Intravenous Thrombolysis. *Front Neurol* [Internet]. 27 août 2019 [cité 19 janv 2021];10. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6719609/>
26. Sung S-F, Chen SC-C, Lin H-J, Chen Y-W, Tseng M-C, Chen C-H. Comparison of Risk-scoring Systems in Predicting Symptomatic Intracerebral Hemorrhage After Intravenous Thrombolysis. *Stroke*. juin 2013;44(6):1561-6.
27. Strbian D, Michel P, Seiffge DJ, Saver JL, Numminen H, Meretoja A, et al. Symptomatic Intracranial Hemorrhage After Stroke Thrombolysis: Comparison of Prediction Scores. *Stroke*. mars 2014;45(3):752-8.
28. Yaghi S, Willey JZ, Cucchiara B, Goldstein JN, Gonzales NR, Khatri P, et al. Treatment and Outcome of Hemorrhagic Transformation After Intravenous Alteplase in Acute Ischemic Stroke: A Scientific Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke* [Internet]. déc 2017 [cité 13 févr 2020];48(12). Disponible sur: <https://www.ahajournals.org/doi/10.1161/STR.0000000000000152>
29. Whiteley William N., Thompson Douglas, Murray Gordon, Cohen Geoff, Lindley Richard I., Wardlaw Joanna, et al. Targeting Recombinant Tissue-Type Plasminogen Activator in Acute Ischemic Stroke Based on Risk of Intracranial Hemorrhage or Poor Functional Outcome. *Stroke*. 1 avr 2014;45(4):1000-6.
30. Flint AC, Faigeles BS, Cullen SP, Kamel H, Rao VA, Gupta R, et al. THRIVE Score Predicts Ischemic Stroke Outcomes and Thrombolytic Hemorrhage Risk in VISTA. *Stroke*. déc 2013;44(12):3365-9.
31. Barber PA, Demchuk AM, Zhang J, Buchan AM. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. *The Lancet*. mai 2000;355(9216):1670-4.
32. Singer Oliver C., Kurre Wiebke, Humpich Marek C., Lorenz Matthias W., Kastrup Andreas, Liebeskind David S., et al. Risk Assessment of Symptomatic Intracerebral Hemorrhage After Thrombolysis Using DWI-ASPECTS. *Stroke*. 1 août 2009;40(8):2743-8.
33. Andrade de Oliveira A, Carthery-Goulart MT, Oliveira Júnior PP de M, Carrettiero DC, Sato JR. Defining multivariate normative rules for healthy aging using neuroimaging and machine learning: an application to Alzheimer's disease. *J Alzheimers Dis*. 2015;43(1):201-12.

34. Greenstein D, Malley JD, Weisinger B, Clasen L, Gogtay N. Using multivariate machine learning methods and structural MRI to classify childhood onset schizophrenia and healthy controls. *Front Psychiatry*. 2012;3:53.
35. An J, Fang P, Wang W, Liu Z, Hu D, Qiu S. Decreased white matter integrity in mesial temporal lobe epilepsy: a machine learning approach. *Neuroreport*. 9 juill 2014;25(10):788-94.
36. Peter R, Korfiatis P, Blezek D, Oscar Beitia A, Stepan-Buksakowska I, Horinek D, et al. A quantitative symmetry-based analysis of hyperacute ischemic stroke lesions in noncontrast computed tomography. *Med Phys*. janv 2017;44(1):192-9.
37. Boldsen JK, Engedal TS, Pedraza S, Cho T-H, Thomalla G, Nighoghossian N, et al. Better Diffusion Segmentation in Acute Ischemic Stroke Through Automatic Tree Learning Anomaly Segmentation. *Front Neuroinform*. 2018;12:21.
38. McKinley R, Häni L, Gralla J, El-Koussy M, Bauer S, Arnold M, et al. Fully automated stroke tissue estimation using random forest classifiers (FASTER). *J Cereb Blood Flow Metab*. août 2017;37(8):2728-41.
39. Alawieh A, Zaraket F, Alawieh MB, Chatterjee AR, Spiotta A. Using machine learning to optimize selection of elderly patients for endovascular thrombectomy. *J Neurointerv Surg*. 2 févr 2019;
40. Asadi H, Dowling R, Yan B, Mitchell P. Machine learning for outcome prediction of acute ischemic stroke post intra-arterial therapy. *PLoS ONE*. 2014;9(2):e88225.
41. Bentley P, Ganesalingam J, Carlton Jones AL, Mahady K, Epton S, Rinne P, et al. Prediction of stroke thrombolysis outcome using CT brain machine learning. *Neuroimage Clin*. 30 mars 2014;4:635-40.
42. Scalzo F, Alger JR, Hu X, Saver JL, Dani KA, Muir KW, et al. Multi-center prediction of hemorrhagic transformation in acute ischemic stroke using permeability imaging features. *Magn Reson Imaging*. juill 2013;31(6):961-9.
43. Yu Y, Guo D, Lou M, Liebeskind D, Scalzo F. Prediction of Hemorrhagic Transformation Severity in Acute Stroke From Source Perfusion MRI. *IEEE Trans Biomed Eng*. 2018;65(9):2058-65.
44. Chartrand G, Cheng PM, Vorontsov E, Drozdal M, Turcotte S, Pal CJ, et al. Deep Learning: A Primer for Radiologists. *Radiographics*. déc 2017;37(7):2113-31.
45. Albers GW, Thijs VN, Wechsler L, Kemp S, Schlaug G, Skalabrin E, et al. Magnetic resonance imaging profiles predict clinical response to early reperfusion: The diffusion and perfusion imaging evaluation for understanding stroke evolution (DEFUSE) study. *Annals of Neurology*. 2006;60(5):508-17.
46. Schröder Julian, Cheng Bastian, Ebinger Martin, Köhrmann Martin, Wu Ona, Kang Dong-Wha, et al. Validity of Acute Stroke Lesion Volume Estimation by Diffusion-Weighted Imaging—Alberta Stroke Program Early Computed Tomographic Score Depends on Lesion Location in 496 Patients With Middle Cerebral Artery Stroke. *Stroke*. 1 déc 2014;45(12):3583-8.

47. Tong DC, Adami A, Moseley ME, Marks MP. Prediction of Hemorrhagic Transformation Following Acute Stroke: Role of Diffusion- and Perfusion-Weighted Magnetic Resonance Imaging. *Arch Neurol.* 1 avr 2001;58(4):587-93.
48. Huang J, Liu B, Yang C, Chen H, Eunice D, Yuan Z. Acute hyperglycemia worsens ischemic stroke-induced brain damage via high mobility group box-1 in rats. *Brain Res.* 16 oct 2013;1535:148-55.
49. Tsvigoulis G, Frey JL, Flaster M, Sharma VK, Lao AY, Hoover SL, et al. Pre-tissue plasminogen activator blood pressure levels and risk of symptomatic intracerebral hemorrhage. *Stroke.* nov 2009;40(11):3631-4.
50. del Zoppo GJ, von Kummer R, Hamann GF. Ischaemic damage of brain microvessels: inherent risks for thrombolytic treatment in stroke. *J Neurol Neurosurg Psychiatry.* juill 1998;65(1):1-9.
51. Knight RA, Barker PB, Fagan SC, Li Y, Jacobs MA, Welch KM. Prediction of impending hemorrhagic transformation in ischemic stroke using magnetic resonance imaging in rats. *Stroke.* janv 1998;29(1):144-51.
52. Lemasson B, Pannetier N, Coquery N, Boisserand LSB, Collomb N, Schuff N, et al. MR Vascular Fingerprinting in Stroke and Brain Tumors Models. *Sci Rep.* 24 nov 2016;6:37071.
53. Gong E, Pauly JM, Wintermark M, Zaharchuk G. Deep learning enables reduced gadolinium dose for contrast-enhanced brain MRI. *J Magn Reson Imaging.* août 2018;48(2):330-40.
54. Bang OY, Saver JL, Kim SJ, Kim G-M, Chung C-S, Ovbiagele B, et al. Collateral Flow Averts Hemorrhagic Transformation After Endovascular Therapy for Acute Ischemic Stroke. *Stroke.* août 2011;42(8):2235-9.
55. Shi Zhong-Song, Loh Yince, Liebeskind David S., Saver Jeffrey L., Gonzalez Nestor R., Tateshima Satoshi, et al. Leukoaraiosis Predicts Parenchymal Hematoma After Mechanical Thrombectomy in Acute Ischemic Stroke. *Stroke.* 1 juill 2012;43(7):1806-11.



Le serment d'Hippocrate

Texte revu par l'Ordre des médecins en 2012

“

Au moment d'être admis(e) à exercer la médecine, je promets et je jure d'être fidèle aux lois de l'honneur et de la probité.

Mon premier souci sera de rétablir, de préserver ou de promouvoir la santé dans tous ses éléments, physiques et mentaux, individuels et sociaux.

Je respecterai toutes les personnes, leur autonomie et leur volonté, sans aucune discrimination selon leur état ou leurs convictions. J'interviendrai pour les protéger si elles sont affaiblies, vulnérables ou menacées dans leur intégrité ou leur dignité. Même sous la contrainte, je ne ferai pas usage de mes connaissances contre les lois de l'humanité.

J'informerai les patients des décisions envisagées, de leurs raisons et de leurs conséquences. Je ne tromperai jamais leur confiance et n'exploiterai pas le pouvoir hérité des circonstances pour forcer les consciences.

Je donnerai mes soins à l'indigent et à quiconque me les demandera. Je ne me laisserai pas influencer par la soif du gain ou la recherche de la gloire.

Admis(e) dans l'intimité des personnes, je tairai les secrets qui me seront confiés. Reçu(e) à l'intérieur des maisons, je respecterai les secrets des foyers et ma conduite ne servira pas à corrompre les mœurs.

Je ferai tout pour soulager les souffrances. Je ne prolongerai pas abusivement les agonies. Je ne provoquerai jamais la mort délibérément.

Je préserverai l'indépendance nécessaire à l'accomplissement de ma mission. Je n'entreprendrai rien qui dépasse mes compétences. Je les entretiendrai et les perfectionnerai pour assurer au mieux les services qui me seront demandés.

J'apporterai mon aide à mes confrères ainsi qu'à leurs familles dans l'adversité. Que les hommes et mes confrères m'accordent leur estime si je suis fidèle à mes promesses ; que je

sois déshonoré(e) et méprisé(e) si j'y manque. ”