



HAL
open science

A comprehensive study of medically serious suicide attempts hospitalized in France: incidence and associated factors

Julien Corbé

► **To cite this version:**

Julien Corbé. A comprehensive study of medically serious suicide attempts hospitalized in France: incidence and associated factors. Human health and pathology. 2021. dumas-03563477

HAL Id: dumas-03563477

<https://dumas.ccsd.cnrs.fr/dumas-03563477v1>

Submitted on 9 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

UNIVERSITE DE MONTPELLIER
FACULTE DE MEDECINE MONTPELLIER-NIMES

THESE

Pour obtenir le titre de

DOCTEUR EN MEDECINE

Présentée et soutenue publiquement
Par

Julien CORBÉ

Le 29 septembre 2021

TITRE

**A Comprehensive Study Of Medically Serious Suicide Attempts
Hospitalized In France: Incidence And Associated Factors**

Directeur de thèse : Monsieur le Professeur Jorge LOPEZ CASTROMAN

JURY

Président : Monsieur le Professeur Jorge LOPEZ CASTROMAN

Assesseurs :

Madame le Professeur Emilie OLIE

Monsieur le Docteur Mathieu LACAMBRE

UNIVERSITE DE MONTPELLIER
FACULTE DE MEDECINE MONTPELLIER-NIMES

THESE

Pour obtenir le titre de

DOCTEUR EN MEDECINE

Présentée et soutenue publiquement
Par

Julien CORBÉ

Le 29 septembre 2021

TITRE

**A Comprehensive Study Of Medically Serious Suicide Attempts
Hospitalized In France: Incidence And Associated Factors**

Directeur de thèse : Monsieur le Professeur Jorge LOPEZ CASTROMAN

JURY

Président : Monsieur le Professeur Jorge LOPEZ CASTROMAN

Asseseurs :

Madame le Professeur Emilie OLIE

Monsieur le Docteur Mathieu LACAMBRE

Professeurs Honoraires

ALBAT Bernard	BRUNEL Michel	HUMEAU Claude	MIRO Luis
ALLIEU Yves	CANAUD Bernard	JAFFIOL Claude	NAVARRO Maurice
ALRIC Robert	CHAPTAL Paul-André	JANBON Charles	NAVRATIL Henri
ARNAUD Bernard	CIURANA Albert-Jean	JANBON François	OTHONIEL Jacques
ASENCIO Gérard	CLOT Jacques	JARRY Daniel	PAGES Michel
ASTRUC Jacques	COSTA Pierre	JOURDAN Jacques	PEGURET Claude
AUSSILLOUX Charles	D'ATHIS Françoise	KLEIN Bernard	PELLISSIER Jacques
AVEROUS Michel	DEMAILLE Jacques	LAFFARGUE François	PETIT Pierre
AYRAL Guy	DESCOMPS Bernard	LALLEMANT Jean Gabriel	POUGET Régis
BAILLAT Xavier	DIMEGLIO Alain	LAMARQUE Jean-Louis	PUJOL Henri
BALDET Pierre	DUBOIS Jean Bernard	LAPEYRIE Henri	RABISCHONG Pierre
BALDY-MOULINIER Michel	DUJOLS Pierre	LEROUX Jean-Louis	RAMUZ Michel
BALMES Jean-Louis	DUMAS Robert	LESBROS Daniel	REBOUL Jean
BANSARD Nicole	DUMAZER Romain	LOPEZ François Michel	RIEU Daniel
BAYLET René	ECHENNE Bernard	LORIOT Jean	ROCHEFORT Henri
BILLIARD Michel	FABRE Serge	LOUBATIERES Marie Madeleine	ROUANET DE VIGNE LAVIT Jean Pierre
BLARD Jean-Marie	FREREBEAU Philippe	MAGNAN DE BORNIER Bernard	SAINT AUBERT Bernard
BLAYAC Jean Pierre	GALIFER René Benoît	MARTY ANE Charles	SANCHO-GARNIER Hélène
BLOTMAN Francis	GODLEWSKI Guilhem	MARY Henri	SANY Jacques
BONNEL François	GRASSET Daniel	MATHIEU-DAUDE Pierre	SEGNARBIEUX François
BOURGEOIS Jean-Marie	GUILHOU Jean-Jacques	MEYNADIER Jean	SENAC Jean-Paul
BOUSQUET Jean	GUI TER Pierre	MICHEL François-Bernard	SERRE Arlette
BRUEL Jean Michel	HEDON Berbard	MION Charles	SOLASSOL Claude
BUREAU Jean-Paul	HERTAULT Jean	MION Henri	VIDAL Jacques
			VISI ER Jean Pierre

Professeurs Emérites

ARTUS Jean-Claude	LE QUELLEC Alain
BLANC François	MARES Pierre
BONAFE Alain	MAUDELONDE Thierry
BOULENGER Jean-Philippe	MAURY Michèle
BOURREL Gérard	MESSNER Patrick
BRINGER Jacques	MILLAT Bertrand
CLAUSTRES Mireille	MONNIER Louis
DAURES Jean-Pierre	MOURAD Georges
DAUZAT Michel	PREFAUT Christian
DAVY Jean-Marc	PUJOL Rémy
DEDET Jean-Pierre	RIBSTEIN Jean
ELEDJAM Jean-Jacques	SCHVED Jean-François
GROLLEAU RAOUX Robert	SULTAN Charles
GUERRIER Bernard	TOUCHON Jacques
GUILLOT Bernard	UZIEL Alain
JONQUET Olivuer	VOISIN Michel
LANDAIS Paul	ZANCA Michel
LARREY Dominique	

Docteurs Emérites

PRAT Dominique
PUJOL Joseph

ANNEE UNIVERSITAIRE 2020 - 2021

PERSONNEL ENSEIGNANT

Professeurs des Universités - Praticiens Hospitaliers

PU-PH de classe exceptionnelle

ALRIC Pierre	Chirurgie vasculaire ; médecine vasculaire (option chirurgie vasculaire)
BACCINO Eric	Médecine légale et droit de la santé
BASTIEN Patrick	Parasitologie et mycologie
BEREGI Jean-Paul	Radiologie et imagerie médicale
BLAIN Hubert	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
BOULOT Pierre	Gynécologie-obstétrique ; gynécologie médicale
CAPDEVILA Xavier	Anesthésiologie-réanimation et médecine péri-opératoire
CHAMMAS Michel	Chirurgie orthopédique et traumatologique
COLSON Pascal	Anesthésiologie-réanimation et médecine péri-opératoire
COMBE Bernard	Rhumatologie
COSTES Valérie	Anatomie et cytologie pathologiques
COTTALORDA Jérôme	Chirurgie infantile
COUBES Philippe	Neurochirurgie
COURTET Philippe	Psychiatrie d'adultes ; addictologie
CRAMPETTE Louis	Oto-rhino-laryngologie
CRISTOL Jean Paul	Biochimie et biologie moléculaire
CYTEVAL Catherine	Radiologie et imagerie médicale
DE LA COUSSAYE Jean Emmanuel	Médecine d'urgence
DE WAZIERES Benoît	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
DELAPORTE Eric	Maladies infectieuses ; maladies tropicales
DEMOLY Pascal	Pneumologie ; addictologie

DOMERGUE Jacques	Chirurgie viscérale et digestive
DUFFAU Hugues	Neurochirurgie
ELIAOU Jean François	Immunologie
FABRE Jean Michel	Chirurgie viscérale et digestive
FRAPIER Jean-Marc	Chirurgie thoracique et cardiovasculaire
HAMAMAH Samir	Biologie et Médecine du développement et de la reproduction ; gynécologie médicale
HERISSON Christian	Médecine physique et de réadaptation
JABER Samir	Anesthésiologie-réanimation et médecine péri-opératoire
JEANDEL Claude	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
JORGENSEN Christian	Thérapeutique ; médecine d'urgence ; addictologie
KOTZKI Pierre Olivier	Biophysique et médecine nucléaire
LABAUGE Pierre	Neurologie
LEFRANT Jean-Yves	Anesthésiologie-réanimation et médecine péri-opératoire
LEHMANN Sylvain	Biochimie et biologie moléculaire
LUMBROSO Serge	Biochimie et Biologie moléculaire
MERCIER Jacques	Physiologie
MEUNIER Laurent	Dermato-vénéréologie
MONDAIN Michel	Oto-rhino-laryngologie
MORIN Denis	Pédiatrie
PAGEAUX Georges-Philippe	Gastroentérologie ; hépatologie ; addictologie
PUJOL Pascal	Biologie cellulaire
QUERE Isabelle	Chirurgie vasculaire ; médecine vasculaire (option médecine vasculaire)
RENARD Eric	Endocrinologie, diabète et maladies métaboliques ; gynécologie médicale
REYNES Jacques	Maladies infectieuses, maladies tropicales
RIPART Jacques	Anesthésiologie-réanimation et médecine péri-opératoire
ROUANET Philippe	Cancérologie ; radiothérapie

SOTTO Albert	Maladies infectieuses ; maladies tropicales
TAOUREL Patrice	Radiologie et imagerie médicale
TOUITOU Isabelle	Génétique
VANDE PERRE Philippe	Bactériologie-virologie ; hygiène hospitalière
YCHOU Marc	Cancérologie ; radiothérapie

PU-PH de 1^{re} classe

AGUILAR MARTINEZ Patricia	Hématologie ; transfusion
ASSENAT Éric	Gastroentérologie ; hépatologie ; addictologie
AVIGNON Antoine	Nutrition
AZRIA David	Cancérologie ; radiothérapie
BAGHDADLI Amaria	Pédopsychiatrie ; addictologie
BLANC Pierre	Gastroentérologie ; hépatologie ; addictologie
BORIE Frédéric	Chirurgie viscérale et digestive
BOURDIN Arnaud	Pneumologie ; addictologie
CAMBONIE Gilles	Pédiatrie
CAMU William	Neurologie
CANOVAS François	Anatomie
CAPTIER Guillaume	Anatomie
CARTRON Guillaume	Hématologie ; transfusion
CAYLA Guillaume	Cardiologie
CHANQUES Gérald	Anesthésiologie-réanimation et médecine péri-opératoire
CORBEAU Pierre	Immunologie
COULET Bertrand	Chirurgie orthopédique et traumatologique
CUVILLON Philippe	Anesthésiologie-réanimation et médecine péri-opératoire
DADURE Christophe	Anesthésiologie-réanimation et médecine péri-opératoire

DAUVILLIERS Yves	Physiologie
DE TAYRAC Renaud	Gynécologie-obstétrique, gynécologie médicale
DE VOS John	Histologie, embryologie et cytogénétique
DEMARIA Roland	Chirurgie thoracique et cardio-vasculaire
DEREURE Olivier	Dermatologie - vénéréologie
DROUPY Stéphane	Urologie
DUCROS Anne	Neurologie
DUPEYRON Arnaud	Médecine physique et de réadaptation
FESLER Pierre	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
GARREL Renaud	Oto-rhino-laryngologie
GENEVIEVE David	Génétique
GUILLAUME Sébastien	Psychiatrie d'adultes ; addictologie
GUIU Boris	Radiologie et imagerie médicale
HAYOT Maurice	Physiologie
HOUEDE Nadine	Cancérologie ; radiothérapie
KLOUCHE Kada	Médecine intensive-réanimation
KOENIG Michel	Génétique
KOUYOUMDJIAN Pascal	Chirurgie orthopédique et traumatologique
LAFFONT Isabelle	Médecine physique et de réadaptation
LAVABRE-BERTRAND Thierry	Histologie, embryologie et cytogénétique
LAVIGNE Jean-Philippe	Bactériologie-virologie ; hygiène hospitalière
LE MOING Vincent	Maladies infectieuses ; maladies tropicales
LECLERCQ Florence	Cardiologie
MARIANO-GOULART Denis	Biophysique et médecine nucléaire
MATECKI Stéfan	Physiologie
MORANNE Olivier	Néphrologie

MOREL Jacques	Rhumatologie
NAVARRO Francis	Chirurgie viscérale et digestive
NOCCA David	Chirurgie viscérale et digestive
PASQUIE Jean-Luc	Cardiologie
PERNEY Pascal	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
PRUDHOMME Michel	Anatomie
PUJOL Jean Louis	Pneumologie ; addictologie
PURPER-OUAKIL Diane	Pédopsychiatrie ; addictologie
ROGER Pascal	Anatomie et cytologie pathologiques
TRAN Tu-Anh	Pédiatrie
VERNHET Hélène	Radiologie et imagerie médicale

PU-PH de 2ème classe

BOURGIER Céline	Cancérologie ; radiothérapie
CANAUD Ludovic	Chirurgie vasculaire ; médecine vasculaire (option chirurgie vasculaire)
CAPDEVIELLE Delphine	Psychiatrie d'Adultes ; addictologie
CLARET Pierre-Géraud	Médecine d'urgence
COLOMBO Pierre-Emmanuel	Cancérologie ; radiothérapie
COSTALAT Vincent	Radiologie et imagerie médicale
DAIEN Vincent	Ophtalmologie
DORANDEU Anne	Médecine légale et droit de la santé
FAILLIE Jean-Luc	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
FUCHS Florent	Gynécologie-obstétrique ; gynécologie médicale
GABELLE DELOUSTAL Audrey	Neurologie
GAUJOUX Viala Cécile	Rhumatologie
GODREUIL Sylvain	Bactériologie-virologie ; hygiène hospitalière

GUILPAIN Philippe	Médecine Interne, gériatrie et biologie du vieillissement ; addictologie
HERLIN Christian	Chirurgie plastique, reconstructrice et esthétique, brulologie
IMMEDIATO DAIEN Claire	Rhumatologie
JACOT William	Cancérologie ; Radiothérapie
JEZIORSKI Eric	Pédiatrie
JUNG Boris	Médecine intensive-réanimation
KALFA Nicolas	Chirurgie infantile
LACHAUD Laurence	Parasitologie et mycologie
LALLEMANT Benjamin	Oto-rhino-laryngologie
LE QUINTREC DONNETTE Moglie	Néphrologie
LETOUZEY Vincent	Gynécologie-obstétrique ; gynécologie médicale
LONJON Nicolas	Neurochirurgie
LOPEZ CASTROMAN Jorge	Psychiatrie d'Adultes ; addictologie
LUKAS Cédric	Rhumatologie
MENJOT de CHAMPFLEUR Nicolas	Radiologie et imagerie médicale
MILLET Ingrid	Radiologie et imagerie médicale
MURA Thibault	Biostatistiques, informatique médicale et technologies de la communication
NAGOT Nicolas	Biostatistiques, informatique médicale et technologies de la communication
OLIE Emilie	Psychiatrie d'adultes ; addictologie
PANARO Fabrizio	Chirurgie viscérale et digestive
PARIS Françoise	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PELLESTOR Franck	Histologie, embryologie et cytogénétique
PEREZ MARTIN Antonia	Chirurgie vasculaire ; médecine vasculaire (option médecine vasculaire)
POUDEROUX Philippe	Gastroentérologie ; hépatologie ; addictologie
RIGAU Valérie	Anatomie et cytologie pathologiques
RIVIER François	Pédiatrie

ROSSI Jean François	Hématologie ; transfusion
ROUBILLE François	Cardiologie
SEBBANE Mustapha	Médecine d'urgence
SIRVENT Nicolas	Pédiatrie
SOLASSOL Jérôme	Biologie cellulaire
STOEBNER Pierre	Dermato-vénéréologie
SULTAN Ariane	Nutrition
THOUVENOT Éric	Neurologie
THURET Rodolphe	Urologie
TUAILLON Edouard	Bactériologie-virologie; hygiène hospitalière
VENAIL Frédéric	Oto-rhino-laryngologie
VILLAIN Max	Ophtalmologie
VINCENT Denis	Médecine interne ; gériatrie et biologie du vieillissement, médecine générale, addictologie
VINCENT Thierry	Immunologie
WOJTUSCISZYN Anne	Endocrinologie-diabétologie-nutrition

PROFESSEURS DES UNIVERSITES

1^{re} classe :

COLINGE Jacques (Cancérologie, Signalisation cellulaire et systèmes complexes)

LAOUDJ CHENIVESSE Dalila (Biochimie et biologie moléculaire)

VISIER Laurent (Sociologie, démographie)

PROFESSEURS DES UNIVERSITES - Médecine générale

1^{re} classe :

LAMBERT Philippe

AMOUYAL Michel

PROFESSEURS ASSOCIES - Médecine Générale

CLARY Bernard

DAVID Michel

GARCIA Marc

PROFESSEURS ASSOCIES - Médecine

BESSIS Didier (Dermato-vénérologie)

MEUNIER Isabelle (Ophtalmologie)

MULLER Laurent (Anesthésiologie-réanimation et médecine péri-opératoire)

PERRIGAULT Pierre-François (Anesthésiologie-réanimation et médecine péri-opératoire)

QUANTIN Xavier (Pneumologie)

ROUBERTIE Agathe (Pédiatrie)

VIEL Eric (Soins palliatifs et traitement de la douleur)

ANNEE UNIVERSITAIRE 2020 - 2021

PERSONNEL ENSEIGNANT

Maîtres de Conférences des Universités - Praticiens Hospitaliers

MCU-PH Hors classe - Echelon Exceptionnel

RICHARD Bruno	Médecine palliative
SEGONDY Michel	Bactériologie-virologie ; hygiène hospitalière

MCU-PH Hors classe

BADIOU Stéphanie	Biochimie et biologie moléculaire
BOULLE Nathalie	Biologie cellulaire
CACHEUX-RATABOUL Valère	Génétique
CARRIERE Christian	Bactériologie-virologie ; hygiène hospitalière
CHARACHON Sylvie	Bactériologie-virologie ; hygiène hospitalière
FABBRO-PERAY Pascale	Epidémiologie, économie de la santé et prévention
GIANSILY-BLAIZOT Muriel	Hématologie ; transfusion

MCU-PH de 1^{re} classe

BERTRAND Martin	Anatomie
BOUDOUSQ Vincent	Biophysique et médecine nucléaire
BRET Caroline	Hématologie biologique
BROUILLET Sophie	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
COSSEE Mireille	Génétique
GIRARDET-BESSIS Anne	Biochimie et biologie moléculaire
LAVIGNE Géraldine	Hématologie ; transfusion
LESAGE François-Xavier	Médecine et Santé au Travail
MARTRILLE Laurent	Médecine légale et droit de la santé
MATHIEU Olivier	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie

MOUZAT Kévin	Biochimie et biologie moléculaire
PANABIÈRES Catherine	Biologie cellulaire
RAVEL Christophe	Parasitologie et mycologie
SCHUSTER-BECK Iris	Physiologie
STERKERS Yvon	Parasitologie et mycologie
THEVENIN-RENE Céline	Immunologie

MCU-PH de 2^{ème} classe

BERGOUGNOUX Anne	Génétique
CHIRIAC Anca	Immunologie
DE JONG Audrey	Anesthésiologie-réanimation et médecine péri-opératoire
DU THANH Aurélie	Dermato-vénérologie
FITENI Frédéric	Cancérologie ; radiothérapie
GOUZI Farès	Physiologie
HERRERO Astrid	Chirurgie viscérale et digestive
HUBERLANT Stéphanie	Gynécologie-obstétrique ; Gynécologie médicale
KUSTER Nils	Biochimie et biologie moléculaire
MAKINSON Alain	Maladies infectieuses, Maladies tropicales
PANTEL Alix	Bactériologie-virologie ; hygiène hospitalière
PERS Yves-Marie	Thérapeutique ; addictologie
ROUBILLE Camille	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
SZABLEWSKY	Anatomie et cytologie pathologiques

Maîtres de Conférences des Universités - Médecine Générale

MCU-MG de 1^{re} classe

COSTA David

OUDE ENGBERINK Agnès

MCU-MG de 2^{ème} classe

FOLCO-LOGNOS Béatrice

CARBONNEL François

Maîtres de Conférences associés - Médecine Générale

CAMPAGNAC Jérôme

LOPEZ Antonio

MILLION Elodie

PAVAGEAU Sylvain

REBOUL Marie-Catherine

SERAYET Philippe

ANNEE UNIVERSITAIRE 2020 - 2021

PERSONNEL ENSEIGNANT

Maitres de Conférences des Universités

Maîtres de Conférences hors classe

BADIA Eric	Sciences biologiques fondamentales et cliniques
CHAZAL Nathalie	Biologie cellulaire

Maîtres de Conférences de classe normale

BECAMEL Carine	Neurosciences
BERNEX Florence	Physiologie
CHAUMONT-DUBEL Séverine	Sciences du médicament et des autres produits de santé
DELABY Constance	Biochimie et biologie moléculaire
GUGLIELMI Laurence	Sciences biologiques fondamentales et cliniques
HENRY Laurent	Sciences biologiques fondamentales et cliniques
HERBET Guillaume	Neurosciences
LADRET Véronique	Mathématiques appliquées et applications des mathématiques
LAINE Sébastien	Sciences du Médicament et autres produits de santé
LE GALLIC Lionel	Sciences du médicament et autres produits de santé
LOZZA Catherine	Sciences physico-chimiques et technologies pharmaceutiques
MAIMOUN Laurent	Sciences physico-chimiques et ingénierie appliquée à la santé
MOREAUX Jérôme	Science biologiques, fondamentales et cliniques
MORITZ-GASSER Sylvie	Neurosciences
MOUTOT Gilles	Philosophie
PASSERIEUX Emilie	Physiologie
RAMIREZ Jean-Marie	Histologie
RAYNAUD Fabrice	Sciences du Médicament et autres produits de santé
TAULAN Magali	Biologie Cellulaire

ANNEE UNIVERSITAIRE 2020 - 2021

PERSONNEL ENSEIGNANT

Praticiens Hospitaliers Universitaires

BARATEAU Lucie	Physiologie
BASTIDE Sophie	Epidémiologie, économie de la santé et prévention
CAZAUBON Yoann	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
DAGNEAUX Louis	Chirurgie orthopédique et traumatologique
DUFLOS Claire	Biostatistiques, informatique médicale et technologies de la communication
GOULABCHAND Radjiv	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
LATTUCA Benoit	Cardiologie
MARIA Alexandre	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
MIOT Stéphanie	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
SARRABAY Guillaume	Génétique
SOUICHE François-Régis	Chirurgie viscérale et digestive

ANNEE UNIVERSITAIRE 2020 - 2021

PERSONNEL ENSEIGNANT

PH chargés d'enseignements

ABOUKRAT Patrick	BLANCHET Catherine	COROIAN Flavia-Oana	GINIES Patrick
AKKARI Mohamed	BLATIERE Véronique	COUDRAY Sarah	GRECO Frédéric
ALRIC Jérôme	BOBBIA Xavier	CRANSAC Frédéric	GUEDJ Anne Marie
AMEDRO Pascal	BOGE Gudrun	CUNTZ Danielle	GUYON Gaël
AMOUREUX Cyril	BOURRAIN Jean Luc	DARDALHON Brigitte	HENRY Vincent
ANTOINE Valéry	BOUYABRINE Hassan	DAVID Aurore	JAMMET Patrick
ARQUIZAN Caroline	BRINGER-DEUTSCH Sophie	DE BOUTRAY Marie	JEDRYKA François
ATTALIN Vincent	BRINGUIER BRANCHEREAU Sophie	DE LA TRIBONNIÈRE Xavier	JREIGE Riad
AYRIGNAC Xavier	BRISOT Dominique	DEBIEN Blaise	KINNE Mélanie
BADR Maliha	BRONER Jonathan	DELPONT Marion	LABARIAS Coralie
BAIS Céline	CADE Stéphane	DENIS Hélène	LACAMBRE Mathieu
BARBAR Saber Davide	CAIMMI Davide Paolo	DEVILLE de PERIERE Gilles	LANG Philippe
BASSET Didier	CARR Julie	DJANIKIAN Flora	LAZERGES Cyril
BATIFOL Dominique	CARTIER César	DONNADIEU-RIGOLE Hélène	LE GUILLOU Cédric
BATTISTELLA Pascal	CASPER Thierry	FAIDHERBE Jacques	LEGLISE Marie Suzanne
BAUCHET Luc	CASSINOTTO Christophe	FATTON Brigitte	LOPEZ Régis
BENEZECH Jean-Pierre	CATHALA Philippe	FAUCHERRE Vincent	LUQUIENS Amandine
BENNYS Karim	CAZABAN Michel	FILLERON Anne	MANZANERA Cyril
BERNARD Nathalie	CHARBIT Jonathan	FITENI Frédéric	MARGUERITTE Emmanuel
BERTCHANSKY Ivan	CHEVALLIER Thierry	FOURNIER Philippe	MARTIN Lucille
BIBOULET Philippe	CHEVALLIER-MICHAUD Josyane	GAILLARD Nicolas	MATTATIA Laurent
BIRON-ANDREANI Christine	COLIN Olivier	GALMICHE Sophie	MEROUEH Fadi
BLANC Brigitte	CONSEIL Mathieu	GENY Christian	MEYER Pierre
BLANCHARD Sylvie	CORBEAU Catherine	GERONIMI Laetitia	MILESI Christophe

MORAU Estelle	SASSO Milène
MOSER Camille	SCHULDINER Sophie
MOUSTY Eve	SEGURET Fabienne
MOUTERDE Gaël	SENESSE Pierre
PANSARD Nicole	SKALLI EI Medhi
PERNIN Vincent	SOLA Christelle
PERRIGAULT Pierre François	SOULLIER Camille
PEYRON Pierre-Antoine	STOEBNER DELBARRE Anne
PICARD Eric	TEOT Luc
PICOT Marie Christine	THIRION Marina
PIERONI Laurence	VACHIERY-LAHAYE Florence
POQUET Hélène	VERNES Eric
PUJOL Sarah-Lise	VINCENT Laure
PUPIER Florence	WAGNER Laurent
QUANTIN Xavier	ZERKOWSKI Laetitia
RAFFARD Laurence	
RAPIDO Francesca	
RIBRAULT Alice	
RICHAUD-MOREL Brigitte	
RIDOLFO Jérôme	
RIPART Sylvie	
RONGIERES Michel	
ROULET Agnès	
RUBENOVITCH Josh	
SANTONI Fannie	

REMERCIEMENTS

A mon jury,

A Monsieur le Professeur Jorge LOPEZ CASTROMAN,

Je tiens à te remercier chaleureusement pour m'avoir proposé cette thèse puis dirigé avec la plus grande patience et bienveillance. Merci de ta confiance, de ton accompagnement et de ta gentillesse à la fois dans ce travail, qui me semblait dans un premier temps insurmontable, mais également dans le service de gérontopsychiatrie à Nîmes. Je suis content d'avoir pu participer à ma manière à un projet de recherche à tes cotés.

A Madame le Professeur Emilie OLIE,

Vous me faites l'honneur d'être dans mon jury. Merci pour votre présence et vos conseils lors de mes passages à l'UPUP et à la Traversière. Merci également pour votre présence lors des staffs à Lapeyronie qui nous pousse sans cesse à nous améliorer et à aller plus loin dans notre réflexion clinique et théorique.

A Monsieur le Docteur Mathieu LACAMBRE,

Un immense merci pour votre présence et votre confiance lors de mon passage à l'Unité sanitaire. Vos conseils de prise en charge m'ont été précieux. Vous m'avez sensibilisé à la finesse clinique et fait découvrir l'art de l'entretien psychothérapeutique. J'ai pu apprécier à leur juste valeur nos discussions les mardis après-midi au CRIAVS. Je suis extrêmement reconnaissant que vous puissiez participer au jury de ma thèse. J'espère que nos chemins se recroiseront.

Aux soignants rencontrés au fil de ma formation,

Aux équipes Uzétiennes,

Aux Docteurs Guylaine ROCHE, Jérôme BOBO et Jérôme LAVAL et à l'ensemble des équipes pour leur gentillesse et leur initiation à la psychiatrie. Que de bons moments passés au Mas Careiron.

Aux équipes Nîmoises,

Au Docteur Ariane MOLENAT pour ton accueil dans le monde de la pédopsychiatrie et des adolescents, ce fut un vrai retour en enfance. Ta douceur et ton investissement furent très agréables.

Aux Docteurs Ismaël CONEJERO et Jorge LOPEZ CASTROMAN pour votre accompagnement en gérontopsychiatrie et vos enseignements à Nîmes. Merci à Christine BENISTAND pour ton aide dans la prise en charge des problèmes somatiques, sans toi cela aurait été bien plus complexe.

A Mathilde SOUCHON, pour nos discussions et nos repas partagés à la cafet'.

Aux équipes Montpelliéraines,

Aux équipes de l'UPUP, Docteurs Lucile VILLAIN et Chloé GIROD et à toute l'équipe soignante (Sylvie, Marie-Christine, Philippe, Jocelyn, Sabrina, Marie...) pour vos qualités professionnelles et humaines, dans un stage qui fut éprouvant mais enrichissant. Je n'oublie pas l'équipe suicide « le top du top, le haut du panier » avec mon cher Dimitri FIEDOS, vivement qu'on se le boive ce verre ! Je remercie également les équipes rencontrées aux urgences, les Docteurs Radu FLOREA, Audrey HENRION, Marie GACHET, Edith DURET pour votre bienveillance et votre dynamisme.

A la famille de la Traversière, avec qui j'ai pu grandement m'épanouir tant sur le plan professionnel, de la formation mais également personnel. Ce fut un réel plaisir de partager tous ces moments avec vous. Merci au Docteur Blandine DEFFINIS pour ta supervision, ta présence sans faille et ta gentillesse. Merci au Docteur Lionnelle SCHENCK pour votre accessibilité, votre humour et votre sens clinique. C'est toujours un plaisir de vous voir lors de mes passages fugaces sur l'unité.

Au Professeur Eric BACCINO pour votre accueil en médecine légale (et du vivant !), votre bienveillance et vos connaissances. Je garde un bon souvenir de nos échanges et de vos idées toutes défendues avec conviction. Merci pour votre enseignement et votre expertise. Peut-être que nous aurons l'occasion de collaborer ensemble dans le futur. J'espère que la grenouille va bien !

A toute l'Unité Sanitaire de VLM, merci pour votre accompagnement. Je remercie grandement Cyril MANZANERA pour ta supervision, ta précision clinique et tes stratégies thérapeutiques. Merci de m'avoir fait découvrir le microcosme des soins en détention. Merci également au Docteur Mathieu LACAMBRE pour vos conseils et votre accueil sur l'USIP. A toi Pauline, nous avons commencé ensemble à l'Unité Sanitaire, avec les mêmes questionnements sur les prises en charge, ce fut une belle collaboration. Au Docteur Céline BAIS, pour ton sourire,

ton accompagnement des patients et tes présentations formatrices au CRIAVS. Merci à la fabuleuse équipe des psychologues.

Merci au Docteur Françoise CAUSSE, pour ta gentillesse et ton initiation à l'expertise psychiatrique ; tu as su susciter ma volonté de poursuivre dans cette voie.

Enfin merci à tous les membres de l'équipe de l'UJA de Montpellier pour les moments que je passe actuellement avec vous. Merci au Docteur Alexandra CARRE pour ton professionnalisme et ta grande bienveillance, j'espère que je pourrai assister à une CAARMS avant mon départ ! Merci à toi Camille pour ton accompagnement, tes conseils et le temps que tu m'as accordé pour rédiger cette thèse !

A mes amis

A Freddy, Thibaut, Pierre, Florent, Tanneguy, Luc, Nono, Pauline, Kelly, Alex, Adrien, Raphy pour toutes ces années passées ensemble à Lyon et dans nos chambres de l'Ecole de Santé des Armées. A ces soirées passées ensemble qui ont plus ou moins bien finies. Mais également à nos voyages à l'autre bout du monde et quelque accident de voiture au pays des Pyramides.

A toi Florent, dont je suis resté proche, j'espère que l'on pourra continuer à se voir avec Ophélie dans le Sud.

A tous mes amis que j'ai rencontré durant l'internat, en commençant par Lionel. Sacré stage avec toi mon Kiki, que de bonne humeur et de complicité, je te souhaite tout le meilleur pour les heureux événements à venir, mais aussi bien du courage pour ta thèse (quand tu t'y mettras). A la fine équipe d'Uzès : Pauline, Raphael et Noé, il faut que l'on continue à se voir chez Raphael (ou ailleurs), pour se conter nos nouvelles aventures (sanibroyeur et cie) ! A la team de Nîmes : Marion, Vanessa, Charlotte, Charlène, Claire, Pauline (x2), Manu et Kiki, une sacrée rigolade avec vous, je pense qu'on nous regrette encore ! Aux Montpelliérains : Rémi, Suzy, Elodie, Sarah, Emma.

Aux cadurciens et d'autres contrées lointaines, nous nous sommes rencontrés il y a peu de temps, mais pourvu que cela se poursuive encore longtemps, avec modération bien sûr !

A ma famille

Merci à toute ma famille restée en Bretagne. A mes parents, Isabelle et Bernard et à ma sœur Anne-Sophie, qui m'ont soutenu en toutes circonstances au gré de mes péripéties et changement de parcours, malgré la distance. A mes grands-parents, à qui je pense fort. A Hubert et Marie-Hélène qui ont toujours été présents dans la famille mais également dans mon parcours universitaire. Sachez que malgré la distance je pense à vous, j'espère que vous pourrez venir nous voir dans le Sud.

Enfin, merci à mon amour Charline pour ta tendresse, ton soutien et ta folie. Tu auras su être patiente malgré la longueur des études, sache que cela n'est que le début d'une longue aventure.

SOMMAIRE

PERSONNEL ENSEIGNANT	3
REMERCIEMENTS	20
SOMMAIRE	24
Abstract	25
Introduction	26
Rationnel	26
Objectif	27
INTRODUCTION	28
Rationale	28
Objective	29
METHODS	29
Study design and data source	29
Study population	30
Study variables	31
Statistical analysis	31
RESULTS	32
Sample description by sex	32
Comparison by fatal outcome	33
Discharge to a psychiatric service	34
Incidence	34
DISCUSSION	35
Strengths and Limitations of our Study	39
Conclusions	39
CONCLUSIONS	41
REFERENCES	42
TABLES	46
APPENDIX	60
RESUME	62
MOTS-CLES	62

A COMPREHENSIVE STUDY OF MEDICALLY SERIOUS SUICIDE ATTEMPTS HOSPITALIZED IN FRANCE: INCIDENCE AND ASSOCIATED FACTORS

Julien Corbé, Christine Montout, Asma Fares, Ihssen Belhadj, Thierry Boudemaghe, Thibault Mura, Jorge Lopez-Castroman, MD, Ph.D

Abstract

Background: People who make medically serious suicide attempts (MSSAs) share a number of features with those who complete suicide, and are at a high risk of suicide themselves. Studies to date have mostly focused on clinical samples; an epidemiological examination at the national level will help to establish risk profiles and pathways of care in this population.

Methods: We explored the French nationwide hospital discharge database (Programme de Médicalisation des Systèmes d'Information, PMSI) to identify any MSSA taking place between 2012 and 2019. Relevant demographic and medical information were collected about the first MSSA of each attempter with a two-year washout period.

Results: First occurrences of MSSAs amounted to 81 959 cases over 8 years, with a mean age of 45.8, and 53.6% women. Incidence was higher in women (18.1 versus 17.3 per 100 000). The most common suicide method was deliberate self-poisoning (64.9% of cases). In comparison, violent methods associated higher mortality and comorbidity and were more frequent in men. The most common mental disorders were mood disorders (55.6%) and substance use disorders (46.2%). A minority of MSSA survivors were hospitalized in psychiatry (32.5%), mostly women.

Conclusions: MSSAs are frequent and easy to identify. There is a need to reinforce the continuity of psychiatric care for this population given the high risk of subsequent suicide, and the poor rates of psychiatric hospitalization after an MSSA. Specific care targeting this population could reduce treatment gaps.

Introduction

Rationnel

On estime que pour chaque décès par suicide, il y a environ, selon les pays, 10 à 30 tentatives de suicide (TS) non mortelles. Un antécédent de TS constitue le meilleur prédicteur de suicide puisqu'environ 40 % des victimes de suicide ont des antécédents de tentatives, multipliant le risque de suicide dans l'année par 49. Cependant, toutes les TS ne véhiculent pas le même risque.

Une tentative de suicide médicalement grave (medically serious suicide attempt (MSSA)) a été définie comme une TS qui aurait été fatale sans l'accès aux soins d'urgence et qui a ensuite nécessité une hospitalisation de plus de 24 heures dans une unité de soins intensifs ou de réanimation, ou une intervention chirurgicale sous anesthésie générale, quelle que soit la violence de l'acte. Les personnes qui font une MSSA sont phénotypiquement plus proches de celles qui se suicident. Elles sont plus âgées, plus susceptibles d'avoir des antécédents de TS, et rapportent une létalité plus élevée lors de TS antérieures par rapport aux individus n'ayant jamais fait de MSSA. Les troubles mentaux et les méthodes de suicide dans les MSSAs sont également à mi-chemin entre les suicides et les TS. De nombreuses études rapportent que l'auto-intoxication est majoritaire dans les échantillons de MSSA (57-79%) mais que les méthodes violentes comme la pendaison sont surreprésentées (10-17%). Le trouble bipolaire, le trouble de l'usage des substances et le trouble du comportement alimentaire sont également surreprésentés dans les groupes de MSSA en comparaison des TS avec des conséquences moins sérieuses. A l'inverse, les diagnostics de psychose non affective semblent être beaucoup plus fréquents dans les échantillons de suicidés (OR=8,5) que dans les échantillons de MSSA (OR=8,5).

Parmi les personnes ayant fait une tentative de suicide, le fait de faire une MSSA double le risque de se suicider à court terme. Un individu sur vingt-cinq (4%) se suicide dans les 18 mois qui suivent une MSSA, mais seulement 1,8% dans l'année qui suit une TS. Ceci est également vrai à moyen terme : 5,3% à 7% de suicides sur 5 ans après une MSSA contre 3,8% sur 5 ans pour une TS. La mortalité globale par suicide après une MSSA serait également multipliée par 5 par rapport

à la population générale. Il est important de noter que l'évaluation directe des individus qui ont fait une MSSA, et qui auraient pu mourir par suicide, fournit des informations sur les troubles mentaux, les processus cognitifs ou les traits psychologiques dans les décès par suicide.

Nous disposons de peu d'informations épidémiologiques sur les MSSAs. Seules quelques études cliniques ont examiné la comorbidité somatique et psychiatrique associée aux MSSAs.

Objectif

Dans notre étude, nous avons voulu décrire d'un point de vue démographique et clinique la population qui fait une première MSSA et son parcours de soins hospitalier au niveau national. Nous avons également calculé l'incidence des MSSAs car, à notre connaissance, cela n'a jamais été fait auparavant. En effet, la plupart des études précédentes ont été réalisées sur des échantillons cliniques ne dépassant pas 1500 individus. Notre base de données fournit des informations exhaustives sur l'ensemble de la population française admise dans les hôpitaux entre 2012 et 2019. En raison de la faible quantité de données disponibles sur ce sujet et du profil à haut risque des survivants de MSSA, nous pensons que cette étude fournira des informations utiles aux soignants dans la gestion des comportements suicidaires.

INTRODUCTION

Rationale

It is estimated that for every suicide death there are about 10 to 30, depending on the country, non-lethal suicide attempts (SAs) (Bachmann, 2018; Han et al., 2016; Kessler et al., 2005; Mościcki et al., 1988). Having attempted suicide constitutes the best predictor of suicide since about 40% of victims of suicide have records of previous attempts (Hawton et al., 2005; Hawton and van Heeringen, 2009), multiplying the risk of suicide in the year by 49 (Hawton et al., 2015). However, not all SAs convey the same risk.

A medically serious suicide attempt (MSSA) has been defined as a SA that would have been fatal without access to emergency care and that subsequently required hospitalization for more than 24 hours in an intensive care unit (ICU), or surgery under general anesthesia, regardless of the violence of the act (Levi-Belz and Beautrais, 2016). Individuals who make MSSA are phenotypically closer to those who complete suicide (Beautrais, 2001; Mościcki, 1995). They are older, more likely to have prior records of SAs, and report higher lethality in previous SAs compared to attempters that never made a MSSA (Giner et al., 2014). Mental disorders and suicide methods in MSSAs are also half-way between suicides and SAs. Many studies report that self-poisoning is majoritarian in MSSA samples (57-79%) but violent methods such as hanging are overrepresented (10-17%) (Beautrais, 2003b; Horesh et al., 2012; Kim et al., 2020; Sun et al., 2015). Bipolar disorder, substance misuse and eating disorder are also overrepresented in MSSAs samples in comparison SAs with less serious consequences (Giner et al., 2014). Conversely, the diagnoses of non-affective psychosis seem to be much more common in samples of suicide completers (OR=8.5) than in MSSA samples (OR=8.5) (Beautrais, 2001).

Among suicide attempters, making a MSSA more than doubles the risk of completing suicide in the short term. One in twenty-five individuals (4%) complete suicide in the 18 months that follow a MSSA (Beautrais, 2004), but only 1.8% in the year following a SA (Probert-Lindström et al., 2020). This is also true in the medium term: 5.3% to 7% over 5 years (Beautrais, 2004, 2003a)

versus 3.8% over 5 years (Probert-Lindström et al., 2020). The overall suicide mortality after an MSSA would also be multiplied by 5 compared to the general population (Beautrais, 2003a). Importantly, the direct assessment of individuals who made a MSSA, and could have died by suicide, provides proxy information about mental disorders, cognitive processes or psychological traits in suicide deaths (Clark and Horton-Deutsch, 1992; Hawton, 2001; Hawton et al., 1998).

We have scarce epidemiological information about MSSAs. Only some clinical studies investigated the somatic and psychiatric comorbidity associated to MSSAs.

Objective

Here, we aimed to describe from a demographic and clinical standpoint the population that makes a first MSSA and what use they make of hospital care at a national level. We will also calculate the incidence of MSSAs since, to our knowledge, it has never been done before. Indeed, most previous studies were conducted on clinical samples of no more than 1500 individuals (Beautrais et al., 1996a; Beautrais, 2001, 2004; Kim et al., 2020). Our database provides exhaustive information on the entire French population admitted to hospitals between 2012 and 2019. Because of the small amount of data available on this topic and the high-risk profile of MSSA survivors, we believe that this study will provide useful information to caregivers in the management of suicidal behavior.

METHODS

Study design and data source

We conducted a nationwide observational study in French hospitals using the French nationwide hospital discharge database (Programme de Médicalisation des Systèmes d'Information, PMSI). The PMSI database contains synthesized and anonymized data about, among other kind of care, acute care hospitalizations in France, encompassing medicine, surgery, and obstetrics sector (Medecine, chirurgie, obstétrique (MCO)). MCO sector is made up of all the

medical units of an establishment, providing acute care in medicine, surgery and obstetrics. Conventional psychiatric units are not included in the MCO sector. PMSI data are prospectively collected by all public and private hospitals in a standardized way for care reimbursement purposes. Every discharge summary contains sociodemographic information about the patient, medical information about the hospital stay and data related to the trajectory of the patient. Medical information is coded with the Tenth revision of the International Classification of Diseases (ICD-10).

National PMSI data are anonymized. Access to these data is authorized for research purpose, and does not require the individual information and written consent of the patients. Approval of the National Data Protection Commission (Commission Nationale Informatique et Liberte, CNIL) was obtained for this study and data were handled and analyzed on the secured electronic platform of the National agency for the information on hospitalizations (Agence Technique de l'Information sur l'Hospitalisation, ATIH).

Study population

All the inpatient admissions related to a first episode of MSSA in the acute care sector of French hospitals between 2010 and 2019 were considered for the present study. A hospital stay for a MSSA implied: 1) the presence of an intentional self-harm ICD-10 code X60-X84 in the discharge summary; 2) the admission to an ICU for somatic conditions, including critical care units; and, 3) the absence of previous hospitalizations caused by a MSSA for at least the two preceding years. The two-year period comprises the years 2010 and 2011 for MSSA occurred in the year 2012, or the longest period available since 2010 for MSSA from 2012. To ensure that we were studying a homogenous sample, only first admissions fulfilling the criteria for MSSA were selected.

Study variables

1. Patients' demographics (sex and age), and comorbidities according to the Charlson index algorithm. The Charlson Comorbidity Index is a method of categorizing a range of non-psychiatric comorbidities of patients based on the ICD to predict short-term mortality. The one year mortality rates for the different scores are: "0", 12%; "1-2", 26%; "3-4", 52%; and ">5", 85% (Charlson et al., 1987).
2. Medical variables: primary/main and secondary diagnosis coded with the ICD-10 related to organic and psychiatric comorbidities, method of the SA. To summarize diagnostic data we used ICD-10 chapter codes for mental disorders (F0 to F9). Three specific diagnoses were analyzed separately because of their strong association with suicide risk: bipolar disorder (F30-F31), alcohol use disorder (AUD) (F10) and schizophrenia (F20).
3. Variables related to the patients' intra and inter-hospital trajectory: lengths of stay, admission in ICU, origin of referral and destination, vital status at discharge.
4. Admission in a psychiatric unit at least 3 days after discharge.

Statistical analysis

A total of 735 416 MSSAs were found between January 1, 2012 and December 31, 2019. Among them, 81 959 first-time MSSAs were identified and analyzed (see data flowchart in Figure 1). As we consider only first MSSAs for the present study, one patient corresponds to one discharge summary.

We achieve exclusively descriptive analysis for their characteristics. Qualitative variables were reported with numbers and percentages and quantitative ones, with means and standard deviations (SD). Pearson χ^2 tests were carried out for qualitative variables. Fischer exact tests were used, as appropriate, for small groups. For quantitative variables, T-tests. were realized.

To estimate the incidence rates of first MSSA in the French general population by sex and age, denominators were INSEE (Institut national de la statistique et des études économiques) figures of the general population and numerators were numbers of first MSSA from this study. Incidences of first MSSA were estimated and reported per person-year overall and among subgroups.

Analyses were performed with a bilateral alpha level of 0.05 using the SAS Enterprise guide 7.15 (SAS Institute, Cary, North Carolina).

RESULTS

Sample description by sex

Most MSSAs cases are women (N=43 953, 53.62%; $p<.0001$), a trend that is particularly evident in extreme age groups (under 25 or over 55 years) (Table 1). In the age group between 25 and 55 years, which counts with the largest caseload (N=46 150, 56.3%), men are overrepresented (N=23 258, 50.4%). MSSAs before 15 are rare in both sexes (N=412, 0.5%).

Deliberate self-poisoning (DSP) by drugs is the most common method of suicide attempt in MSSA (64.9% of the cases), especially among women (72.1% compared to 56.5% among men). Men are more likely to use violent methods such as hanging (7.9% versus 1.9% in women) or firearms (3.1% versus 0.2%). Men are also more likely to die after their first MSSA (68.4% of deaths).

Concerning mental disorders, the most frequent diagnoses are mood disorders (55.6%), including bipolar disorder (7.2%), and substance use disorders (SUD) (46.2%), including alcohol dependence (34.3%). Men seem to suffer significantly more from SUDs (56.3% vs. 37.5%), particularly alcohol dependence (42.8% vs. 27.1%), and schizophrenia (5.5% vs. 2.1%). Women are more concerned by mood disorders (64% vs 45.8%), including bipolar disorder (8.9% vs 5.1%). Personality disorders are balanced (approximately 3% in both genders).

The average length of stay is statistically higher for men, with 5.55 days spent in ICU compared with 4.4 days for women ($p < .0001$). The Charlson Score is also higher in men (4.1% of men with Charlson Score >3 versus 2.5% of women). Women are more likely to enter a psychiatric ward 3 days after their discharge from ICU (N=16 912; 38.5% of women) than men (N=12 309; 32.4% of men).

Comparison by fatal outcome

Following a first MSSA, 4 519 patients died, representing 5.5% of the MSSA population (Table 2). MSSAs are significantly more lethal in men, resulting in death in 8.1% of cases versus 3.3% of women ($p < .0001$). On average, the deceased are older (54.69 ± 19.23 years) than the survivors (45.28 ± 17.41 years). A high case fatality rate is observed over 75 years of age, with 15% of deaths compared to 4.4% in the [25 to 55] age group. The proportion of deceased patients admitted from psychiatry (N=264; 88.6%) is smaller than the proportion of deceased patients admitted from other places ($>93\%$). The number of deaths to MSSA is stable along the study period (every year about 12-13% of the total).

Deliberate self-poisoning (DSP) by drugs appears to be less lethal (1.9% deaths) than any other method (12.2% deaths). Among the deceased, the method chosen to attempt suicide is commonly violent. Hanging/strangulation is used in 38.5% of deaths and 4.7% of MSSAs, firearms are used in 9.8% of deaths and 1.6% of MSSAs. On the contrary, DSP accounts for 67.3% of MSSAs and 22.4% of deaths.

Mortality is particularly important in patients diagnosed with mood disorders or organic mental disorders (F3 and F0) when compared to any other diagnosis. On the contrary, diagnoses of SUD (F1), psychotic disorder (F2) or personality disorder (F6) are less often associated with a fatal outcome. Deceased patients have longer lengths of stay (8.57 ± 20.31 days) than survivors do (7.13 ± 13.23 days). In addition, the Charlson score is statistically higher in the decedents (15.9% with Charlson Score >1 in the decedents vs. 10.1% in the survivors).

Discharge to a psychiatric service

One third of surviving MSSA patients are discharged directly to a psychiatric service (N=25 177, 32.5%), while almost half are discharged directly to their home (N=37 652; 48.6%) (Table 3). These numbers are stable over the study period. Women are more likely to be discharged to psychiatry than men. They represent 58.5% of discharges to a psychiatric service, whereas they constitute 54.9% of survivors. The extreme age groups are less likely to be hospitalized in psychiatry after ICU, less than 25% in the youngest and oldest age groups (8-15 years and over 75 years) compared to more than 30% in the other age groups (25-75 years).

MSSA survivors after DSP are more frequently hospitalized in psychiatry than those using different methods (34.2% vs. 28.9%). Discharges to psychiatric inpatient units for MSSAs using violent methods such as hanging (N=925; 24%) or firearm (N=167;12.9%) are particularly infrequent.

Patients with psychotic disorder (N=1 969; 45.1%) or mood disorder (N=11 843; 37.4%) are more often hospitalized in psychiatry, in contrast to those with SUD (N=7 639 ;28.7%). Nearly three quarters (N=205; 77.7%) of psychiatric inpatients are re-admitted after intensive care in a psychiatric ward. On the other hand, about one-third of MSSA survivors coming from the emergency room (N=17 525; 33.9%) or home (N=6 928; 29.5%) are subsequently admitted to a psychiatric unit.

Patients who have less comorbidities are more discharged to psychiatry (33.2% with Charlson Score=0 vs. 17.7% with Charlson Score>5). They spend less time in intensive care (4.61 ± 7.2 vs. 8.34 ± 15.17 days) than those discharged elsewhere. Methods of attempting suicide are represented in Table 4.

Incidence

We find an overall increase in the incidence of MSSA over the study period, from 16.57 in 2012 to 17.75 in 2019 (cases per 100 000 person-years). This tendency was particularly marked

in females aged 15-25 years with a ten-point increase (Table 5). In women, the incidence is consistently higher than in men, with a trend towards a narrowing of this gap. In 2012, the incidence is 17.42 in women and 15.66 in men. In 2019, 18.14 in women versus 17.33 in men.

DISCUSSION

To our knowledge, this is largest study to date focused on MSSA. It provides an 8-year coverage of first MSSAs in the entire French population aged 8 or more. According to the literature, first MSSAs were mostly made by women and the most common method in both sexes was DSP, although violent suicide methods were over-represented in men (Beautrais, 2003b, 2001; Giner et al., 2014; Kim et al., 2020). Mood disorders (55.6%) and SUDs (46.2%) were by large the most common psychiatric disorders and showed a clear gender pattern, women more often diagnosed with mood disorders and men with SUDs. Importantly, less than one-third of MSSA survivors are hospitalized in psychiatry after discharge and the patients most at risk for completed suicide are the least hospitalized (for instance, men with substance use disorders or using violent methods). The use of violent methods in MSSAs was associated with the highest lethality and comorbidity.

Similar to suicide completers (Bachmann, 2018; Cibis et al., 2012; Värnik et al., 2008), MSSA attempters who die seem to be mostly men over 65 years that use hanging as a method. In our sample death was frequent (one in twenty attempters), representing more than fivefold the death rate by any SA leading to hospitalization in France between 2004 and 2011 (Christine Chan Chee, Delphine Jezewski-Serra, 2014). MSSA perpetrators are older compared to other suicide attempters (Giner et al., 2014; Horesh et al., 2012; Kim et al., 2020), but younger than suicide victims (Beautrais, 2001). Women make more attempts but men die more often which is consistent with the “gender paradox” in suicide (Canetto and Sakinofsky, 1998; Schrijvers et al., 2012). Indeed, men represent 68.4% of deaths in our study, spend more time in intensive care and have

more somatic comorbidity than women. These differences could be explained by the choice of more lethal methods in men (Beautrais, 2003b) but even when using similar methods men have higher mortality rates in our sample. It has been reported that one in ten survivors of a MSSA die within 5 years, mostly by suicide (59%) (Beautrais, 2003a). Overall mortality is sharply increased in the years that follow a violent self-harm, especially the first year, and also among men and physically vulnerable persons (Bergen et al., 2012; Chen et al., 2011; Goldman-Mellor et al., 2019; Stenbacka and Jokinen, 2015; Vuagnat et al., 2019). This may explain the increase in natural mortality.

One in ten patients with MSSA (10.5%) had multiple somatic comorbidities. Coherently, physical illnesses have been found to precipitate MSSA (Kim et al., 2020). Physical sequelae secondary to MSSA may also be independent risk factors for suicide. Patients with somatic pathology complete suicide 2 to 3 times more than general population in Taiwan (Cheng et al., 2000), even more so in the presence of multiple co-morbidities or social isolation due to disability (Ahmedani et al., 2017; Kennedy and Garmon-Jones, 2017). Besides, intensive care survivors (all reasons for hospitalization combined) had higher risk of suicide after discharge than other inpatients (Fernando et al., 2021). In a national population study, the interaction of psychiatric and somatic illnesses increased the risk of suicide (Qin et al., 2014). This trend was even more significant in men and the elderly (Blasco-Fontecilla et al., 2010). The physical consequences of a MSSA can also hinder psychiatric care, especially after a violent SA (Persett et al., 2018).

In our study, only 32.5% of MSSA survivors were subsequently admitted in a psychiatric clinic. Between 2004 and 2011, 20% of all SAs hospitalized in French hospitals (MCO) were subsequently admitted in psychiatry (Christine Chan Chee, Delphine Jezewski-Serra, 2014). These weak rates are partly explained by admissions in psychiatric units within general hospitals, which are not counted in our study. In France these units are mostly conceived to provide emergency psychiatric care in academic hospitals, and despite an increase in recent years they are still uncommon and most psychiatric care takes place in psychiatric clinics (public or private).

Besides, stays in emergency or crisis units are short and most patients in our study were discharged to their home (48.6%). Male sex and advanced age decreased the likelihood of admission in psychiatry after critical care according to our results. In addition, the longer the time spent in ICU, the fewer psychiatric admissions. These patients had also more physical comorbidities and made the most serious attempts. In summary, they had the more severe risk profile.

Non-psychiatric medical departments are not well-adapted to the management of suicide attempters (less communication, untrained teams) (Betz et al., 2013; Chen et al., 2016; Cheng et al., 2009). Non-fatal self-harm attempters admitted to non-psychiatric departments are more likely to complete suicide in the following year than those admitted to psychiatry (Vuagnat et al., 2019). Consultation-Liaison Psychiatry (CLP) should identify and manage these cases to reinforce continuous care (Stein et al., 2020), but the accessibility to CLP varies widely among hospitals (Olfson et al., 2014; Wand et al., 2016; Wood and Wand, 2014). According to one study, when CLP intervenes only 29.2% of violent or serious attempts are discharged (Cooper-Kazaz, 2013). Of note, the risk of suicide seems to be higher among women after a MSSA (Beautrais, 2004) but higher among men after a nonfatal SA (Hawton and Fagg, 1988).

Two out of three MSSA used DSP as a method for suicide. Similar results were found in previous studies with intoxications accounting for 57-79% of MSSA (Beautrais, 2003b; Horesh et al., 2012; Kim et al., 2020; Sun et al., 2015). Suicide attempt methods in our sample are thus close to those of non-lethal SAs if we compare with previous studies: 1) 81.7% of the suicides used a highly lethal method against 17.6% in MSSA (Beautrais, 2003b), 2) DSP accounts for approximately two out of three of SAs not requiring intensive care (Cibis et al., 2012), but 3) only for 12.7% of suicides in Europe (Värnik et al., 2008).

High levels of psychopathology increase the susceptibility to MSSA (Beautrais, 2003b). The diagnostic profile of MSSA patients (mood disorders, SUDs) in our study is similar to previous reports by Beautrais's group, which found high rates of current and lifetime mental

disorder (around 90%), and previous SAs (23.6 to 52.7%) (Beautrais, 2001; Beautrais et al., 1998, 1996b). Post-traumatic stress disorder (PTSD) was also more prevalent among MSSA survivors (Lopez-Castroman et al., 2015). Consistently with previous studies (Beautrais et al., 1999; Gvion and Levi-Belz, 2018), SUDs and particularly AUDs were very frequent in our sample, one in three MSSA survivors (31.1%) had an AUD in the previous month (Beautrais et al., 1999). Alcohol dependence more than doubles the risk of making a MSSA compared to the general population (Conner et al., 2003a), and drinking within 3 hours of a SA causes a 6-fold increase in the risk of MSSA (Powell et al., 2001). In a prior study, the comorbidity of alcohol dependence and a mood disorder had lower odds of making a MSSA compared to a mood disorder alone (OR=6 vs 17), but alcohol-dependent patients making MSSA were more likely to have a mood disorder (Conner et al., 2003a, 2003b). Other SUDs, such as cannabis or tobacco dependence, have been strongly associated with MSSA (Beautrais et al., 1999; Lopez-Castroman et al., 2016).

Strong similarities in psychiatric diagnoses have been shown between MSSAs and suicides: mood disorders, substance use disorders and anxiety disorders are overrepresented compared to the general population (Arsenault-Lapierre et al., 2004; Beautrais, 2001). Nevertheless, non-affective psychosis was a risk factor for suicide and not for MSSA (5.9% vs. 1.1%), whereas anxiety disorders and mood disorders were more present in MSSA (Beautrais, 2003b, 2001). Compared with non-critical care SAs, psychotic disorders, bipolar disorder, eating disorders, SUDs and SA recidivism were overrepresented in MSSA (Giner et al., 2014; Kim et al., 2020).

Given the low rate of psychiatric hospitalizations in our study, it seems essential to improve care pathways for MSSA survivors. The two-year period following discharge after a SA is at high risk for suicide (Parra-Uribe et al., 2017; Tejedor et al., 1999). Continuous psychiatric and somatic outpatient care is very important. A 7-day follow-up after discharge from a psychiatric hospital was associated with an important decrease in suicide rates in the 3-month period following discharge (While et al., 2012). Specific training or sensibilization of non-

psychiatric teams about the risk of suicide and its management is important. Less than 70% of emergency physicians know how to assess the risk of suicide. (Betz et al., 2013).

The incidence rate in our study is lower (45.7 per 100 000 person-years) than in a Chinese study (Sun et al., 2015), which aimed to calculate the incidence rate in 3 counties from two public health surveillance systems during 2009-2011, but this one included deaths of MSSA before their arrival at the hospital. Our figures are lower than SAs (148.8 per 100 000 person-years) according to a US study (Kuo et al., 2001).

Strengths and Limitations of our Study

The main limitation of this study is that it is based on administrative data, ICD-10 codes are mainly used for billing purposes by physicians on the hospital ward. They might vary according to the practitioner or the institution. Besides, in some university hospitals in France, patients are hospitalized in psychiatric departments associated to the emergency care. These cases have not been counted as psychiatric stays and may lead to an underestimation of psychiatric hospitalizations. Also, short stays, especially in services other than intensive care, may raise questions about the severity of the suicidal act, as simple monitoring may be sufficient. Finally, important information such as suicidal intent or records of previous suicidality is not available in the database. However, PMSI data provides exhaustive and updated information on all health establishments at the national level for almost 10 years. To our knowledge, this is the first epidemiological study of this magnitude conducted on MSSA.

Conclusions

In conclusion, the phenotype of MSSA survivors is midway between that of suicide victims and suicide attempters. MSSA survivors are more often female, use more violent methods of suicide, and have more serious somatic and psychiatric pathologies. Given the risk of suicide and somatic complications after a MSSA (Beautrais, 2003a; Chen et al., 2011; Goldman-Mellor et al., 2019; Vuagnat et al., 2019) and the low rate of psychiatric hospitalizations in our study, it

seems essential to improve care pathways for MSSA survivors. This retrospective study needs to be completed by prospective studies mixing somatic and psychiatric aspects in order to better understand the consequences of MSSA and thus adapt the management.

CONCLUSIONS

En conclusion, le phénotype des survivants de MSSA est à mi-chemin entre celui des suicidés et de ceux faisant des tentatives de suicide. En effet, les survivants de MSSA sont plus souvent des femmes, d'un plus grand âge, utilisant des moyens plus violents, décédant plus, et présentant des pathologies psychiatriques et somatiques plus graves que ceux ayant fait une TS moins sérieuse. Compte tenu du haut risque de suicide et de complications somatiques après une MSSA et du faible taux d'hospitalisation en psychiatrie dans notre étude, il semble essentiel d'améliorer les parcours de soins des survivants d'une MSSA en intra-hospitalier mais également en ambulatoire. Cette étude rétrospective doit être complétée par des études prospectives mêlant les aspects somatiques et psychiatriques afin de mieux comprendre les conséquences des MSSAs et ainsi adapter la prise en charge.

REFERENCES

- Ahmedani, B.K., Peterson, E.L., Hu, Y., Rossom, R.C., Lynch, F., Lu, C.Y., Waitzfelder, B.E., Owen-Smith, A.A., Hubley, S., Prabhakar, D., Williams, L.K., Zeld, N., Mutter, E., Beck, A., Tolsma, D., Simon, G.E., 2017. Major Physical Health Conditions and Risk of Suicide. *Am. J. Prev. Med.* 53, 308–315. <https://doi.org/10.1016/j.amepre.2017.04.001>
- Arsenault-Lapierre, G., Kim, C., Turecki, G., 2004. Psychiatric diagnoses in 3275 suicides: a meta-analysis. *BMC Psychiatry* 4, 37. <https://doi.org/10.1186/1471-244X-4-37>
- Bachmann, S., 2018. Epidemiology of Suicide and the Psychiatric Perspective. *Int. J. Environ. Res. Public Health* 15. <https://doi.org/10.3390/ijerph15071425>
- Beautrais, A.L., 2004. Further suicidal behavior among medically serious suicide attempters. *Suicide Life Threat. Behav.* 34, 1–11. <https://doi.org/10.1521/suli.34.1.1.27772>
- Beautrais, A.L., 2003a. Subsequent mortality in medically serious suicide attempts: a 5 year follow-up. *Aust. N. Z. J. Psychiatry* 37, 595–599. <https://doi.org/10.1046/j.1440-1614.2003.01236.x>
- Beautrais, A.L., 2003b. Suicide and serious suicide attempts in youth: a multiple-group comparison study. *Am. J. Psychiatry* 160, 1093–1099. <https://doi.org/10.1176/appi.ajp.160.6.1093>
- Beautrais, A.L., 2001. Suicides and serious suicide attempts: two populations or one? *Psychol. Med.* 31, 837–845. <https://doi.org/10.1017/s0033291701003889>
- Beautrais, A.L., Joyce, P.R., Mulder, R.T., 1999. Cannabis abuse and serious suicide attempts. *Addict. Abingdon Engl.* 94, 1155–1164. <https://doi.org/10.1046/j.1360-0443.1999.94811555.x>
- Beautrais, A.L., Joyce, P.R., Mulder, R.T., 1998. Psychiatric illness in a New Zealand sample of young people making serious suicide attempts. *N. Z. Med. J.* 111, 44–48.
- Beautrais, A.L., Joyce, P.R., Mulder, R.T., Fergusson, D.M., Deavoll, B.J., Nightingale, S.K., 1996a. Prevalence and comorbidity of mental disorders in persons making serious suicide attempts: a case-control study. *Am. J. Psychiatry* 153, 1009–1014. <https://doi.org/10.1176/ajp.153.8.1009>
- Beautrais, A.L., Joyce, P.R., Mulder, R.T., Fergusson, D.M., Deavoll, B.J., Nightingale, S.K., 1996b. Prevalence and comorbidity of mental disorders in persons making serious suicide attempts: a case-control study. *Am. J. Psychiatry* 153, 1009–1014. <https://doi.org/10.1176/ajp.153.8.1009>
- Bergen, H., Hawton, K., Waters, K., Ness, J., Cooper, J., Steeg, S., Kapur, N., 2012. Premature death after self-harm: a multicentre cohort study. *Lancet Lond. Engl.* 380, 1568–1574. [https://doi.org/10.1016/S0140-6736\(12\)61141-6](https://doi.org/10.1016/S0140-6736(12)61141-6)
- Betz, M.E., Sullivan, A.F., Manton, A.P., Espinola, J.A., Miller, I., Camargo, C.A., Boudreaux, E.D., ED-SAFE Investigators, 2013. Knowledge, attitudes, and practices of emergency department providers in the care of suicidal patients. *Depress. Anxiety* 30, 1005–1012. <https://doi.org/10.1002/da.22071>
- Blasco-Fontecilla, H., Baca-Garcia, E., Duberstein, P., Perez-Rodriguez, M.M., Dervic, K., Saiz-Ruiz, J., Courtet, P., de Leon, J., Oquendo, M.A., 2010. An exploratory study of the relationship between diverse life events and specific personality disorders in a sample of suicide attempters. *J. Personal. Disord.* 24, 773–784. <https://doi.org/10.1521/pedi.2010.24.6.773>
- Canetto, S.S., Sakinofsky, I., 1998. The gender paradox in suicide. *Suicide Life Threat. Behav.* 28, 1–23.
- Charlson, M.E., Pompei, P., Ales, K.L., MacKenzie, C.R., 1987. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J. Chronic Dis.* 40, 373–383. [https://doi.org/10.1016/0021-9681\(87\)90171-8](https://doi.org/10.1016/0021-9681(87)90171-8)
- Chen, K.Y., Evans, R., Larkins, S., 2016. Why are hospital doctors not referring to Consultation-Liaison Psychiatry? - a systemic review. *BMC Psychiatry* 16, 390. <https://doi.org/10.1186/s12888-016-1100-6>
- Chen, V.C.H., Tan, H.K.L., Chen, C.-Y., Chen, T.H.H., Liao, L.-R., Lee, C.T.C., Dewey, M., Stewart, R., Prince, M., Cheng, A.T.A., 2011. Mortality and suicide after self-harm: community cohort study in Taiwan. *Br. J. Psychiatry J. Ment. Sci.* 198, 31–36. <https://doi.org/10.1192/bjp.bp.110.080952>
- Cheng, A.T., Chen, T.H., Chen, C.C., Jenkins, R., 2000. Psychosocial and psychiatric risk factors for suicide. Case-control psychological autopsy study. *Br. J. Psychiatry J. Ment. Sci.* 177, 360–365. <https://doi.org/10.1192/bjp.177.4.360>
- Cheng, I.-C., Hu, F.-C., Tseng, M.-C.M., 2009. Inpatient suicide in a general hospital. *Gen. Hosp. Psychiatry* 31, 110–115. <https://doi.org/10.1016/j.genhosppsy.2008.12.008>

- Christine Chan Chee, Delphine Jezewski-Serra, 2014. Hospitalisations et recours aux urgences pour tentative de suicide en France métropolitaine à partir du PMSI-MCO 2004-2011 et d'Oscour® 2007-2011 [WWW Document]. URL <https://www.f2rsmpsy.fr/544-hospitalisations-recours-aux-urgences-pour-tentative-suicide-france-metropolitaine-partir-pmsi-mco-2004-2011-droscou> (accessed 3.29.21).
- Cibis, A., Mergl, R., Bramesfeld, A., Althaus, D., Niklewski, G., Schmidtke, A., Hegerl, U., 2012. Preference of lethal methods is not the only cause for higher suicide rates in males. *J. Affect. Disord.* 136, 9–16. <https://doi.org/10.1016/j.jad.2011.08.032>
- Clark, D.C., Horton-Deutsch, S.L., 1992. Assessment in absentia: The value of the psychological autopsy method for studying antecedents of suicide and predicting future suicides, in: *Assessment and Prediction of Suicide*. Guilford Press, New York, NY, US, pp. 144–182.
- Conner, K.R., Beautrais, A.L., Conwell, Y., 2003a. Moderators of the relationship between alcohol dependence and suicide and medically serious suicide attempts: analyses of Canterbury Suicide Project data. *Alcohol. Clin. Exp. Res.* 27, 1156–1161. <https://doi.org/10.1097/01.ALC.0000075820.65197.FD>
- Conner, K.R., Beautrais, A.L., Conwell, Y., 2003b. Risk factors for suicide and medically serious suicide attempts among alcoholics: analyses of Canterbury Suicide Project data. *J. Stud. Alcohol* 64, 551–554. <https://doi.org/10.15288/jsa.2003.64.551>
- Cooper-Kazaz, R., 2013. Psychiatric consultation of all suicide-attempt patients during a one year period in a tertiary hospital. *Isr. Med. Assoc. J. IMAJ* 15, 424–429.
- Fernando, S.M., Qureshi, D., Sood, M.M., Pugliese, M., Talarico, R., Myran, D.T., Herridge, M.S., Needham, D.M., Rochweg, B., Cook, D.J., Wunsch, H., Fowler, R.A., Scales, D.C., Bienvenu, O.J., Rowan, K.M., Kisilewicz, M., Thompson, L.H., Tanuseputro, P., Kyeremanteng, K., 2021. Suicide and self-harm in adult survivors of critical illness: population based cohort study. *The BMJ* 373, n973. <https://doi.org/10.1136/bmj.n973>
- Giner, L., Jausent, I., Olié, E., Béziat, S., Guillaume, S., Baca-Garcia, E., Lopez-Castroman, J., Courtet, P., 2014. Violent and serious suicide attempters: one step closer to suicide? *J. Clin. Psychiatry* 75, e191-197. <https://doi.org/10.4088/JCP.13m08524>
- Goldman-Mellor, S., Olfson, M., Lidon-Moyano, C., Schoenbaum, M., 2019. Association of Suicide and Other Mortality With Emergency Department Presentation. *JAMA Netw. Open* 2, e1917571. <https://doi.org/10.1001/jamanetworkopen.2019.17571>
- Gvion, Y., Levi-Belz, Y., 2018. Serious Suicide Attempts: Systematic Review of Psychological Risk Factors. *Front. Psychiatry* 9. <https://doi.org/10.3389/fpsy.2018.00056>
- Han, B., Kott, P.S., Hughes, A., McKeon, R., Blanco, C., Compton, W.M., 2016. Estimating the rates of deaths by suicide among adults who attempt suicide in the United States. *J. Psychiatr. Res.* 77, 125–133. <https://doi.org/10.1016/j.jpsychires.2016.03.002>
- Hawton, K., 2001. Studying Survivors of Nearly Lethal Suicide Attempts: An Important Strategy in Suicide Research. *Suicide Life. Threat. Behav.* 32, 76–84. <https://doi.org/10.1521/suli.32.1.5.76.24215>
- Hawton, K., Appleby, L., Platt, S., Foster, T., Cooper, J., Malmberg, A., Simkin, S., 1998. The psychological autopsy approach to studying suicide: a review of methodological issues. *J. Affect. Disord.* 50, 269–276. [https://doi.org/10.1016/s0165-0327\(98\)00033-0](https://doi.org/10.1016/s0165-0327(98)00033-0)
- Hawton, K., Bergen, H., Cooper, J., Turnbull, P., Waters, K., Ness, J., Kapur, N., 2015. Suicide following self-harm: findings from the Multicentre Study of self-harm in England, 2000-2012. *J. Affect. Disord.* 175, 147–151. <https://doi.org/10.1016/j.jad.2014.12.062>
- Hawton, K., Fagg, J., 1988. Suicide, and other causes of death, following attempted suicide. *Br. J. Psychiatry J. Ment. Sci.* 152, 359–366. <https://doi.org/10.1192/bjp.152.3.359>
- Hawton, K., Sutton, L., Haw, C., Sinclair, J., Harriss, L., 2005. Suicide and attempted suicide in bipolar disorder: a systematic review of risk factors. *J. Clin. Psychiatry* 66, 693–704. <https://doi.org/10.4088/jcp.v66n0604>
- Hawton, K., van Heeringen, K., 2009. Suicide. *Lancet Lond. Engl.* 373, 1372–1381. [https://doi.org/10.1016/S0140-6736\(09\)60372-X](https://doi.org/10.1016/S0140-6736(09)60372-X)

- Horesh, N., Levi, Y., Apter, A., 2012. Medically serious versus non-serious suicide attempts: relationships of lethality and intent to clinical and interpersonal characteristics. *J. Affect. Disord.* 136, 286–293. <https://doi.org/10.1016/j.jad.2011.11.035>
- Kennedy, P., Garmon-Jones, L., 2017. Self-harm and suicide before and after spinal cord injury: a systematic review. *Spinal Cord* 55, 2–7. <https://doi.org/10.1038/sc.2016.135>
- Kessler, R.C., Berglund, P., Borges, G., Nock, M., Wang, P.S., 2005. Trends in suicide ideation, plans, gestures, and attempts in the United States, 1990-1992 to 2001-2003. *JAMA* 293, 2487–2495. <https://doi.org/10.1001/jama.293.20.2487>
- Kim, S., Choi, K.H., Lee, K.-S., Kim, D.-J., Hong, S.-C., Lee, H.-K., Kweon, Y.-S., Lee, C.T., Lee, K.-U., 2020. Risk Factors for Serious Suicide Attempts with High Medical Severity. *Suicide Life. Threat. Behav.* 50, 408–421. <https://doi.org/10.1111/sltb.12597>
- Kuo, W.H., Gallo, J.J., Tien, A.Y., 2001. Incidence of suicide ideation and attempts in adults: the 13-year follow-up of a community sample in Baltimore, Maryland. *Psychol. Med.* 31, 1181–1191. <https://doi.org/10.1017/s0033291701004482>
- Levi-Belz, Y., Beautrais, A., 2016. Serious suicide attempts: Toward an integration of terms and definitions. *Crisis J. Crisis Interv. Suicide Prev.* 37, 299–309. <https://doi.org/10.1027/0227-5910/a000386>
- Lopez-Castroman, J., Cerrato, L., Beziat, S., Jaussent, I., Guillaume, S., Courtet, P., 2016. Heavy tobacco dependence in suicide attempters making recurrent and medically serious attempts. *Drug Alcohol Depend.* 160, 177–182. <https://doi.org/10.1016/j.drugalcdep.2016.01.004>
- Lopez-Castroman, J., Jaussent, I., Beziat, S., Guillaume, S., Baca-Garcia, E., Olié, E., Courtet, P., 2015. Posttraumatic Stress Disorder following childhood abuse increases the severity of suicide attempts. *J. Affect. Disord.* 170, 7–14. <https://doi.org/10.1016/j.jad.2014.08.010>
- Mościcki, E.K., 1995. Epidemiology of suicidal behavior. *Suicide Life. Threat. Behav.* 25, 22–35.
- Mościcki, E.K., O'Carroll, P., Rae, D.S., Locke, B.Z., Roy, A., Regier, D.A., 1988. Suicide attempts in the Epidemiologic Catchment Area Study. *Yale J. Biol. Med.* 61, 259–268.
- Olfson, M., Marcus, S.C., Bridge, J.A., 2014. Focusing suicide prevention on periods of high risk. *JAMA* 311, 1107–1108. <https://doi.org/10.1001/jama.2014.501>
- Parra-Urbe, I., Blasco-Fontecilla, H., Garcia-Parés, G., Martínez-Naval, L., Valero-Coppin, O., Cebrià-Meca, A., Oquendo, M.A., Palao-Vidal, D., 2017. Risk of re-attempts and suicide death after a suicide attempt: A survival analysis. *BMC Psychiatry* 17, 163. <https://doi.org/10.1186/s12888-017-1317-z>
- Persett, P.S., Grimholt, T.K., Ekeberg, O., Jacobsen, D., Myhren, H., 2018. Patients admitted to hospital after suicide attempt with violent methods compared to patients with deliberate self-poisoning - a study of background variables, somatic and psychiatric health and suicidal behavior. *BMC Psychiatry* 18, 21. <https://doi.org/10.1186/s12888-018-1602-5>
- Powell, K.E., Kresnow, M.J., Mercy, J.A., Potter, L.B., Swann, A.C., Frankowski, R.F., Lee, R.K., Bayer, T.L., 2001. Alcohol consumption and nearly lethal suicide attempts. *Suicide Life. Threat. Behav.* 32, 30–41. <https://doi.org/10.1521/suli.32.1.5.30.24208>
- Probert-Lindström, S., Berge, J., Westrin, Å., Öjehagen, A., Skogman Pavulans, K., 2020. Long-term risk factors for suicide in suicide attempters examined at a medical emergency in patient unit: results from a 32-year follow-up study. *BMJ Open* 10, e038794. <https://doi.org/10.1136/bmjopen-2020-038794>
- Qin, P., Hawton, K., Mortensen, P.B., Webb, R., 2014. Combined effects of physical illness and comorbid psychiatric disorder on risk of suicide in a national population study. *Br. J. Psychiatry J. Ment. Sci.* 204, 430–435. <https://doi.org/10.1192/bjp.bp.113.128785>
- Schrijvers, D.L., Bollen, J., Sabbe, B.G.C., 2012. The gender paradox in suicidal behavior and its impact on the suicidal process. *J. Affect. Disord.* 138, 19–26. <https://doi.org/10.1016/j.jad.2011.03.050>
- Stein, B., Müller, M.M., Meyer, L.K., Söllner, W., CL Guidelines Working Group, 2020. Psychiatric and Psychosomatic Consultation-Liaison Services in General Hospitals: A Systematic Review and Meta-Analysis of Effects on Symptoms of Depression and Anxiety. *Psychother. Psychosom.* 89, 6–16. <https://doi.org/10.1159/000503177>

- Stenbacka, M., Jokinen, J., 2015. Violent and non-violent methods of attempted and completed suicide in Swedish young men: the role of early risk factors. *BMC Psychiatry* 15, 196. <https://doi.org/10.1186/s12888-015-0570-2>
- Sun, J., Guo, X., Zhang, J., Wang, M., Jia, C., Xu, A., 2015. Incidence and fatality of serious suicide attempts in a predominantly rural population in Shandong, China: a public health surveillance study. *BMJ Open* 5. <https://doi.org/10.1136/bmjopen-2014-006762>
- Tejedor, M.C., Díaz, A., Castellón, J.J., Pericay, J.M., 1999. Attempted suicide: repetition and survival—findings of a follow-up study. *Acta Psychiatr. Scand.* 100, 205–211. <https://doi.org/10.1111/j.1600-0447.1999.tb10847.x>
- Värnik, A., Kölves, K., van der Feltz-Cornelis, C.M., Marusic, A., Oskarsson, H., Palmer, A., Reisch, T., Scheerder, G., Arensman, E., Aromaa, E., Giupponi, G., Gusmäo, R., Maxwell, M., Pull, C., Szekely, A., Sola, V.P., Hegerl, U., 2008. Suicide methods in Europe: a gender-specific analysis of countries participating in the “European Alliance Against Depression.” *J. Epidemiol. Community Health* 62, 545–551. <https://doi.org/10.1136/jech.2007.065391>
- Vuagnat, A., Jollant, F., Abbar, M., Hawton, K., Quantin, C., 2019. Recurrence and mortality 1 year after hospital admission for non-fatal self-harm: a nationwide population-based study. *Epidemiol. Psychiatr. Sci.* 29, e20. <https://doi.org/10.1017/S2045796019000039>
- Wand, A.P.F., Wood, R., Macfarlane, M.D., Hunt, G.E., 2016. Comparison of consultation-liaison psychiatry services for inner-city, district or regional general hospitals using a common tool: Does one size fit all? *J. Psychosom. Res.* 84, 13–21. <https://doi.org/10.1016/j.jpsychores.2016.03.007>
- While, D., Bickley, H., Roscoe, A., Windfuhr, K., Rahman, S., Shaw, J., Appleby, L., Kapur, N., 2012. Implementation of mental health service recommendations in England and Wales and suicide rates, 1997-2006: a cross-sectional and before-and-after observational study. *Lancet Lond. Engl.* 379, 1005–1012. [https://doi.org/10.1016/S0140-6736\(11\)61712-1](https://doi.org/10.1016/S0140-6736(11)61712-1)
- Wood, R., Wand, A.P.F., 2014. The effectiveness of consultation-liaison psychiatry in the general hospital setting: a systematic review. *J. Psychosom. Res.* 76, 175–192. <https://doi.org/10.1016/j.jpsychores.2014.01.002>

TABLES

Table 1. Description of MSSA population by sex.

	<i>Male</i> <i>N=38006</i> <i>N (%)</i>	<i>Female</i> <i>N=43953</i> <i>N (%)</i>	<i>Total</i> <i>N=81959</i> <i>N (%)</i>	<i>p-value</i>
Age (mean±SD)	45.04±16.81	46.45±18.31	45.8±17.64	<.0001
Age in five classes				<.0001
[8 years; 15 years [89 (21.6%)	323 (78.4%)	412 (0.5%)	
[15 years; 25 years [4550 (40.9%)	6570 (59.1%)	11120 (13.6%)	
[25 years; 55 years [23258 (50.4%)	22892 (49.6%)	46150 (56.3%)	
[55 years; 75 years [7817 (42.1%)	10766 (57.9%)	18583 (22.7%)	
75+ years	2292 (40.3%)	3402 (59.7%)	5694 (6.9%)	
Drug poisoning (X60-X64)	21472 (40.4%)	31695 (59.6%)	53167 (64.9%)	<.0001
Death because of the attempt	3090 (68.4%)	1429 (31.6%)	4519 (5.5%)	<.0001
Admission				
From emergency room	25073 (46.2%)	29185 (53.8%)	54258 (66.2%)	0.31
From a psychiatric unit	134 (45%)	164 (55%)	298 (0.4%)	
From home	11731 (46.6%)	13443 (53.4%)	25174 (30.7%)	
Other	1048 (47.7%)	1147 (52.3%)	2195 (2.7%)	
Discharge				
To home	17638 (46.8%)	20014 (53.2%)	37652 (45.9%)	<.0001
To psychiatric unit	10456 (41.5%)	14721 (58.5%)	25177 (30.7%)	
Other	6822 (46.7%)	7789 (53.3%)	14611 (17.8%)	
Days in intensive care (mean±SD)	5.55±10.12	4.4±7.39	4.93±8.78	<.0001
Charlson Score				
0 (one-year mortality=12%)	33553 (45.7%)	39839 (54.3%)	73392 (89.5%)	<.0001
1-2 (one-year mortality=26%)	2897 (48.8%)	3041 (51.2%)	5938 (7.2%)	
3-4 (one-year mortality=52%)	1035 (61.4%)	652 (38.6%)	1687 (2.1%)	
5+ (one-year mortality=85%)	521 (55.3%)	421 (44.7%)	942 (1.1%)	
ICD 10 Codes for Mental and behavioural disorders				
F01-F09	1212 (50.5%)	1187 (49.5%)	2399 (4.1%)	<.0001
F10-F19	15423 (56.4%)	11923 (43.6%)	27346 (46.2%)	<.0001
F20-F29	2698 (59.8%)	1814 (40.2%)	4512 (7.6%)	<.0001
F30-F39	12548 (38.2%)	20339 (61.8%)	32887 (55.6%)	<.0001
F40-F48	2989 (37.7%)	4942 (62.3%)	7931 (13.4%)	<.0001
F50-F59	101 (15%)	573 (85%)	674 (1.1%)	<.0001
F60-F69	908 (44.4%)	1139 (55.6%)	2047 (3.5%)	0.07
F70-F79	79 (60.8%)	51 (39.2%)	130 (0.2%)	.0009
F80-F89	37 (62.7%)	22 (37.3%)	59 (0.1%)	0.01
F90-F98	344 (40.9%)	497 (59.1%)	841 (1.4%)	.002

	<i>Male</i> N=38006 N (%)	<i>Female</i> N=43953 N (%)	<i>Total</i> N=81959 N (%)	<i>p-value</i>
F99	293 (41%)	422 (59%)	715 (1.2%)	.004
Alcohol use disorder	11733 (57.7%)	8593 (42.3%)	20326 (34.3%)	<.0001
Bipolar disorder	1407 (33.1%)	2843 (66.9%)	4250 (7.2%)	<.0001
Schizophrenia	1504 (68.8%)	681 (31.2%)	2185 (3.7%)	<.0001
Psychiatric admission in the 3 days following discharge	12309 (42.1%)	16912 (57.9%)	29221 (35.7%)	<.0001

Table 2. Description by living vs. deceased status

	<i>Alive following first MSSA N=77440 N (%)</i>	<i>Died following first MSSA N=4519 N (%)</i>	<i>Total N=81959 N (%)</i>	<i>p-value</i>
Male	34916 (91.9%)	3090 (8.1%)	38006 (46.4%)	
Female	42524 (96.7%)	1429 (3.3%)	43953 (53.6%)	
Age (mean±SD)	45.28±17.41	54.69±19.23	45.8±17.64	<.0001
Age in five classes				
[8 years; 15 years [398 (96.6%)	14 (3.4%)	412 (0.5%)	<.0001
[15 years; 25 years [10848 (97.6%)	272 (2.4%)	11120 (13.6%)	
[25 years; 55 years [44111 (95.6%)	2039 (4.4%)	46150 (56.3%)	
[55 years; 75 years [17242 (92.8%)	1341 (7.2%)	18583 (22.7%)	
75+ years	4841 (85%)	853 (15%)	5694 (6.9%)	
Admission				
From emergency room	51636 (95.2%)	2622 (4.8%)	54258 (66.2%)	<.0001
From a psychiatric unit	264 (88.6%)	34 (11.4%)	298 (0.4%)	
From home	23456 (93.2%)	1718 (6.8%)	25174 (30.7%)	
Other	2053 (93.5%)	142 (6.5%)	2195 (2.7%)	
Discharge				<.0001
To home	37652 (100%)	0	37652 (45.9%)	
To psychiatric unit	25177 (100%)	0	25177 (30.7%)	
Other	14611 (100%)	0	14611 (17.8%)	
MSSA mechanism				
Drug poisoning (X60-64)	52155 (98.1%)	1012 (1.9%)	53167 (64.9%)	<.0001
Other (X65-X84)	25285 (87.8%)	3507 (12.2%)	28792 (35.1%)	
Days in intensive care (mean±SD)	7.13±13.23	8.57±20.31	7.21±13.72	<.0001
Charlson Score				
0 (one-year mortality=12%)	69592 (94.8%)	3800 (5.2%)	73392 (89.5%)	<.0001
1-2 (one-year mortality=26%)	5513 (92.8%)	425 (7.2%)	5938 (7.2%)	
3-4 (one-year mortality=52%)	1509 (89.4%)	178 (10.6%)	1687 (2.1%)	
5+ (one-year mortality=85%)	826 (87.7%)	116 (12.3%)	942 (1.1%)	
Alcohol SUD	19754 (97.2%)	572 (2.8%)	20326 (34.3%)	<0.001
Bipolar disorder	4118 (96.9%)	132 (3.1%)	4250 (7.2%)	0.13
Schizophrenia	2111 (96.6%)	74 (3.4%)	2185 (3.7%)	0.74
ICD 10 Codes for Mental and behavioural disorders				
F01-F09	2291 (95.5%)	108 (4.5%)	2399 (4.1%)	.007
F10-F19	26593 (97.2%)	753 (2.8%)	27346 (46.2%)	<.0001
F20-F29	4368 (96.8%)	144 (3.2%)	4512 (7.6%)	0.22

	<i>Alive following first MSSA N=77440 N (%)</i>	<i>Died following first MSSA N=4519 N (%)</i>	<i>Total N=81959 N (%)</i>	<i>p-value</i>
F30-F39	31624 (96.2%)	1263 (3.8%)	32887 (55.6%)	<.009
F40-F48	7773 (98%)	158 (2%)	7931 (13.4%)	<.0001
F50-F59	657 (97.5%)	17 (2.5%)	674 (1.1%)	0.19
F60-F69	2025 (98.9%)	22 (1.1%)	2047 (3.5%)	<.0001
F70-F79	126 (96.9%)	4 (3.1%)	130 (0.2%)	1
F80-F89	58 (98.3%)	1 (1.7%)	59 (0.1%)	–
F90-F98	822 (97.7%)	19 (2.3%)	841 (1.4%)	.047
F99	697 (97.5%)	18 (2.5%)	715 (1.2%)	0.15
Psychiatric admission in the 3 days following discharge	29221 (100%)	0	29221 (35.7%)	

Table 3. Description of patients discharged to a psychiatric service following the first MSSA (77 440 patients alive following the MSSA)

	<i>Discharge to a psychiatric unit: 0 N=52263 N (%)</i>	<i>Discharge to a psychiatric unit: 1 N=25177 N (%)</i>	<i>Total N=77440 N (%)</i>	<i>p-value</i>
Male	24460 (70.1%)	10456 (29.9%)	34916 (45.1%)	<.0001
Female	27803 (65.4%)	14721 (34.6%)	42524 (54.9%)	
Age (mean±SD)	45.13± 17.93	45.58±16.27	45.28±17.41	.0006
Age in five classes				
[8 years; 15 years [317 (79.6%)	81 (20.4%)	398 (0.5%)	<.0001
[15 years; 25 years [7735 (71.3%)	3113 (28.7%)	10848 (14%)	
[25 years; 55 years [29390 (66.6%)	14721 (33.4%)	44111 (57%)	
[55 years ;75 years [11099 (64.4%)	6143 (35.6%)	17242 (22.3%)	
75+ years	3722 (76.9%)	1119 (23.1%)	4841 (6.3%)	
Admission				
From emergency room	34111 (66.1%)	17525 (33.9%)	51636 (66.7%)	<.0001
From a psychiatric unit	59 (22.3%)	205 (77.7%)	264 (0.3%)	
From home	16528 (70.5%)	6928 (29.5%)	23456 (30.3%)	
Other	1541 (75.1%)	512 (24.9%)	2053 (2.7%)	
Discharge				
To home	37652 (100%)	0	37652 (48.6%)	<.0001
To psychiatric unit	0	25177 (100%)	25177 (32.5%)	
Other	14611 (100%)	0	14611 (18.9%)	
MSSA mechanism				
Drug poisoning (X60-64)	34296 (65.8%)	17859 (34.2%)	52155 (67.3%)	<.0001
Other (X65-X84)	17967 (71.1%)	7318 (28.9%)	25285 (32.7%)	
Days in intensive care (mean±SD)	8.34±15.17	4.61±7.2	7.13±13.23	<.0001
Charlson Score				
0 (one-year mortality=12%)	46460 (66.8%)	23132 (33.2%)	69592 (89.9%)	<.0001
1-2 (one-year mortality=26%)	3955 (71.7%)	1558 (28.3%)	5513 (7.1%)	
3-4 (one-year mortality=52%)	1168 (77.4%)	341 (22.6%)	1509 (1.9%)	
5+ (one-year mortality=85%)	680 (82.3%)	146 (17.7%)	826 (1.1%)	
Alcohol SUD	14142 (71.6%)	5612 (28.4%)	19754 (34.6%)	<.0001
Bipolar disorder	2130 (51.7%)	1988 (48.3%)	4118 (7.2%)	<.0001
Schizophrenia	1120 (53.1%)	991 (46.9%)	2111 (3.7%)	<.0001
ICD 10 Codes for Mental and behavioural disorders				
F01-F09	1735 (75.7%)	556 (24.3%)	2291 (4%)	<.0001
F10-F19	18954 (71.3%)	7639 (28.7%)	26593 (46.6%)	<.0001

	<i>Discharge to a psychiatric unit: 0 N=52263 N (%)</i>	<i>Discharge to a psychiatric unit: 1 N=25177 N (%)</i>	<i>Total N=77440 N (%)</i>	<i>p-value</i>
F20-F29	2399 (54.9%)	1969 (45.1%)	4368 (7.7%)	<.0001
F30-F39	19781 (62.6%)	11843 (37.4%)	31624 (55.4%)	<.0001
F40-F48	5456 (70.2%)	2317 (29.8%)	7773 (13.6%)	<.0001
F50-F59	441 (67.1%)	216 (32.9%)	657 (1.2%)	<.0001
F60-F69	1313 (64.8%)	712 (35.2%)	2025 (3.5%)	0.10
F70-F79	91 (72.2%)	35 (27.8%)	126 (0.2%)	0.18
F80-F89	36 (62.1%)	22 (37.9%)	58 (0.1%)	0.47
F90-F98	572 (69.6%)	250 (30.4%)	822 (1.4%)	0.06
F99	440 (63.1%)	257 (36.9%)	697 (1.2%)	0.06
Psychiatric admission in the 3 days following discharge	8093 (27.7%)	21128 (72.3%)	29221 (37.7%)	<.0001

Table 4. Description of MSSA patients according to the suicide attempt method

	<i>Intentional self-poisoning with drugs (X60-X64)</i> N=53167 N (%)	<i>Intentional self-poisoning with other substances (alcohol, solvents, pesticides, chemicals) (X65-X69)</i> N=7673 N (%)	<i>Intentional self-harm by hanging, strangulation and suffocation (X70)</i> N=3857 N (%)	<i>Intentional self-harm by drowning and submersion (X71)</i> N=256 N (%)	<i>Intentional self-harm by firearm (X72 à X74)</i> N=1297 N (%)	<i>Intentional self-harm by smoke, fire and flames (X75-X77)</i> N=609 N (%)
Male	21472 (56.5%)	4359 (11.5%)	3010 (7.9%)	102 (0.3%)	1196 (3.1%)	399 (1%)
Female	31695 (72.1%)	3314 (7.5%)	847 (1.9%)	154 (0.4%)	101 (0.2%)	210 (0.5%)
Age in five classes						
[8 years; 15 years [265 (64.3%)	30 (7.3%)	23 (5.6%)	1 (0.2%)	1 (0.2%)	2 (0.5%)
[15 years; 25 years [7446 (67%)	875 (7.9%)	375 (3.4%)	12 (0.1%)	64 (0.6%)	56 (0.5%)
[25 years; 55 years [29926 (64.8%)	4471 (9.7%)	2452 (5.3%)	63 (0.1%)	490 (1.1%)	381 (0.8%)
[55 years ;75 years [11905 (64.1%)	1900 (10.2%)	769 (4.1%)	129 (0.7%)	462 (2.5%)	154 (0.8%)
75+ years	3625 (63.7%)	397 (7%)	238 (4.2%)	51 (0.9%)	280 (4.9%)	16 (0.3%)
Admission						
From emergency room	36254 (66.8%)	5000 (9.2%)	2191 (4%)	161 (0.3%)	801 (1.5%)	217 (0.4%)
From a psychiatric unit	165 (55.4%)	20 (6.7%)	52 (17.4%)	2 (0.7%)	2 (0.7%)	4 (1.3%)
From home	15425 (61.3%)	2432 (9.7%)	1527 (6.1%)	89 (0.4%)	445 (1.8%)	337 (1.3%)
Other	1306 (59.5%)	215 (9.8%)	86 (3.9%)	3 (0.1%)	48 (2.2%)	51 (2.3%)
Missing	17	6	1	1	1	0
Discharge						
To home	26324 (69.9%)	4293 (11.4%)	674 (1.8%)	65 (0.2%)	324 (0.9%)	104 (0.3%)
To psychiatric unit	17859 (70.9%)	1903 (7.6%)	925 (3.7%)	82 (0.3%)	167 (0.7%)	54 (0.2%)
Other	7972 (54.6%)	1183 (8.1%)	516 (3.5%)	44 (0.3%)	362 (2.5%)	314 (2.1%)

	<i>Intentional self-poisoning with drugs (X60-X64)</i> N=53167 N (%)	<i>Intentional self-poisoning with other substances (alcohol, solvents, pesticides, chemicals) (X65-X69)</i> N=7673 N (%)	<i>Intentional self-harm by hanging, strangulation and suffocation (X70)</i> N=3857 N (%)	<i>Intentional self-harm by drowning and submersion (X71)</i> N=256 N (%)	<i>Intentional self-harm by firearm (X72 à X74)</i> N=1297 N (%)	<i>Intentional self-harm by smoke, fire and flames (X75-X77)</i> N=609 N (%)
Death because of the attempt	1012 (22.4%)	294 (6.5%)	1742 (38.5%)	65 (1.4%)	444 (9.8%)	137 (3%)
Charlson Score						
0 (one-year mortality=12%)	47569 (64.8%)	6788 (9.2%)	3589 (4.9%)	226 (0.3%)	1102 (1.5%)	542 (0.7%)
1-2 (one-year mortality=26%)	3943 (66.4%)	584 (9.8%)	183 (3.1%)	21 (0.4%)	118 (2%)	51 (0.9%)
3-4 (one-year mortality=52%)	1061 (62.9%)	185 (11%)	59 (3.5%)	5 (0.3%)	45 (2.7%)	13 (0.8%)
5+ (one-year mortality=85%)	594 (63.1%)	116 (12.3%)	26 (2.8%)	4 (0.4%)	32 (3.4%)	3 (0.3%)
ICD 10 Codes for Mental and behavioural disorders						
F01-F09	1178 (49.1%)	220 (9.2%)	113 (4.7%)	17 (0.7%)	80 (3.3%)	13 (0.5%)
F10-F19	17690 (64.7%)	3678 (13.4%)	853 (3.1%)	42 (0.2%)	292 (1.1%)	139 (0.5%)
F20-F29	2600 (57.6%)	247 (5.5%)	115 (2.5%)	15 (0.3%)	34 (0.8%)	101 (2.2%)
F30-F39	22376 (68%)	2419 (7.4%)	1429 (4.3%)	121 (0.4%)	503 (1.5%)	209 (0.6%)
F40-F48	5539 (69.8%)	578 (7.3%)	261 (3.3%)	30 (0.4%)	88 (1.1%)	63 (0.8%)
F50-F59	486 (72.1%)	38 (5.6%)	17 (2.5%)	0	4 (0.6%)	3 (0.4%)
F60-F69	1292 (63.1%)	101 (4.9%)	86 (4.2%)	6 (0.3%)	22 (1.1%)	28 (1.4%)
F70-F79	78 (60%)	11 (8.5%)	3 (2.3%)	1 (0.8%)	1 (0.8%)	3 (2.3%)
F80-F89	31 (52.5%)	3 (5.1%)	3 (5.1%)	1 (1.7%)	0	2 (3.4%)

	<i>Intentional self-poisoning with drugs (X60-X64)</i> N=53167 N (%)	<i>Intentional self-poisoning with other substances (alcohol, solvents, pesticides, chemicals) (X65-X69)</i> N=7673 N (%)	<i>Intentional self-harm by hanging, strangulation and suffocation (X70)</i> N=3857 N (%)	<i>Intentional self-harm by drowning and submersion (X71)</i> N=256 N (%)	<i>Intentional self-harm by firearm (X72 à X74)</i> N=1297 N (%)	<i>Intentional self-harm by smoke, fire and flames (X75-X77)</i> N=609 N (%)
F90-F98	516 (61.4%)	78 (9.3%)	30 (3.6%)	3 (0.4%)	17 (2%)	3 (0.4%)
F99	128 (17.9%)	14 (2%)	6 (0.8%)	1 (0.1%)	3 (0.4%)	2 (0.3%)
Alcohol use disorder	12595 (62%)	3259 (16%)	646 (3.2%)	39 (0.2%)	238 (1.2%)	101 (0.5%)
Bipolar disorder	3054 (71.9%)	281 (6.6%)	86 (2%)	11 (0.3%)	35 (0.8%)	35 (0.8%)
Schizophrenia	1293 (59.2%)	103 (4.7%)	50 (2.3%)	5 (0.2%)	19 (0.9%)	42 (1.9%)
Missing	14487	1998	1596	73	503	201

Table 4. Description of MSSA patients according to the suicide attempt method (Suite)

	<i>Intentional self-harm by sharp or blunt object (X78-X79)</i> N=2308 N (%)	<i>Intentional self-harm by jumping from a high place (X80)</i> N=2386 N (%)	<i>Intentional self-harm by collision (X81-X82)</i> N=316 N (%)	<i>Intentional self-harm by other specified or unspecified means (X83- X84)</i> N=10090 N (%)	<i>Total N=81959</i>
Male	1462 (3.8%)	1214 (3.2%)	209 (0.5%)	4583 (12.1%)	38006 (46.4%)
Female	846 (1.9%)	1172 (2.7%)	107 (0.2%)	5507 (12.5%)	43953 (53.6%)
Age in five classes					
[8 years; 15 years [16 (3.9%)	29 (7%)	0	45 (10.9%)	412 (0.5%)
[15 years; 25 years [329 (3%)	508 (4.6%)	68 (0.6%)	1387 (12.5%)	11120 (13.6%)
[25 years; 55 years [1171 (2.5%)	1287 (2.8%)	181 (0.4%)	5728 (12.4%)	46150 (56.3%)
[55 years ;75 years [554 (3%)	390 (2.1%)	55 (0.3%)	2265 (12.2%)	18583 (22.7%)
75+ years	238 (4.2%)	172 (3%)	12 (0.2%)	665 (11.7%)	5694 (6.9%)
Admission					
From emergency room	1551 (2.9%)	1392 (2.6%)	203 (0.4%)	6488 (12%)	54258 (66.2%)
From a psychiatric unit	15 (5%)	6 (2%)	0	32 (10.7%)	298 (0.4%)
From home*	696 (2.8%)	873 (3.5%)	100 (0.4%)	3250 (12.9%)	25174 (30.7%)
Other	44 (2%)	111 (5.1%)	13 (0.6%)	318 (14.5%)	2195 (2.7%)
Missing	2	4	0	2	34
Discharge					
To home*	892 (2.4%)	576 (1.5%)	98 (0.3%)	4302 (11.4%)	37652 (45.9%)
To psychiatric unit	826 (3.3%)	298 (1.2%)	60 (0.2%)	3003 (11.9%)	25177 (30.7%)
Other	510 (3.5%)	1300 (8.9%)	142 (1%)	2268 (15.5%)	14611 (17.8%)


	<i>Intentional self-harm by sharp or blunt object (X78-X79)</i> N=2308 N (%)	<i>Intentional self-harm by jumping from a high place (X80)</i> N=2386 N (%)	<i>Intentional self-harm by collision (X81-X82)</i> N=316 N (%)	<i>Intentional self-harm by other specified or unspecified means (X83- X84)</i> N=10090 N (%)	<i>Total N=81959</i>
Death because of the attempt	80 (1.8%)	212 (4.7%)	16 (0.4%)	517 (11.4%)	4519 (5.5%)
Charlson Score					
0 (one-year mortality=12%)	2000 (2.7%)	2197 (3%)	299 (0.4%)	9080 (12.4%)	73392 (89.5%)
1-2 (one-year mortality=26%)	186 (3.1%)	132 (2.2%)	10 (0.2%)	710 (12%)	5938 (7.2%)
3-4 (one-year mortality=52%)	79 (4.7%)	41 (2.4%)	6 (0.4%)	193 (11.4%)	1687 (2.1%)
5+ (one-year mortality=85%)	43 (4.6%)	16 (1.7%)	1 (0.1%)	107 (11.4%)	942 (1.1%)
ICD 10 Codes for Mental and behavioural disorders					
F01-F09	78 (3.3%)	170 (7.1%)	13 (0.5%)	517 (21.6%)	2399 (4.1%)
F10-F19	698 (2.6%)	725 (2.7%)	97 (0.4%)	3132 (11.5%)	27346 (46.2%)
F20-F29	247 (5.5%)	454 (10.1%)	36 (0.8%)	663 (14.7%)	4512 (7.6%)
F30-F39	1006 (3.1%)	814 (2.5%)	124 (0.4%)	3886 (11.8%)	32887 (55.6%)
F40-F48	241 (3%)	198 (2.5%)	30 (0.4%)	903 (11.4%)	7931 (13.4%)
F50-F59	20 (3%)	18 (2.7%)	0	88 (13.1%)	674 (1.1%)
F60-F69	114 (5.6%)	65 (3.2%)	12 (0.6%)	321 (15.7%)	2047 (3.5%)
F70-F79	5 (3.8%)	10 (7.7%)	1 (0.8%)	17 (13.1%)	130 (0.2%)
F80-F89	2 (3.4%)	6 (10.2%)	0	11 (18.6%)	59 (0.1%)

	<i>Intentional self-harm by sharp or blunt object (X78-X79)</i> N=2308 N (%)	<i>Intentional self-harm by jumping from a high place (X80)</i> N=2386 N (%)	<i>Intentional self-harm by collision (X81-X82)</i> N=316 N (%)	<i>Intentional self-harm by other specified or unspecified means (X83- X84)</i> N=10090 N (%)	<i>Total N=81959</i>
F90-F98	29 (3.4%)	30 (3.6%)	4 (0.5%)	131 (15.6%)	841 (1.4%)
F99	8 (1.1%)	17 (2.4%)	2 (0.3%)	534 (74.7%)	715 (1.2%)
Alcohol use disorder	492 (2.4%)	516 (2.5%)	70 (0.3%)	2370 (11.7%)	20326 (34.3%)
Bipolar disorder	132 (3.1%)	134 (3.2%)	14 (0.3%)	468 (11%)	4250 (7.2%)
Schizophrenia	114 (5.2%)	248 (11.4%)	20 (0.9%)	291 (13.3%)	2185 (3.7%)
Missing	521	537	88	2779	22783

Table 5. Incidence of the first MSSA per 100 000 person-years by age group and sex

	2012			2013			2014			2015			2016			2017			2018			2019		
	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both	Male	Female	Both
Age in five classes																								
[8 years; 15 years [0.31	1.63	0.95	0.41	1.19	0.79	0.44	1.68	1.05	0.41	1.24	0.82	0.33	1.47	0.89	0.40	1.15	0.77	0.43	1.57	0.99	0.27	1.50	0.87
[15 years; 25 years [13.07	17.30	15.16	12.14	18.55	15.29	15.74	20.58	18.12	13.28	19.49	16.33	13.48	19.00	16.19	13.54	21.73	17.55	15.74	25.19	20.37	16.51	28.13	22.21
[25 years; 55 years [22.29	22.55	22.42	22.08	21.57	21.82	24.41	24.65	24.53	22.93	21.71	22.31	22.23	21.06	21.63	23.31	21.69	22.49	24.27	22.06	23.15	24.34	21.79	23.05
[55 years;75 years [12.42	17.51	15.08	12.87	16.65	14.85	13.85	18.03	16.04	14.12	17.66	15.97	12.81	17.03	15.02	15.11	18.17	16.72	14.54	17.24	15.96	15.10	16.85	16.02
75+ years	12.05	11.08	11.44	11.65	9.29	10.17	12.68	11.35	11.85	12.29	11.47	11.78	12.77	11.91	12.24	12.29	11.24	11.65	13.05	12.04	12.43	11.25	11.25	11.25
All ages	15.66	17.42	16.57	15.48	16.68	16.10	17.30	18.84	18.10	16.31	17.33	16.84	15.68	16.90	16.31	16.67	17.67	17.19	17.25	18.11	17.70	17.33	18.14	17.75

Figure 1 : Data flowchart :



APPENDIX

ICD 10 Codes for Mental and behavioural disorders

F01-F09: Organic, including symptomatic, mental disorders

F10-F19: Mental and behavioural disorders due to psychoactive substance use

F20-F29: Schizophrenia, schizotypal and delusional disorders

F30-F39: Mood [affective] disorders

F40-F48: Neurotic, stress-related and somatoform disorders

F50-F59: Behavioural syndromes associated with physiological disturbances and physical factors

F60-F69: Disorders of adult personality and behaviour

F70-F79: Mental retardation

F80-F89: Disorders of psychological development

F90-F98: Behavioural and emotional disorders with onset usually occurring in childhood and adolescence

F99: Unspecified mental disorder

SERMENT

- *En présence des Maîtres de cette école, de mes chers condisciples et devant l'effigie d'Hippocrate, je promets et je jure, au nom de l'Être suprême, d'être fidèle aux lois de l'honneur et de la probité dans l'exercice de la médecine.*

- *Je donnerai mes soins gratuits à l'indigent et n'exigerai jamais un salaire au-dessus de mon travail.*

- *Admis dans l'intérieur des maisons, mes yeux ne verront pas ce qui s'y passe, ma langue taira les secrets qui me seront confiés, et mon état ne servira pas à corrompre les mœurs, ni à favoriser le crime.*

- *Respectueux et reconnaissant envers mes Maîtres, je rendrai à leurs enfants l'instruction que j'ai reçue de leurs pères.*

- *Que les hommes m'accordent leur estime si je suis fidèle à mes promesses. Que je sois couvert d'opprobre et méprisé de mes confrères si j'y manque.*

RESUME

Contexte : Les personnes faisant des tentatives de suicide médicalement graves (medically serious suicide attempt (MSSA)) partagent un certain nombre de caractéristiques avec celles qui se suicident et présentent un risque élevé de suicide ultérieur. Les études menées jusqu'à présent ont surtout porté sur des échantillons cliniques ; une étude épidémiologique au niveau national permettra d'établir des profils à risque et des parcours de soins dans cette population.

Méthodes : Nous avons exploré la base de données nationale française des sorties d'hôpital (Programme de Médicalisation des Systèmes d'Information, PMSI) pour identifier toute MSSA ayant eu lieu entre 2012 et 2019. Des informations démographiques et médicales pertinentes ont été collectées sur la première MSSA de chaque patient avec une période de *washout* de deux ans.

Résultats : Le nombre de premières MSSAs s'est élevé à 81 959 cas sur 8 ans, avec un âge moyen de 45,8 ans et 53,6% de femmes. L'incidence était plus élevée chez les femmes (18.1 contre 17.3 pour 100 000 personnes-années). La méthode de suicide la plus courante était l'auto-intoxication médicamenteuse (64,9% des cas). En comparaison, les méthodes violentes étaient associées à une mortalité et à des comorbidités plus élevées et étaient plus fréquentes chez les hommes. Les troubles mentaux les plus fréquents étaient les troubles de l'humeur (55,6%) et les troubles de l'usage de substances (46,2%). Une minorité de survivants de MSSAs ont été hospitalisés en psychiatrie (32,5%), principalement des femmes.

Conclusions : Les MSSAs sont fréquentes et faciles à identifier. Il est nécessaire de renforcer la continuité des soins psychiatriques pour cette population étant donné le risque élevé de suicide ultérieur, et les faibles taux d'hospitalisation en psychiatrie après une MSSA. Une prise en charge spécifique ciblant cette population pourrait réduire les écarts de traitement.

MOTS-CLES Tentative de suicide grave, Suicide, Comportement suicidaire, Troubles affectifs, Epidémiologie, Psychiatrie, Programme de Médicalisation des Systèmes d'Information