



Contribution du score Mercuri analysé sur IRM musculaire, dans l'évaluation du handicap fonctionnel, chez des adultes présentant une arthrogrypose multiple congénitale

Marion Finzi

► To cite this version:

Marion Finzi. Contribution du score Mercuri analysé sur IRM musculaire, dans l'évaluation du handicap fonctionnel, chez des adultes présentant une arthrogrypose multiple congénitale. Human health and pathology. 2021. dumas-03736501

HAL Id: dumas-03736501

<https://dumas.ccsd.cnrs.fr/dumas-03736501v1>

Submitted on 20 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



AVERTISSEMENT

Ce document est le fruit d'un long travail approuvé par le jury de soutenance.

La propriété intellectuelle du document reste entièrement celle du ou des auteurs. Les utilisateurs doivent respecter le droit d'auteur selon la législation en vigueur, et sont soumis aux règles habituelles du bon usage, comme pour les publications sur papier : respect des travaux originaux, citation, interdiction du pillage intellectuel, etc.

Il est mis à disposition de toute personne intéressée par l'intermédiaire de [l'archive ouverte DUMAS](#) (Dépôt Universitaire de Mémoires Après Soutenance).

Si vous désirez contacter son ou ses auteurs, nous vous invitons à consulter la page de DUMAS présentant le document. Si l'auteur l'a autorisé, son adresse mail apparaîtra lorsque vous cliquerez sur le bouton « Détails » (à droite du nom).

Dans le cas contraire, vous pouvez consulter en ligne les annuaires de l'ordre des médecins, des pharmaciens et des sages-femmes.

Contact à la Bibliothèque universitaire de Médecine Pharmacie de Grenoble :

bump-theses@univ-grenoble-alpes.fr

Année : 2020-2021

**THE CONTRIBUTION OF MUSCULAR MRI USING MERCURI SCORE
TO EVALUATE DISABILITY PATTERNS
IN ADULTS WITH ARTHROGYPOSIS MULTIPLEX CONGENITA.**

THÈSE
POUR LE DIPLÔME D'ÉTAT DE DOCTEUR EN MÉDECINE

SPÉCIALITÉ :
MÉDECINE PHYSIQUE ET READAPTATION

Par Mme Marion FINZI

[Données à caractère personnel]

THÈSE SOUTENUE PUBLIQUEMENT À LA FACULTÉ DE MÉDECINE DE GRENOBLE

Le 20/09/2021.

DEVANT LE JURY COMPOSÉ DE :

Président du jury :

M. le Pr. Dominic PERENNOD (co-directeur de thèse)

Membres :

M. le Pr. Pierre-Simon JOUK

M. le Dr. Klaus DIETERICH (co-directeur de thèse)

Mme la Dr. Véronique BOURG

Mme la Dr. Chantal DURAND

L'UFR de Médecine de Grenoble n'entend donner aucune approbation ni improbation aux opinions émises dans les thèses ; ces opinions sont considérées comme propres à leurs auteurs.

Doyen de la Faculté : Pr. Patrice MORAND

Année 2020-2021

ENSEIGNANTS DE L'UFR DE MEDECINE

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH	ALBALADEJO Pierre	Anesthésiologie-réanimation et médecine péri-opératoire
PU-PH	APTEL Florent	Ophthalmologie
PU-PH	ARVIEUX-BARTHELEMY Catherine	Chirurgie viscérale et digestive
PU-PH	BAILLET Athan	Rhumatologie
PU-PH	BARONE-ROCHETTE Gilles	Cardiologie
PU-PH	BAYAT Sam	Physiologie
MCF Ass.MG	BENDAMENE Farouk	Médecine Générale
PU-PH	BENHAMOU Pierre-Yves	Endocrinologie, diabète et maladies métaboliques
PU-PH	BERGER François	Biologie cellulaire
MCU-PH	BIDART-COUTTON Marie	Biologie cellulaire
PU-PH	BLAISE Sophie	Chirurgie vasculaire ; médecine vasculaire
PR Ass. Méd.	BOILLOT Bernard	
MCU-PH	BOISSET Sandrine	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	BONAZ Bruno	Gastroentérologie ; hépatologie ; addictologie
PU-PH	BONNETERRE Vincent	Médecine et santé au travail
PU-PH	BOREL Anne-Laure	Nutrition
PU-PH	BOSSON Jean-Luc	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	BOTTARI Serge	Biologie cellulaire
PR Ass.MG	BOUCHAUD Jacques	Médecine Générale
PU-PH	BOUGEROL Thierry	Psychiatrie d'adultes
PU-PH	BOUILLET Laurence	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
MCU-PH	BOUSSAT Bastien	Epidémiologie, économie de la santé et prévention
PU-PH	BOUZAT Pierre	Anesthésiologie-réanimation et médecine péri-opératoire
PU-PH émérite	BRAMBILLA Christian	Pneumologie
PU-PH émérite	BRAMBILLA Elisabeth	Anatomie et cytologie pathologiques
MCU-PH	BRENIER-PINCHART Marie Pierre	Parasitologie et mycologie
PU-PH	BRICAULT Ivan	Radiologie et imagerie médicale
PU-PH	BRICHON Pierre-Yves	Chirurgie thoracique et cardiovasculaire
MCU-PH	BRIOT Raphaël	Thérapeutique-médecine de la douleur ; Addictologie
PU-PH émérite	CAHN Jean-Yves	Hématologie
PU-PH émérite	CARPENTIER Patrick	Chirurgie vasculaire, médecine vasculaire
PR Ass.MG	CARRILLO Yannick	Médecine Générale
PU-PH	CESBRON Jean-Yves	Immunologie
PU-PH	CHABARDES Stephan	Neurochirurgie
PU-PH	CHABRE Olivier	Endocrinologie, diabète et maladies métaboliques
PU-PH	CHAFFANJON Philippe	Anatomie
MCF Ass.MG	CHAMBORDON Benoît	Médecine Générale
PU-PH	CHARLES Julie	Dermato-vénéréologie
MCF Ass.MG	CHAUVET Marion	Médecine Générale
PU-PH	CHAVANON Olivier	Chirurgie thoracique et cardiovasculaire

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH	CHIQUET Christophe	Ophthalmologie
PU-PH	CHIRICA Mircea	Chirurgie viscérale et digestive
PU-PH	CINQUIN Philippe	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	CLAVARINO Giovanna	Immunologie
PU-PH	COHEN Olivier	Histologie, embryologie et cytogénétique
PU-PH	COURVOISIER Aurélien	Chirurgie infantile
PU-PH	COUTTON Charles	Génétique
PU-PH	COUTURIER Pascal	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PU-PH	CRACOWSKI Jean-Luc	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PU-PH	DEBATY Guillaume	Médecine d'Urgence
PU-PH	DEBILLON Thierry	Pédiatrie
PU-PH	DECAENS Thomas	Gastroentérologie ; hépatologie ; addictologie
PR Ass. Méd.	DEFAYE Pascal	Cardiologie
PU-PH	DEGANO Bruno	Pneumologie ; addictologie
PU-PH	DEMATTÉIS Maurice	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PU-PH émérite	DEMONGEOT Jacques	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	DERANSART Colin	Physiologie
PU-PH	DESCOTES Jean-Luc	Urologie
PU-PH	DETANTE Olivier	Neurologie
MCU-PH	DIETERICH Klaus	Génétique
MCU-PH	DOUTRELEAU Stéphane	Physiologie
MCU-PH	DUMESTRE-PERARD Chantal	Immunologie
PU-PH	EPAULARD Olivier	Maladies infectieuses ; Maladies tropicales
PU-PH	ESTEVE François	Biophysique et médecine nucléaire
MCU-PH	EYSERIC Hélène	Médecine légale et droit de la santé
PU-PH	FAUCHERON Jean-Luc	Chirurgie viscérale et digestive
MCU-PH	FAURE Julien	Biochimie et biologie moléculaire
PU-PH	FERRETTI Gilbert	Radiologie et imagerie médicale
PU-PH	FEUERSTEIN Claude	Physiologie
PU-PH	FONTAINE Éric	Nutrition
PU-PH	FRANCOIS Patrice	Epidémiologie, économie de la santé et prévention
MCU-MG	GABOREAU Yoann	Médecine Générale
PU-PH	GARBAN Frédéric	Hématologie ; Transfusion
PU-PH	GAUDIN Philippe	Rhumatologie
PU-PH	GAVAZZI Gaétan	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PU-PH	GAY Emmanuel	Neurochirurgie
MCU-PH	GILLOIS Pierre	Biostatistiques, informatique médicale et technologies de communication
PU-PH	GIOT Jean-Philippe	Chirurgie plastique, reconstructrice et esthétique ; Brûloogie
MCU-PH	GRAND Sylvie	Radiologie et imagerie médicale
PU-PH émérite	GRIFFET Jacques	Chirurgie infantile
MCU-PH	GUZUN Rita	Nutrition
PU-PH	HAINAUT Pierre	Biochimie et biologie moléculaire
PU-PH émérite	HALIMI Serge	Nutrition
PU-PH	HENNEBICQ Sylviane	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PU-PH	HOFFMANN Pascale	Gynécologie-obstétrique ; gynécologie médicale
PU-PH émérite	HOMMEL Marc	Neurologie
PU-MG	IMBERT Patrick	Médecine Générale

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH émérite	JOUK Pierre-Simon	Génétique
PU-PH	KAHANE Philippe	Physiologie
MCU-PH	KASTLER Adrian	Radiologie et imagerie médicale
PU-PH	KRAINIK Alexandre	Radiologie et imagerie médicale
PU-PH	LABARERE José	Epidémiologie, économie de la santé et prévention
MCU-PH	LABLANCHE Sandrine	Endocrinologie, diabète et maladies métaboliques
MCU-PH	LANDELLE Caroline	Bactériologie – virologie ; Hygiène hospitalière
PU-PH	LANTUEJOUL Sylvie	Anatomie et cytologie pathologiques
PR Ass. Méd.	LARAMAS Mathieu	Cancérologie ; radiothérapie
MCU-PH	LARDY Bernard	Biochimie et biologie moléculaire
MCU - PH	LE GOUELLEC LE PISSART Audrey	Biochimie et biologie moléculaire
PU-PH	LECCIA Marie-Thérèse	Dermato-vénérérologie
MCF Ass.MG	LEDOUX Jean-Nicolas	Médecine Générale
PU-PH émérite	LETOUBLON Christian	Chirurgie viscérale et digestive
PU-PH	LEVY Patrick	Physiologie
PU-PH	LONG Jean-Alexandre	Urologie
MCU-PH	Lupo Julien	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	MAIGNAN Maxime	Médecine d'urgence
PU-PH	MAITRE Anne	Médecine et santé au travail
MCU-PH	MALLARET Marie-Reine	Bactériologie – virologie ; Hygiène hospitalière
MCU-PH	MARLU Raphaël	Hématologie ; Transfusion
PR Ass. Méd.	MATHIEU Nicolas	Gastroentérologie ; hépatologie ; addictologie
MCU-PH	MAUBON Danièle	Parasitologie et mycologie
PU-PH	MAURIN Max	Bactériologie-virologie ; Hygiène hospitalière
MCU-PH	MC LEER Anne	Histologie, embryologie et cytogénétique
MCU-PH	MONDET Julie	Histologie, embryologie et cytogénétique
PU-PH	MORAND Patrice	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	MOREAU-GAUDRY Alexandre	Biostatistiques, informatique médicale et technologies de communication
PU-PH	MORO Elena	Neurologie
PU-PH	MORO-SIBILOT Denis	Pneumologie ; addictologie
MCU-PH	MORTAMET Guillaume	Pédiatrie
PU-PH	MOUSSEAU Mireille	Cancérologie ; radiothérapie
PU-PH émérite	MOUTET François	Chirurgie plastique, reconstructrice et esthétique ; brûrologie
MCF Ass.MG	ODDOU Christel	Médecine Générale
PR Ass. Méd.	ORMEZZANO Olivier	Cardiologie
MCU-PH	PACLET Marie-Hélène	Biochimie et biologie moléculaire
PU-PH	PAILHE Régis	Chirurgie orthopédique et traumatologie
PU-PH	PALOMBI Olivier	Anatomie
PU-PH	PARK Sophie	Hématologie ; Transfusion
PU-PH	PASSAGGIA Jean-Guy	Anatomie
PR Ass.MG	PAUMIER-DESBRIERES Françoise	Médecine Générale
PU-PH	PAYEN DE LA GARANDERIE Jean-François	Anesthésiologie-réanimation et médecine péri-opératoire
MCU-PH	PAYSANT François	Médecine légale et droit de la santé
MCU-PH	PELLETIER Laurent	Biologie cellulaire
PU-PH	PELLOUX Hervé	Parasitologie et mycologie
PU-PH	PEPIN Jean-Louis	Physiologie
PU-PH	PERENNOD Dominique	Médecine physique et de réadaptation

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH	PERNOD Gilles	Chirurgie vasculaire ; Médecine vasculaire
PU-PH	PIOLAT Christian	Chirurgie infantile
PU-PH	PISON Christophe	Pneumologie ; Addictologie
PU-PH	PLANTAZ Dominique	Pédiatrie
PU-PH	POIGNARD Pascal	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	POLACK Benoit	Hématologie ; Transfusion
PU-PH	POLOSAN Mircea	Psychiatrie d'adultes ; Addictologie
PU-PH	RAMBEAUD Jean-Jacques	Urologie
PU-PH	RAY Pierre	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PR Ass. Méd.	RECHE Fabian	Chirurgie viscérale et digestive
MCU-PH	RENDU John	Biochimie et biologie moléculaire
MCU-PH émérite	RIALLE Vincent	Biostatistiques, informatique médicale et technologies de communication
PU-PH	RIETHMULLER Didier	Gynécologie-obstétrique ; gynécologie médicale
PU-PH	RIGHINI Christian	Oto-rhino-laryngologie
PU-PH émérite	ROMANET Jean Paul	Ophthalmologie
PU-PH	ROSTAING Lionel	Néphrologie
PU-PH	ROUSTIT Matthieu	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
MCU-PH	ROUX-BUISSON Nathalie	Biochimie et biologie moléculaire
PR Ass.MG	ROYER DE VERICOURT Guillaume	Médecine Générale
PU-PH émérite	SARAGAGLIA Dominique	Chirurgie orthopédique et traumatologie
MCU-PH	SATRE Véronique	Génétique
PU-PH	SAUDOU Frédéric	Biologie cellulaire
PU-PH	SCHMERBER Sébastien	Oto-rhino-laryngologie
PU-PH	SCHWEBEL Carole	Médecine intensive-réanimation
PU-PH	SCOLAN Virginie	Médecine légale et droit de la santé
MCU-PH	SEIGNEURIN Arnaud	Epidémiologie, économie de la santé et prévention
PU-PH émérite	STAHL Jean-Paul	Maladies infectieuses ; Maladies tropicales
PU-PH	STANKE Françoise	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
MCU-PH	STASIA Marie-José	Biochimie et biologie moléculaire
PU-PH	STURM Nathalie	Anatomie et cytologie pathologiques
PU-PH	TAMISIER Renaud	Physiologie
PU-PH	TERZI Nicolas	Médecine intensive-réanimation
PU-PH	THEVENON Julien	Génétique
MCU-PH	TOFFART Anne-Claire	Pneumologie ; Addictologie
PU-PH	TONETTI Jérôme	Chirurgie orthopédique et traumatologie
PU-PH	TOUSSAINT Bertrand	Biochimie et biologie moléculaire
PU-PH	VALMARY-DEGANO Séverine	Anatomie et cytologie pathologiques
PU-PH	VANZETTO Gérald	Cardiologie
PU-PH	VUILLEZ Jean-Philippe	Biophysique et médecine nucléaire
PU-PH	WEIL Georges	Epidémiologie, économie de la santé et prévention
PU-PH	ZAOUI Philippe	Néphrologie
PU-PH émérite	ZARSKI Jean-Pierre	Gastroentérologie ; hépatologie ; addictologie

PU-PH : Professeur des Universités - Praticiens Hospitaliers
 MCU-PH : Maître de Conférences des Universités - Praticiens Hospitaliers
 PU-MG : Professeur des Universités de Médecine Générale
 MCU-MG : Maître de Conférences des Universités de Médecine Générale
 PR Ass. Méd. : Professeur des Universités Associé de Médecine
 PR Ass.MG : Professeur des Universités Associé de Médecine Générale
 MCF Ass.MG : Maître de Conférences Associé de Médecine Générale

Table des matières

Remerciements	8
Liste des abréviations	10
Table des illustrations	11
Résumé	12
Abstract.....	13
Introduction	14
Methods	16
1. Design	16
2. Standard protocol approvals, registrations, and patient consents.	16
3. Participants	16
4. Disability assessments	17
5. Muscle MRI.....	17
6. Clinical deficits.....	18
7. Activity limitations	19
8. Statistics analysis.....	19
Results	21
1. Participants	21
2. Mercuri scores per muscle.....	22
3. Relationship between Mercuri score and muscular weakness.....	26
4. Relationship between Mercuri score and PROM	29
5. Activity limitations	29

Discussion.....	32
Conclusion.....	37
References.....	38
Annexes : formulaire d'information et d'opposition	41
Serment d'hippocrate.....	44

Remerciements

A M. le Professeur Dominic Pérennou, *professeur universitaire - praticien hospitalier* : vous me faites le grand honneur de présider mon jury de thèse et êtes à l'origine de ce sujet qui a été le lien parfait avec mon master 2 de génétique et la spécialité médicale que j'ai choisie. Je vous remercie pour vos conseils, et pour le temps que vous m'avez offert pour en discuter ensemble. Veuillez trouver ici ma gratitude et l'expression de mon profond respect.

A M. le Professeur Pierre-Simon Jouk, *professeur universitaire - praticien hospitalier* : vous êtes à l'origine du centre de référence de l'arthrogrypose. Votre présence à ce jury est un immense honneur. J'espère que cette thèse aux confluences de la rééducation, la génétique et l'imagerie saura refléter au mieux la multidisciplinarité du fruit de votre travail.

A Mme la Docteur Véronique Bourg, *praticienne hospitalière* : Je vous remercie pour l'accueil que vous m'avez réservé dans votre service, pour tout ce que j'ai appris à vos côtés. La pédiatrie est un monde à part, je l'ai découvert avec plaisir et en chanson. Enfin, l'enfance est une étape cruciale dans la prise en charge de l'AMC, votre présence à ce jury était indispensable.

A Mme la Docteur Chantal Durand, *praticienne hospitalière* : vous me faites l'honneur de participer à mon jury de thèse. Il était important à mes yeux de présenter mon travail à un radiologue qui a participé activement à l'expertise des IRM musculaires du centre de référence de l'arthrogrypose. Je vous remercie de l'intérêt que vous avez porté à ma thèse.

A M. le Docteur Klaus Dieterich, *maître de conférence universitaire - praticien hospitalier* : Je n'aurai pas assez de place pour exprimer ma profonde reconnaissance pour ton aide précieuse dans l'élaboration de cette thèse. Depuis ces réunions zoom les soirs du premier confinement, aux dernières impromptues à l'HCE quand je débarquais à l'improviste en génétique. Merci pour ta confiance, tes conseils et ton enthousiasme au cours de ces deux dernières années.

A Louis, pour ta précieuse aide en statistique. **A Mariya et Thu**, pour l'incroyable travail d'analyse d'IRM : plus de 5000 muscles à classer un par un ! Sans vous, cette thèse n'aurait jamais été possible.

A tous les rééducateurs qui ont partagé avec moi leurs passions, leurs connaissances. A Justine en diabétologie, pour ces discussions musculaires entre deux ajustements de glycémie, un lien très apprécié avec ma spécialité en territoire sucré.

A toutes les équipes qui m'ont accueillie durant l'internat : Merci de m'avoir formée et accompagnée dans mon cursus, dans une ambiance toujours chaleureuse et bienveillante.

A ma famille : A mes parents, pour leur amour et leur soutien pendant toutes ces années, pour m'avoir offert les meilleures conditions possibles pour débuter et réussir mes études, pour leurs relectures assidues et les cours d'Excel par téléphone. A mes sœurs Anaïs et Célia, qui malgré les kilomètres qui nous séparent, sont présentes pour moi, à mes grands-parents et toute ma famille pour leurs encouragements. A Hélène pour cette bataille de « S » menée ensemble.

A B.

A mes co-internes : Solène et Antoine, je n'aurai pas pu espérer de meilleurs co-pioupious pour partager cette aventure qu'est l'internat, pour parcourir la France au fil des cours de DES. Je suis ravie d'avoir pu découvrir ma nouvelle région d'adoption à vos côtés, et d'avoir partagé avec vous ma passion pour la couture.

A tous les internes et assistants de MPR grenoblois que j'ai eu la chance de rencontrer et côtoyer pendant ces années d'internat : Rémi, Lucie, Lénaïc, Lisa, Camille, Jean, David, Ophélie, Adélie, Alicia et Alicia, Juliette, Cyrielle et Nicolas. Toujours prêts pour une sortie ski de fond – tartiflette, une après-midi jeux de société ou une soirée *MP-biR*. Sans vous l'internat aurait eu moins de saveurs.

A mes amis de médecine parisiens, Camille, Edouard, Flora, Lucie, Marion, Rémi... pour ces soirées cartes. **A mes amis urgentistes grenoblois** pour m'avoir fait découvrir le monde si vaste du houblon, **et amis chambériens** pour ces diners concerts à l'internat pendant le couvre-feu et ces sorties à la Féclaz, A tous les autres qui ont été présents pour moi.

A Arthur : pour ton soutien depuis toutes ces années, toujours présent dans les moments difficiles comme dans les meilleurs. Bientôt la fin de nos études, c'est un chapitre qui s'achève. Heureusement je n'ai pas le syndrome de la page blanche, j'ai plein d'idées pour la suite ! Prends ton stylo, on l'écrira ensemble. Je t'aime.

Liste des abréviations

6MWT : Six-minute-walking test

AMC : Arthrogryposis multiplex congenita

AUC : Area under curve

BMI : Body mass index

CI : Confidence intervals

DA : Distal Arthrogryposis

DMD : Duchenne muscular dystrophy

FIM : Functional Independence Measure

GOF : Gain of function

LL : Lower limbs

LOF : Loss of function

MACHAON : Mercuri Analysis Contribution on Handicap evaluation in ArthrogypOsis, a congenital Neuromuscular disease

MPS : Multiple pterygium syndrom

MRI : Magnetic Resonance Imaging

NMJ : Neuromuscular junction

NPV : Negative predictive values

PPV : Positive predictive values

PROM : Passive range of motion

RF : Rectus Femoris

SD : Standard deviation

T1W-TSE : T1-weighted turbo spin echo

UL : Upper limbs

VM : Vaste Medialis

Table des illustrations

Figure 1. Flow chart	21
Figure 2 : Individual muscles analysed by Mercuri scale.	23
Figure 3: MRI with individual muscle analysis of Mercuri score.....	24
Figure 4 : Upper limb, trunk and lower limb Mercuri score for Amyoplasia and DA.	25
Figure 5 : Relationship between Mercuri scores and muscular weakness of upper limbs (A) and lower limbs (B) for Amyoplasia (blue) and DA (red).....	27
Table 1 : General characteristics of the two groups. a.u = arbitrary unit.	22
Table 2 : Characteristics of individuals with Distal Arthrogryposis.	28
Table 3 : Ccorrelations of 6MWT with Mercuri score, muscle score, and joint testing score of lower limbs.....	31

Résumé

Introduction : L'arthrogrypose multiple congénitale (AMC) se définit par des limitations articulaires congénitales touchant au moins deux niveaux articulaires, conduisant à des handicaps multiples. La fonction musculaire est altérée. Le but de notre étude était d'évaluer la corrélation entre l'infiltration graisseuse musculaire avec les déficiences et limitations d'activité chez des adultes présentant une AMC.

Méthode: Cette étude évalue des adultes non sélectionnés adressés au centre de références de l'AMC entre 2010 et 2020. Tous ont bénéficié d'une évaluation pluridisciplinaire, ainsi qu'une IRM musculaire corps entier. Les muscles ont été analysés grâce à l'échelle de Mercuri, quantifiant l'infiltration graisseuse. Deux groupes étiologiques ont été formés selon l'étiologie Amyoplasie et Arthrogryposes Distales (AD), les deux formes les plus communes d'AMC.

Résultats : La moyenne d'âge des 53 adultes (34 femmes) était de 34 ans, 35 avaient une Amyoplasie, et 18 une Arthrogrypose Distale. Dans les deux groupes d'individus avec Amyoplasie et AD, nous avons retrouvé une corrélation significative entre le score Mercuri moyen et l'évaluation manuelle de la faiblesse musculaire aux membres supérieurs et inférieurs. De plus, dans l'Amyoplasie, les scores Mercuri des membres supérieurs et inférieurs étaient corrélés à l'amplitude articulaire passive, le score Mercuri des membres inférieurs était corrélé au test de marche de 6 minutes, et celui des membres supérieurs à la mobilité de la main dans l'espace. Dans l'AD, ces corrélations n'ont pas été retrouvées.

Conclusion : Notre étude est la première à étudier le lien entre imagerie médicale et aspects fonctionnels dans l'AMC. L'IRM musculaire, un puissant outil diagnostic, est aussi hautement adapté pour évaluer le handicap fonctionnel. Des scores Mercuri élevés étaient corrélés à de plus grandes déficiences et une moindre mobilité.

Mots clés : *AMC, arthrogrypose, IRM, infiltration graisseuse musculaire, handicap.*

Abstract

Introduction: Arthrogryposis multiplex congenita (AMC) defines multiple congenital contractures involving two or more body areas, leading to various disabilities. Muscle function is impaired. The purpose of this study was to evaluate the correlation between muscle fat infiltration and deficiencies and activity limitations in adults with AMC.

Methods: This analysis of data for unselected adults referred to the French AMC reference center from 2010 to 2020. All underwent a pluriprofessionnal disability evaluation and had an all-body muscular MRI. Each muscle was analysed with Mercuri score, which quantify muscle fat infiltration. Participants were divided into Amyoplasia and Distal Arthrogryposis (DA), the two most common forms of AMC.

Results: Mean age of the 53 participants (34 females) was 34 years; 35 had Amyoplasia and 18 had DA. In both groups of individuals with Amyoplasia and DA, we demonstrate a significant correlation between mean Mercuri score and manual testing of muscular weakness, for both upper and lower limbs. Furthermore, in Amyoplasia, Mercuri scores of upper and lower limb muscles were correlated with joint stiffness, Mercuri score of lower limb muscles was correlated with the six-minute-walking test, and Mercuri scores of upper limb and trunk muscles were correlated with the reaching capacity. In DA, no further correlation was found.

Conclusion: Our study is the first focus on the link between functional and imaging aspects in AMC. We show that Muscular MRI, a powerful diagnostic tool, is also highly useful for disability evaluation. Higher mercuri score were correlated with higher deficiencies and lower mobility.

Key words: *AMC, arthrogryposis, MRI, muscular fat infiltration, disability.*

Introduction

Arthrogryposis multiplex congenita (AMC) defines congenital contractures involving two or more body areas. The incidence is estimated between 1/3000 and 1/12 000¹⁻³.

Disability in adults is multiple: physical with reduced passive range of motion (PROM), muscle weakness, leading to activity limitation, and altered autonomy. Invisible and psychological issues like pain, anxiety, fatigue, difficulty in sexual life, altered self-esteem have also been described⁴.

In the two most common forms of arthrogryposis, Amyoplasia (“no muscle formation”) and Distal Arthrogryposis (DA), the primary condition leads to muscle alteration. In Amyoplasia, muscle tissue is absent or underdeveloped, with fatty or connective tissue replacement. DA mechanisms are multiple and can be linked to muscle weakness in sarcomeric structures, a lack of relaxation, or connective tissue dysfunctions. DA is characterized by mainly distal joint contractures^{1,5-8}.

Disability differs between Amyoplasia and DA with greater deficiencies in Amyoplasia than in other AMC types. In adults with Amyoplasia, muscle weakness is higher, grip strength is lower, passive joint mobility in major lower limb articulations are more altered. Most of them use compensatory strategies to perform daily life activity, and the majority needs some assistance⁹. Motor component of the Functional Independence Measure (FIM) is lower, requiring technical aids is more frequent, and adults with Amyoplasia are more often unable to drive a car, than other AMC type⁴.

Muscle strength is a major determining factor of functional abilities. In Duchenne muscular dystrophy, where muscles are progressively replaced by fat with age, distance walked in 6 minutes and muscle strength decreases with disease progression^{10,11}. The decline of muscle

strength is related to motor function loss^{12,13}. In Amyoplasia, Kroksmark *et al.* (2006) studied different ambulation capacities with isometric muscle strength. The most severely reduced isometric muscle strength was found in the non-ambulatory group, and they found good correlation between muscle strength and motor function¹⁴.

However, in AMC, relation between muscle weakness and motor function is sometimes difficult to analyse given articular deformation. Medical imaging with Magnetic Resonance Imaging (MRI) which is non-irradiating, can be a complementary method to analyse muscles.

Clinical correlation between muscle weakness, passive range of motion and fatty infiltration on MRI imaging has often not been studied, especially in AMC. Few studies have been performed on other muscular diseases like Pompe disease, muscular dystrophies, Duchenne muscular dystrophy (DMD) or GNE myopathy, where muscular MRI can be used to evaluate and follow the neuromuscular disease progression as it is correlated with muscle function and activities^{15–18}. Although AMC is a congenital non-progressive condition, by analogy with these neuromuscular diseases, we hypothesise that muscular fat infiltration evaluated with Mercuri score on MRI will be correlated with muscular weakness and PROM, and then with mobility namely the six-minute-walking test (6MWT) and reaching capacity.

The purpose of this study is to evaluate the correlation between muscle fat infiltration using Mercuri score based on muscular MRI, and deficiencies and activity limitations, in adults with Amyoplasia and Distal Arthrogryposes.

Methods

1. Design

This was an observational study of the cohort *Mercuri Analysis Contribution on Handicap evaluation in ArthrogypOsis, a congenital Neuromuscular disease* (MACHAON) of adults referred to the French AMC reference centre (Genetic and Neurorehabilitation Departments, University Hospital Grenoble-Alpes, France). Deficits, activity limitations, and participation restrictions were systematically analysed according to the international classification functioning. Here, we focus on the relationship between muscular infiltration and deficits and activity limitations.

2. Standard protocol approvals, registrations, and patient consents.

The study was performed in compliance with the Helsinki Declaration. All participants provided informed consent. According to French laws, as a noninterventional (observational) study, this research did not require approval by a national ethics committee. Individuals were informed by letters and have not objected to the use of their clinical data for research.

3. Participants

In total, 53 individuals with AMC were assessed from 2010 to 2020, and had an all-body MRI. They were divided into 2 subgroups according to clinical signs and molecular diagnosis: 35 with Amyoplasia and 18 with DA.

DA regroups a large variety of etiology, many gene mutations can lead to a gain of function of neuromuscular junction or sarcomere structure on the one hand, loss of function of neuronal system or multiple pterygium syndrome can be observed on the other hand.

4. Disability assessments

All participants underwent a 4-day multidisciplinary evaluation involving physicians (geneticists, physical medicine and rehabilitation physicians, physiotherapists, occupational therapists, and nurses). Systematic assessments involved analysis of the medical history in available medical records, clinical examinations, and a battery of tests.

5. Muscle MRI

Procedure

All MRI images were performed during the usual clinical care which was justified by the disease. Muscle MRI was performed on a 1.5T Siemens or General Electric MRI system at the department of Radiology and Medical Imaging. Participants were in supine position with forearms placed on the abdomen allowing acquisition of axial slices, at the trunk, upper limbs (UL) except for forearms, and lower limbs (LL). The imaging sequence was a T1-weighted turbo spin echo (T1W-TSE), with 8 mm section thickness, interspaced by 9 mm slice gaps. Forty-eight slices were respectively performed for upper and lower body parts. Slices of the upper body parts assessed shoulder girdle and arm muscles, whereas slices of the lower body parts assessed pelvic girdle and lower limb muscles.

Muscle targeted

We targeted one hundred muscles, fifty on each side, with upper limbs n=11, trunk n=8 and lower limbs n=31.

Mercuri score

The degree of muscle fat infiltration was assessed with at least 2 visible slices. We used the four-point scale proposed by Mercuri *et al.*¹⁹. Each muscle was staged as follows: 1. Normal appearance, 2. Mild involvement. 3. Moderate involvement. 4. Severe involvement.

This scaling was systematically performed by two independent operators, an experienced geneticist specialized in neuromuscular diseases (KD), and one radiologist who was blinded to the deficits (HTN for 39, MR for 15). In case of discordant assessments, a consensus was reached after discussion.

We a priori decided that to be kept in the final analysis, a muscle had to be scored in at least 80% of individuals. Final analysis has been performed on Mercuri scores of every available muscle average for left and right hemibodies.

A Mercuri score for each bilateral articulation (shoulder girdle, elbow, pelvic girdle, knee, and ankle) was determined with the mean of individual Mercuri analysis of muscles involved. Then an upper limb Mercuri score was determined (mean of shoulder girdle and elbow), a lower limb Mercuri score (mean of pelvic girdle, knee and ankle) and trunk Mercuri score.

6. Clinical deficits

Two principal clinical deficits were assessed by trained physiotherapists, muscular weakness, and passive range of motion. Because of the complexity of clinical pictures, muscular weakness was assessed by function (flexion, extension etc.) and not by each muscle, using the Medical Research Council scale (0–5) for muscle weakness. The grading was as follows: 0, no contraction; 1, flicker or trace of contraction; 2, active movement possible only with gravity eliminated; 3, active movement against gravity in the whole range of motion; 4, active movement against gravity and resistance; 5, quasi-normal strength. PROM was investigated for all main joints by conventional clinical examination and quantified in degrees. The range of motion of each movement (i.e. amplitude) was normalized with respect to the normal maximal movement, expressed as percentages.

Scores of muscular weakness and PROM were averaged for each joint and limb.

7. Activity limitations

Two principal activity limitations were assessed by trained physiotherapists and occupational therapists. Mobility was assessed by the 6MWT²⁰, performed by individuals who were asked to walk as self-preferred speed in a straight line on the floor with sequences of 30 meters, with a half-turn at each end. In case of interruption (fatigue, pain, dyspnea), patients were allowed to rest, until they felt able to restart.

The ability to reach some body targets with upper limbs was assessed by ad hoc test. Individuals sitting on a chair were instructed to successively touch their mouths, heads, necks, backs, opposite shoulders, ipsilateral shoulders, knees and feet, with both hands. Each successful touch scored two points if easily performed, and one point if performed with difficulty. The total score was calculated on both sides averaged and ranged from 0 to 16 points.

8. Statistics analysis

Descriptive statistics were performed on demographic and clinical variables. Continuous data were presented as mean (SD) and dichotomized data as number (%). Data for two groups (Amyoplasia vs DA) were compared by one-way ANOVA test and otherwise by chi-square test. Effect sizes were calculated by using the F values of the one-way ANOVA test, $\eta^2 = F * \text{degree of freedom}_{\text{between}} / (F * \text{degree of freedom}_{\text{between}} + \text{degree of freedom}_{\text{within}})$, and the χ^2 of the chi-square test, $\eta^2 = \sqrt{(\chi^2 / n) * \text{degree of freedom}}$. Effect sizes are given with $p < 0.05$; The values 0.01, 0.06 and 0.14 represent a small, medium and large effect for the one-way ANOVA test; values 0.1, 0.3, and 0.5 represent a small, medium, and large effect for the chi-square test.

The association between primary outcome (Mercuri score) and other clinical variables (muscle and joint testing, 6MWT and ad-hoc reaching test) were analyzed by linear regression, r and R² with 95% confidence intervals (CIs) were given. Because of Mercuri score for 3 parts supposed to be dependent on each other, corrected P values were set at 0.0083 (0.05/3 parts*2 groups)

for the associations between Mercuri scores and muscle testing, joint testing, and 6MWT, and at 0.0125 (0.05/2 parts*2 groups) for the associations between Mercuri scores and ad-hoc reaching test.

We further tested the value of Mercuri, muscle and joint testing scores to predict the good walking ability (6MWT ≥ 450) and satisfactory upper limbs reaching ability (ad-hoc test ≥ 14) in individuals with Amyoplasia. We calculated areas under the receiver operating characteristic (ROC) curve (AUC). Optimal cut-off values were determined by the Youden Index. From the cut-off values, we calculated sensitivity, specificity, positive predictive values (PPV) and negative predictive values (NPV).

Missing data were few, so imputation was not performed. Unless otherwise indicated (corrections), two tailed P<0.05 was considered statistically significant. Statistical analysis involved using Stata 15.1.

Results

1. Participants

Fifty-three individuals with AMC were consecutively assessed from 2010 to 2020, among whom 35/53 (66%) with Amyoplasia and 18/53 (34%) with DA (see flow chart in Figure 1). They were young (mean age 34 (11.9) years) with a normal body mass index (mean 23 (6.3) kg/m²); mostly female (34/53, 64.2%). Their disability was moderate with a FIM score of 112/126 (13.7). General characteristics of the two groups, Amyoplasia and DA are presented in Table 1. They were comparable in terms of age, sex ratio, BMI, distance walked during 6 min, passive range of motion of the upper limbs. Individuals with Amyoplasia had greater muscular weakness of upper and lower limbs, lower PROM of the lower limbs, worst reaching ability, and finally a greater global disability score FIM and (higher disability). No data were missing for the primary variables, mean Mercuri score and muscular testing score. There were few missing data for secondary variables: PROM (n=3, 5.7%), mobility and disability (n=1, 1.9%).




Figure 1. Flow chart.

Table 1 : General characteristics of the two groups. a.u = arbitrary unit.

	Amyoplasia (n=35)	DA (n=18)	P-value
Clinical characteristics			
Age, y	33.9 (12.8)	33.2 (11.4)	F=0, p=0.856
Female, (%)	22 (62,8)	12 (66.7)	X ² =0.1, p=0.784
BMI, kg/m ²	23.96 (6.5)	22.10 (5.8)	F=1, p=0.313
Muscle testing (n=53)			
Upper limbs, a.u (0-5)	3.3 (1.3)	4.4 (0.6)	F=11.9, p=0.001, η ² =0.19
Lower limbs, a.u (0-5)	2.8 (1.2)	3.9 (1.1)	F=11.8, p=0.001, η ² =0.19
Articular testing, a.u (0-1), (n=50)			
Upper limbs,	0.7 (0.2)	0.8 (0.2)	F=0.6, p=0.435
Lower limbs,	0.5 (0.2)	0.7 (0.2)	F=5.9, p=0.019, η ² =0.11
Activity limitations, (n=52)			
6MWT, m	310.0 (200.2)	357.1 (176.9)	F=0.7, p=0.405
Reaching, a.u (0-16)	9.3 (5.3)	13.1 (3.2)	F=7.7, p=0.008, η ² =0.13
Disability (FIM 18-126) (n=52)			
	109.7(15)	117.7(7.2)	F=4.3, p=0.043, η ² =0.08

2. Mercuri scores per muscle

Figure 2 presents all muscles analysed. Some muscles were non analysable because of incomplete muscle imaging, orthopaedic material artefact, or because of movement artefact and anatomic difficulties to recognize each muscle (scoliosis for example). Muscles with less than 80% of Mercuri score determined were discarded: supraspinatus (less than two MRI sections), and sternocleidomastoid muscle, teres major, muscles of abdominal wall, quadratus lumborum, psoas, iliacus, gluteus medius, gluteus minimum, pectineus (part of muscle no explored by MRI section).

Final analyses were performed on 80 muscles. UL n=9, Trunk n=5, LL n=26. Figure 3 presents muscular MRI sections of three patients with different genotype and phenotype.




Figure 2 : Individual muscles analysed by Mercuri scale. The first group ($n=6$) was included in shoulder analysis, the second group ($n=3$) in elbow analysis: both were included in upper limb Mercuri score ($n=9$). The third group ($n=5$) was included in trunk Mercuri score. The fourth group ($n=9$) was included in pelvic girdle analysis, the fifth group ($n=8$) in knee analysis, and the sixth group in ankle analysis ($n=9$). The last three groups were included in the lower limb Mercuri score ($n=26$). Muscles discarded were Sternocleidomastoid muscle, Supraspinatus, Teres major, Muscles of abdominal wall, Quadratus lumborum, Psoas, Iliacus, Gluteus medius, Gluteus minimum, Pectineus.







A Female, 25 years, DA : TNNT3 mutation GOF on protein function	UL Mercuri score : 1.00  Pectoralis major Deltoid Pectoralis minor Infra- supinatus Sub- scapularis	LL Mercuri score: 1.19  Rectus femoris Vastus intermedius Vastus medialis Vastus lateralis Sartorius Adductor longus Adductor brevis Adductor magnus Gracilis Biceps femoris Semimembranosus Semitendinosus
B Male, 27 years DA : TTN mutation LOF on protein function	UL Mercuri score: 2.63  4 2 4 2 3	LL Mercuri score: 2.92  4 1 2 2 2 2 2 4 4 4
C Female, 28 years Amyoplasia	UL Mercuri score: 4.00  All 4	LL Mercuri score: 3.35  RF 3 VM 3 All the rest 4

Figure 3: MRI with individual muscle analysis of Mercuri score. Line A, TNNT3: mutation in tropomyosin gene. GOF: gain of function, Muscles are identified, all have a Mercuri score = 1. Line B, TTN: mutation in titin gene. LOF: loss of function. Line A and B: muscles are analyzed in the same order. Line C, Amyoplasia etiology. RF = Rectus Femoris, VM = Vaste Medialis. All muscle except these two have a Mercuri score= 4.

Figure 4 compares the mean Mercuri scores of the two groups of individuals for the upper, lower limb muscles and the trunk muscles. For Amyoplasia, upper limb Mercuri score was 2.40 (0.93) lower limb Mercuri score was 2.64 (0.80) and trunk Mercuri score was 1.93 (0.82). For DA, upper limb Mercuri score was 1.29 (0.52), lower limb Mercuri score was 1.64 (0.77) and trunk Mercuri score was 1.50 (0.66). Fat infiltration was higher in Amyoplasia than in DA for the upper limbs ($F=21.58$; $p<0.001$; $\eta^2=0.3$), and the lower limbs ($F=19.12$; $p<0.001$; $\eta^2=0.27$), but not for the trunk ($F=3.75$; $p=0.059$).




Figure 4 : Upper limb, trunk and lower limb Mercuri score for Amyoplasia (blue) and DA (red). * means $p<0.001$. UL n=53, trunk n=52 and LL n=53.

3. Relationship between Mercuri score and muscular weakness

For Amyoplasia, we found a strong correlation between Mercuri scores and manual muscular testing scores for both upper limbs ($r = -0.93$, 95%CI [-0.96; -0.87], $p < 0.001$) and the lower limbs ($r = -0.9$, 95%CI [-0.94; -0.83], $p < 0.001$). These results indicated that the mean Mercuri scores explained almost all the information contained in muscle weakness, 86.2% for the upper limbs (95%CI [78.2; 94.2]), and 81.7% for the lower limbs (95%CI [71.5; 92]).

For DA, the correlation between mean Mercuri scores and manual muscular score were lower but significant both for upper limb muscles ($r = -0.66$, 95%CI [-0.87; -0.33], $p = 0.003$) and lower limb muscles ($r = -0.68$, 95%CI [-0.88; -0.31], $p = 0.002$). These results indicated that the mean Mercuri scores explained part of information contained in muscle weakness, 42.9% for the upper limbs (95%CI [12.1; 73.7]), and 46% for the lower limbs (95%CI [15.8; 76.2]). These relationships are displayed in Figure 5. Table 2 presents characteristics of individuals with DA, with two subgroups according to the effect on protein function: gain of function (GOF) or loss of function (LOF). These two subgroups present a different repartition with a lower fat infiltration and less muscular weakness in the GOF subgroup in comparison with the LOF subgroup. For upper limb muscles, the mean Mercuri score was 1.09 (0.17) vs 1.70 (0.74) and manual muscular weakness score was 4.47 (0.30) vs 4.30 (0.62) for respectively GOF subgroup and LOF subgroup. For lower limbs, the mean Mercuri score was 1.27 (0.57) vs 2.38 (0.92) and manual muscular weakness score was 4.10 (0.7) vs 3.60 (1.60), for respectively GOF subgroup and LOF subgroup.




Figure 5 : Relationship between Mercuri scores and muscular weakness of upper limbs (A) and lower limbs (B) for Amyoplasia (blue) and DA (red).

Table 2 : Characteristics of individuals with Distal Arthrogryposis. General characteristics, genetic, effect on protein function, Mercuri scores and muscular testing scores for upper limbs and lower limbs. GOF: gain of function, LOF loss of function. NMJ: neuromuscular junction, MPS: multiple pterygium syndrome, UL: upper limbs, LL: lower limbs.

Patients	Age	Sex	Gene muted	Effect on protein function	Structure involved or syndrome	Mercuri score		Muscle testing	
						UL	LL	UL	LL
Patient 14	31	F	<i>PIEZ02</i>	GOF	NMJ	1.00	1.15	4.50	3.79
Patient 15	26	H	<i>PIEZ02</i>	GOF	NMJ	1.00	1.00	5.00	4.64
Patient 29	37	H	<i>PIEZ02</i>	GOF	NMJ	1.00	1.00	4.17	3.57
Patient 34	22	F	<i>PIEZ02</i>	GOF	NMJ	1.00	1.00	5.00	4.83
Patient 40	32	H	<i>PIEZ02</i>	GOF	NMJ	1.00	1.15	4.13	3.63
Patient 54	21	F	<i>PIEZ02</i>	GOF	NMJ	1.48	1.11	3.21	2.98
Patient 51	28	F	<i>MYH3</i>	GOF	Sarcomere	1.00	2.03	4.47	2.97
Patient 31	22	F	<i>TNNI2</i>	GOF	Sarcomere	1.33	1.60	4.00	4.92
Patient 10	25	F	<i>TNNT3</i>	GOF	Sarcomere	1.00	1.19	5.00	4.72
Patient 41	30	F	<i>TNNT3</i>	GOF	Sarcomere	1.00	1.21	4.13	3.48
Patient 56	34	H	<i>TPM2</i>	GOF	Sarcomere	1.00	1.38	5.00	4.67
Patient 02	50	H	Unknown	GOF	unknown	1.25	1.39	5.00	5.00
Mean score (SD) (n=12)						1.09 (0.17)	1.27 (0.57)	4.47 (0.30)	4.10 (0.7)
Patient 24	33	F	<i>ECEL1</i>	LOF	Neuronal	1.13	1.70	4.75	4.56
Patient 25	42	F	<i>ECEL1</i>	LOF	Neuronal	1.13	1.87	4.63	4.63
Patient 07	36	F	<i>ECEL1</i>	LOF	Neuronal	1.79	2.48	4.27	4.13
Patient 01	68	F	<i>TRPV4</i>	LOF	Neuronal	2.54	3.88	3.63	0.83
Patient 06	27	H	<i>TTN</i>	LOF	MPS	2.63	2.92	3.50	2.50
Patient 46	34	F	<i>CHRNG</i>	LOF	MPS	1.00	1.42	5.00	4.97
Mean score (SD) (n=6)						1.70 (0.74)	2.38 (0.92)	4.30 (0.62)	3.60 (1.6)

4. Relationship between Mercuri score and PROM

For Amyoplasia, we found a correlation between Mercuri scores and PROM scores for both upper limbs ($r = -0.8$, 95%CI [-0.88; -0.69], $p < 0.001$) and the lower limbs ($r = -0.74$, 95%CI [-0.85; -0.57], $p < 0.001$). These results indicated that the mean Mercuri scores explained around half of information contained in PROM, 63.6% for the upper limbs (95%CI [44.7; 82.3]), and 55.3% for the lower limbs (95%CI [33.8; 76.8]).

For DA, no correlation was found between mean Mercuri scores and PROM score for upper limbs ($r = -0.26$, 95%CI [-0.67; 0.39], $p = 0.29$) and for lower limbs ($r = -0.53$, 95%CI [-0.74; -0.07], $p = 0.025$). These results indicated that the mean Mercuri scores explained only a small part of PROM, only 7% for the upper limbs (95%CI [-13.3; 27.2]), and 27.6% for the lower limbs (95%CI [-3.8; 58.9]).

5. Activity limitations

Mobility and Mercuri Score

Seven individuals could not walk more than 100 meters in 6MWT, 6 with Amyoplasia (17%) and one with DA (6%).

We found a strong correlation between lower limb Mercuri score and the 6MWT for Amyoplasia ($r = -0.78$, 95%CI [-0.89; -0.63], $p < 0.001$), with Mercuri score explaining 60.9% of 6MWT distance (95%CI [41.7; 80.1]). In contrast, trunk or upper limb muscles did not influence gait, with no correlation between the 6MWT and the mean Mercuri scores in Amyoplasia of trunk muscles ($r = -0.32$, 95%CI [-0.6; -0.1], $p = 0.072$), or of the upper limb muscles ($r = 0.23$, 95%CI [-0.06; 0.52], $p = 0.188$). Thus, only little information was contained in the mean Mercuri scores of trunk (10.1%, 95%CI [1; 35.9]) and of upper limbs (4.6%, 95%CI [-3.6; 27]).

For DA, there was no significant correlation between Mercuri scores and 6MWT, in upper limb, trunk and lower limb muscles, respectively: $r = -0.59$, 95%CI [-0.87; 0.11], $p=0.009$; $r = -0.42$, 95%CI [-0.77; 0.67], $p=0.082$; and $r = -0.44$, 95%CI [-0.79; 0.36], $p=0.068$. The capacity of 6MWT could be partly explained by the Mercuri scores of upper limbs (35.2%, 95%CI [-1.2; 75.7]), trunk (17.7%, 95%CI [-44.9; 59.3]), and lower limbs (19.3%, 95%CI [-13; 62.4]).

Reaching and Mercuri Score

For Amyoplasia, we found a correlation between Mercuri scores and reaching score for both upper limb muscles ($r = -0.88$, 95%CI [0.8; 0.94], $p<0.001$), and trunk muscles ($r = -0.88$, 95%CI [0.8; 0.94], $p<0.001$). These results indicated that the mean Mercuri scores explained part of information contained the mobility of the hand in space, 71% for upper limbs (95%CI [56.3; 86.7]) and 52% for the trunk (95%CI [30.2; 73.8]).

For DA, no correlation was fond between mean Mercuri scores and reaching score for both upper limb muscles ($r = -0.23$, 95%CI [-0.78; 0.26], $p=0.37$) and the trunk muscles ($r = -0.17$, 95%CI [-0.77; 0.48], $p=0.514$). These results indicated that the mean Mercuri scores don't explain information contained reaching capacity, 5% (95% CI [(-6.8; 60.1)]) for upper limbs and 0.3% for the trunk (95%CI [-23; 59.3]).

Predicting walking and reaching capacity

For individuals with Amyoplasia, we tried to predict the walking capacity (6MWT ≥ 450 m) from three muscular factors of the lower limbs: the Mercuri score of muscular imaging on one hand, the manual testing of muscle weakness and the manual testing of PROM on the other hand. All factors predicted walking capacity. For the manual testing of muscular weakness results were as follow: AUC =0.9 (95%CI [0.79; 1], $p<0.001$), the cut-off =3.46, sensitivity 0.88, specificity 0.89, PPV 0.73, NPV 0.96. For Mercuri score, results were as follows: AUC=

0.92 (95%CI [0.81; 1], p<0.001), the cut-off =2.17, sensitivity 0.96, specificity 0.88, PPV 0.96, NPV 0.89. When using the joint testing score, results were as follow: AUC =0.88 (95%CI [0.76; 1], p<0.001), the cut-off =0.61, sensitivity 0.71, specificity 0.83, PPV 0.58, NPV 0.89. We further ran a stepwise multiple linear regression to explore the associations between 6MWT and Mercuri score, muscle weakness and joint testing scores (Table 3). Finally, we found that Mercuri score of lower limb muscles was the best variable to explain the 6MWT (Beta= -196, 95%CI [-253; -140], p<0.001).

Table 3 : Correlations of 6MWT with Mercuri score, muscle score, and joint testing score of lower limbs.

Variables	6MWT	Mercuri score	Muscle testing score	Joint testing score
6MWT	1	r = -0.78 p< 0.001	r = 0.73 p< 0.001	r = 0.7 p< 0.001

We also employed the Mercuri score, muscle weakness and joint testing scores of upper limbs to predict the normal capacity of reaching (ad hoc test ≥ 14), respectively. The prediction accuracies were all high. For Mercuri score, results were as follows: AUC= 0.93 (95%CI [0.85;1], p<0.001), the cut-off =1.75, sensitivity 0.91, specificity 0.73, PPV 0.85, NPV 0.85. When using the muscular testing score, results were as follow: AUC =0.95 (95%CI [0.88;1], p<0.001), the cut-off=4.13, sensitivity 0.91, specificity 0.96, PPV 0.92, NPV 0.96. When using the joint testing score, results were as follow: AUC =0.92 (95%CI [0.82;1], p<0.001), the cut-off =0.79, sensitivity 0.9, specificity 0.81, PPV 0.67, NPV 0.95.

Discussion

This study focused on muscular imaging, extends the first study published from our cohort of adults with AMC referred to the French reference centre from 2010 to 2020⁴. We retrospectively analysed data of 53 individuals: 35 with Amyoplasia and 18 with DA. All had an all-body MRI, and muscles were analysed with the Mercuri scale. A mean Mercuri score of upper limbs, lower limbs and trunk were calculated with their respective muscles.

A relationship between muscular weakness and muscular MRI has already been found in some neuromuscular diseases like Duchenne muscular dystrophy¹⁷, Pompe disease¹⁸, and GNE myopathy¹⁵. It remained to be investigated in AMC. By analogy with these diseases, we hypothesized that in AMC also, muscular fat infiltration evaluated with Mercuri score on MRI, will be correlated with muscular weakness and PROM, and then with mobility (6MWT and reaching capacity). It has been confirmed for Amyoplasia, and partially for DA.

Our primary outcome was the Mercuri score. Most muscles in adults with AMC were analysable. Nine muscles had been discharged because of missing MRI sections, which can easily be resolved with new MRI protocol. Mercuri scores were lower in DA than in Amyoplasia with significant results on limbs, which is concordant with the differences in physiopathology of the two groups. Amyoplasia is characterized by the absence of muscles, whereas in DA, mechanisms are multiple and present less fat infiltration in muscles⁶.

On functional evaluations: our study finds a higher muscular weakness and less PROM in Amyoplasia than in DA, which is concordant with other studies^{4,6,7}. On activity limitations, there were no difference on 6MWT between Amyoplasia and DA, but in DA, the reaching capacity was more efficient, and disability was less important.

Is MRI useful?

Our study focuses on the link between functional and imaging aspects in AMC. We show that muscular MRI, a powerful diagnostic tool, is also highly useful for disability evaluation.

For Amyoplasia, we demonstrate a significant correlation between mean Mercuri score evaluated on muscular MRI, and manual testing of muscular weakness or PROM for both upper and lower limbs. The deficiencies were significantly linked with the replacement of muscles by fat tissue (high Mercuri score). Mercuri scores of limbs were correlated and predicted their linked activity: upper limbs and trunk with reaching capacity, and lower limbs with 6MWT.

For DA, Mercuri score was significantly correlated with manual muscular weakness testing. Nevertheless, no correlation between Mercuri score and PROM or activity limitations, was found. AMC is defined by a phenotype, and DA regroups many physiopathologies in the same group. Many systems have been proposed to classify various forms of arthrogryposis and all have limits²¹. In our study, four individuals present implication of neurogenic gene (*ECEL1*, *TRPV4*), two Escobar syndromes (*TTN* and *CHRNG* gene mutated), inducing a loss of function of protein involved. The 12 other individuals present a gain of function of protein involved, with five individuals with sarcomere protein altered (*TNNI2*, *TNN3*, *TPM2*, *MYH3*), six with mutation in a mechanosensitive cation channel (*PIEZ02*) and one unknown. As we have shown in Table 2, fat infiltration was variable between these subgroups. More studies are necessary with more individuals to identify subgroups of DA where Mercuri score evaluation on muscular MRI can be pertinent to study deficits and activity limitations.

What impact?

Parents of children with AMC are concerned by their future autonomy. Prediction of walking ability constitutes a recurrent question during medical consultations. In the study of functional

outcome conducted by Sells *et al.* in 1996, 7 in 38 children were non ambulant²⁷. Thus, finding prognostic factors is a challenge.

We found that Mercuri score was the best predictor of 6MWT results in comparison with manual muscular testing and PROM. It is tempting to hypothesize that the muscular fat infiltration evaluated on MRI during childhood could be a predictive element of future capacities. Our results should be analysed carefully. On the one hand, our population was adults with AMC, with a single functional evaluation. On the other hand, recruitment at the reference centre allows us to examine individuals with heterogeneous medical management. Other elements have been reported to have an impact on achievement of motor function like joint contractures and joint positions at birth^{14,28}. Furthermore, the multidisciplinary care during childhood with rehabilitation and orthotic management, can improve PROM, muscular strength⁷, and modify ambulation ability. Surgery history in our cohort was multiple and individually based⁴.

Study limitations

DA is mainly characterized by distal joint contractures. The lack of muscle testing data and Mercuri score or the forearm in the upper limb analysis might have underestimated mean muscle weakness and Mercuri score. This is also true for the lack of wrist PROM and upper limb joint stiffness. From a functional point of view, grasping capacity is an important part of the upper limb function in association with reaching capacity. These are key factors for autonomy in daily life activities⁹. This aspect could not be studied.

For Mercuri analysis based on muscular MRI, one hundred muscles for each individual have been analysed. This represents non negligible time for untrained radiologists, and thus can't be used in clinical routine. The first possibility to decrease the time spent on this analysis would

be to reduce the number of muscles by selecting the ones with the most important statistical impact for each function during the muscular testing. One could suggest selecting muscles from one side of the body. Indeed, the typical presentation of individual with AMC is symmetrical. In this study we decide to analyse both left and right sides together, to minimize dominance influences with missing data. Time-efficient muscle segmentation is a crucial issue to allow a complete imaging analysis in clinical practice. Nakayama *et al.* developed an algorithm to automatically calculate the Mercuri grades from computed tomography (CT) and T1-weighted MR images, on seven patients affected by limb girdle muscular dystrophy (LGMD). Correlation between the Mercuri grade calculated and visual was significant²².

Mercuri score has also some limits: it is semiquantitative, and skill based. Moreover, muscular atrophy is not considered. Other methods should be investigated. Recent progress in artificial intelligence allows us to consider deep learning for an automatic analysis of fat infiltration, and muscle volume. When fat infiltration was low, the visual score was strongly correlated with a quantitatively obtained fat fraction²³. Semiautomatic segmentation method using active shape model²⁴, or MRI analysis of an entire muscle region using texture features without distinguishing any muscle groups²⁵ have been explored. Ding *et al.* have demonstrated that in four muscles of the thigh, compared with manual segmentation, the automatic segmentation performed in less than two minutes, were higher reproductive in fat fraction estimation, and can give muscle volume with a limited absolute difference²⁶.

These approaches are promising but need more research to be transferred into clinical practice in AMC. Indeed, these automatic analyses were not suitable for individuals with severe fat infiltration, which represents the most important proportion in our cohort. Population sizes were reduced in all studies, and only few muscles, mostly thighs, have already been analysed.

Perspectives

Finally, AMC is a congenital non-progressive condition. Further studies are needed to evaluate the correlation between Mercuri score and deficiencies or activity limitations at different ages from birth to adulthood, relying on a long-term follow-up with repetitive muscular MRIs. It would also be interesting to further study the relationship between Mercuri score and restriction of participation elements. Indeed, individuals with AMC developed multiple compensations to improve their autonomy in daily life activities⁹.

Conclusion

Muscular MRI is routinely performed in adults evaluated in the French AMC reference centre. It has become a powerful tool for diagnostic purposes. Our study confirms that muscular MRI with Mercuri score is also interesting to appreciate muscular capacities in Amyoplasia and in Distal Arthrogryposis, as it was correlated with muscular weakness. Furthermore, in Amyoplasia, Mercuri score of lower limb muscles was correlated with ambulation ability (6MWT), and Mercuri score of upper limb muscles was correlated with reaching capacity. In Amyoplasia, muscle is the main determinant of ambulation capacity. We have shown that both lower limb muscular characteristics, namely fat infiltration of muscle evaluated on imaging, and muscular weakness evaluated by manual testing, had a strong capacity to predict the ability to walk, and Mercuri score was the best predictor of 6MWT results in comparison with manual muscular testing and passive range of motion.

These findings will help us to tailor the multidisciplinary supports like surgery, orthopaedic devices, and rehabilitation including training with physiotherapist or occupational therapist, on an individual basis, to maximize functional capacities.


VU ET PERMIS D'IMPRIMER

Grenoble le 10 septembre 2021

LE DOYEN



LE PRESIDENT DE LA THESE



Pr Dominic PERENNOM
Médecine Physique et Réadaptation
Institut de Rééducation
Hôpital Sud - CHU de Grenoble
BP 338 - 38484 ECHIROLLES Cedex
Tél. 04 76 76 80 88
N° 33061099999/00

References

1. Hall, J. G. Arthrogryposis multiplex congenita: etiology, genetics, classification, diagnostic approach, and general aspects. *J. Pediatr. Orthop. Part B* **6**, 159–166 (1997).
2. Kimber, E. AMC: amyoplasia and distal arthrogryposis. *J. Child. Orthop.* **9**, 427–432 (2015).
3. Dahan-Oliel, N. *et al.* International multidisciplinary collaboration toward an annotated definition of arthrogryposis multiplex congenita. *Am. J. Med. Genet. C Semin. Med. Genet.* **181**, 288–299 (2019).
4. Dai, S. *et al.* Disability in adults with arthrogryposis is severe, partly invisible, and varies by genotype. *Neurology* **90**, e1596–e1604 (2018).
5. Bamshad, M., Jorde, L. B. & Carey, J. C. A revised and extended classification of the distal arthrogryposes. *5*.
6. Kimber, E. Causes, Consequences and Clinical Course in Amyoplasia and Distal Arthrogryposis. *64*.
7. Kowalczyk, B. & Feliś, J. Arthrogryposis: an update on clinical aspects, etiology, and treatment strategies. *Arch. Med. Sci.* **1**, 10–24 (2016).
8. Ravenscroft, G. *et al.* Fetal akinesia: review of the genetics of the neuromuscular causes. *J. Med. Genet.* **48**, 793–801 (2011).
9. Steen, U., Wekre, L. L. & Vøllestad, N. K. Physical functioning and activities of daily living in adults with amyoplasia, the most common form of arthrogryposis. A cross-sectional study. *Disabil. Rehabil.* **40**, 2767–2779 (2018).
10. McDonald, C. M. *et al.* The 6-minute walk test and other clinical endpoints in duchenne muscular dystrophy: Reliability, concurrent validity, and minimal clinically important differences from a multicenter study. *Muscle Nerve* **48**, 357–368 (2013).

11. Hogrel, J.-Y. *et al.* Normalized grip strength is a sensitive outcome measure through all stages of Duchenne muscular dystrophy. *J. Neurol.* **267**, 2022–2028 (2020).
12. Scott, O. M., Hyde, S. A., Goddard, C. & Dubowitz, V. Quantitation of muscle function in children: A prospective study in duchenne muscular dystrophy. *Muscle Nerve* **5**, 291–301 (1982).
13. Nunes, M. F. *et al.* Relationship between muscle strength and motor function in Duchenne muscular dystrophy. *Arq. Neuropsiquiatr.* **74**, 530–535 (2016).
14. Kroksmark, A.-K., Kimber, E., Jerre, R., Beckung, E. & Tulinius, M. Muscle involvement and motor function in amyoplasia. *Am. J. Med. Genet. A.* **140A**, 1757–1767 (2006).
15. Liu, C.-Y. *et al.* Skeletal muscle magnetic resonance biomarkers in GNE myopathy. *Neurology* 10.1212/WNL.0000000000011231 (2020) doi:10.1212/WNL.0000000000011231.
16. Xie, Z. *et al.* Value of muscle magnetic resonance imaging in the differential diagnosis of muscular dystrophies related to the dystrophin-glycoprotein complex. *Orphanet J. Rare Dis.* **14**, 250 (2019).
17. Brogna, C. *et al.* Functional levels and MRI patterns of muscle involvement in upper limbs in Duchenne muscular dystrophy. *PLOS ONE* **13**, e0199222 (2018).
18. Figueiroa-Bonaparte, S. *et al.* Muscle MRI Findings in Childhood/Adult Onset Pompe Disease Correlate with Muscle Function. *PLOS ONE* **11**, e0163493 (2016).
19. Mercuri, E. *et al.* A short protocol for muscle MRI in children with muscular dystrophies. *Eur. J. Paediatr. Neurol.* **6**, 305–307 (2002).
20. Guyatt, G. H., Sullivan, M. J., Thompson, P. J. & Fallen, E. L. The 6-minute walk: a new measure of exercise capacity in patients with chronic heart failure. *CAN MED ASSOC J* **132**, 5 (1985).
21. Hall, J. G., Kimber, E. & Dieterich, K. Classification of arthrogryposis. *Am. J. Med. Genet. C Semin. Med. Genet.* **181**, 300–303 (2019).

22. Nakayama, T., Ishiyama, A., Murakami, T., Kimura, E. & Kuru, S. Automatic calculation of Mercuri grades from CT and MR muscle images. *Brain Dev.* **41**, 870–877 (2019).
23. Wokke, B. H. *et al.* Comparison of dixon and T1-weighted MR methods to assess the degree of fat infiltration in duchenne muscular dystrophy patients: Visual and Quantitative MRI in DMD. *J. Magn. Reson. Imaging* **38**, 619–624 (2013).
24. Kemnitz, J. *et al.* Clinical evaluation of fully automated thigh muscle and adipose tissue segmentation using a U-Net deep learning architecture in context of osteoarthritic knee pain. *Magn. Reson. Mater. Phys. Biol. Med.* **33**, 483–493 (2020).
25. Rodrigues, R. & Pinheiro, A. M. G. Segmentation of Skeletal Muscle in Thigh Dixon MRI Based on Texture Analysis. *ArXiv190404747 Cs* (2019).
26. Ding, J. *et al.* Deep learning-based thigh muscle segmentation for reproducible fat fraction quantification using fat–water decomposition MRI. *Insights Imaging* **11**, 128 (2020).
27. Sells, J. M., Jaffe, K. M. & Hall, J. G. Amyoplasia, the Most Common Type of Arthrogryposis: The Potential for Good Outcome. *Pediatrics* **97**, 225–231 (1996).
28. Donohoe, M., Pruszynski, B., Rogers, K. & Bowen, J. R. Predicting Ambulatory Function Based on Infantile Lower Extremity Posture Types in Amyoplasia Arthrogryposis. *J. Pediatr. Orthop.* **39**, e531–e535 (2019).

Annexes : formulaire d'information et d'opposition



NOTICE D'INFORMATION SUR UN PROJET DE RECHERCHE

Mercuri Analysis Contribution on Handicap evaluation in ArthrogypOsis, a congenital Neuromuscular disease.

MACHAON

Madame, Monsieur,

Cette fiche vous informe d'un projet de recherche, planifié ou en cours, organisé par les services de génétique et de MPR neurologique, en collaboration avec le service d'imagerie médicale du C.H.U. Grenoble Alpes. **Nous vous remercions d'avance d'en prendre connaissance.**

DESCRIPTIF DE L'ETUDE

Cette recherche consiste à évaluer l'intérêt d'une imagerie musculaire par IRM dans l'évaluation des déficits et du handicap chez les patients présentant une arthrogrypose multiple congénitale.

Cette étude devrait débuter en juin 2021 et il est prévu qu'elle s'achève en octobre 2021.

BENEFICES, CONTRAINTES ET RISQUES

Ce projet utilisera les données du dossier médical de patients pris en charge dans ces services, pour faire progresser les connaissances et les pratiques médicales. Seules des données acquises dans le cadre d'une prise en charge seront utilisées.

La prise en charge d'un patient n'est aucunement modifiée par l'utilisation de ses données médicales ou par son opposition à cette recherche.

CONFIDENTIALITE DES DONNEES

Dans le cadre de cette recherche, un traitement de données personnelles va être mis en œuvre.

Les données que nous prévoyons de collecter sont les suivantes :

- Âge, sexe, poids si mentionné
- Antécédents médicaux et chirurgicaux
- Imagerie par IRM
- Quotation de la force musculaire
- Mesure des amplitudes articulaires
- Distance parcourue au test de marche de 6 minutes
- Mobilité de la main dans l'espace et différentes préhensions possible
- Score MIF (mesure d'indépendance fonctionnelle)
- Capacité ambulatoire et d'équilibre
- Aide quotidienne (humaine ou technique)
- Activité professionnelle
- Autonomie dans les activités de la vie quotidienne (conduite, loisir ect.)

VOS DROITS

Vous disposez de droits d'accès et de rectification, d'opposition et d'un droit de réclamation à une autorité de contrôle (CNIL).

QUE DEVEZ-VOUS FAIRE ?

- Si vous ne vous opposez pas à l'utilisation des données de votre dossier médical, vous n'avez rien à faire.
- Si vous souhaitez vous opposer à l'utilisation de vos données pour ce projet, vous pouvez :
 - En parler directement au personnel du service que vous rencontrerez ou que vous avez rencontré au cours du bilan multidisciplinaire.
 - Contacter le responsable de l'étude ou le délégué à la protection des données (voir ci-dessous).

VOS CONTACTS

Pour exercer vos droits, ou pour toute question à ce sujet :

- Vous pouvez contacter le délégué à la protection des données du CHU Grenoble Alpes par mail (protection-donnees@chu-grenoble.fr) ou par téléphone au 04.76.76.82.02
- Une demande peut aussi être envoyée par courrier à CHU Grenoble Alpes, Délégué à la Protection des Données, CS 10217, 38043 GRENOBLE Cedex 09.
- Vous pouvez demander un formulaire d'opposition à l'accueil des services de MPR neurologique et de génétique clinique.

Si vous avez des questions sur l'étude, vous pouvez contacter le responsable ci-dessous :

Investigateur responsable de l'étude : Pr PERENNOU et Dr DIETERICH Coordonnées : CHU Grenoble Alpes, kdieterich@chu-grenoble.fr dperennou@chu-grenoble.fr CS10217 CHU, 38043 GRENOBLE Cedex 09	Organisme responsable du traitement des données : CHU Grenoble Alpes Coordonnées : CHU Grenoble Alpes - DR CI Pavillon Dauphiné – Rez-de-chaussée CS 10217 38043 GRENOBLE Cedex 9 Tel : 04 76 76 59 57
--	---



FORMULAIRE D'OPPOSITION

**Mercuri Analysis Contribution on Handicap evaluation in ArthrogypOsis,
a congenital Neuromuscular disease.
MACHAON**

A REMPLIR PAR LE PATIENT

Je soussigné(e) Madame/Monsieur (nom et prénom) demande à faire valoir mon droit d'opposition à l'utilisation des données à caractère personnel de mon dossier médical (ou du dossier de mon enfant) dans le cadre de cette recherche.

Date : ____ / ____ / ____

Signature :

Formulaire à renvoyer à :

Secrétariat de rééducation neurologique

Institut de Rééducation

Avenue de Kimberley


38130 ECHIROLLES

**CHU Grenoble Alpes, Délégué à la Protection des Données
CS 10217, 38043 GRENOBLE Cedex 09**

Ou à remettre au secrétariat des services médicaux qui transmettra

Ou en version scannée par email à protection-donnees@chu-grenoble.fr

Serment d'hippocrate



SERMENT D'HIPPOCRATE

En présence des Maîtres de cette Faculté, de mes chers condisciples et devant l'effigie d'HIPPOCRATE,

Je promets et je jure d'être fidèle aux lois de l'honneur et de la probité dans l'exercice de la Médecine.

Je donnerai mes soins gratuitement à l'indigent et n'exigerai jamais un salaire au dessus de mon travail. Je ne participerai à aucun partage clandestin d'honoraires.

Admis dans l'intimité des maisons, mes yeux n'y verront pas ce qui s'y passe ; ma langue taira les secrets qui me seront confiés et mon état ne servira pas à corrompre les mœurs, ni à favoriser le crime.

Je ne permettrai pas que des considérations de religion, de nation, de race, de parti ou de classe sociale viennent s'interposer entre mon devoir et mon patient.

Je garderai le respect absolu de la vie humaine.

Même sous la menace, je n'admettrai pas de faire usage de mes connaissances médicales contre les lois de l'humanité.

Respectueux et reconnaissant envers mes Maîtres, je rendrai à leurs enfants l'instruction que j'ai reçue de leurs pères.

Que les hommes m'accordent leur estime si je suis fidèle à mes promesses.

Que je sois couvert d'opprobre et méprisé de mes confrères si j'y manque.