



**HAL**  
open science

## ERAS wrong turn: each step costs carbon

Charles Moret, Diana Kabanova

► **To cite this version:**

Charles Moret, Diana Kabanova. ERAS wrong turn: each step costs carbon. Human health and pathology. 2022. dumas-04387299

**HAL Id: dumas-04387299**

**<https://dumas.ccsd.cnrs.fr/dumas-04387299>**

Submitted on 11 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## AVERTISSEMENT

Ce document est le fruit d'un long travail approuvé par le jury de soutenance.

La propriété intellectuelle du document reste entièrement celle du ou des auteurs. Les utilisateurs doivent respecter le droit d'auteur selon la législation en vigueur, et sont soumis aux règles habituelles du bon usage, comme pour les publications sur papier : respect des travaux originaux, citation, interdiction du pillage intellectuel, etc.

Il est mis à disposition de toute personne intéressée par l'intermédiaire de [l'archive ouverte DUMAS](#) (Dépôt Universitaire de Mémoires Après Soutenance).

Si vous désirez contacter son ou ses auteurs, nous vous invitons à consulter en ligne les annuaires de l'ordre des médecins, des pharmaciens et des sages-femmes.

Contact à la Bibliothèque universitaire de Médecine Pharmacie de Grenoble :

[bump-theses@univ-grenoble-alpes.fr](mailto:bump-theses@univ-grenoble-alpes.fr)

Année : 2022

**LA RAAC COMME NORME ÉCO-RESPONSABLE :  
CHAQUE ÉTAPE COÛTE DU CARBONE  
ERAS WRONG TURN: EACH STEP COSTS CARBON**

THÈSE  
POUR LE DIPLÔME D'ÉTAT DE DOCTEUR EN MÉDECINE  
SPÉCIALITÉ : ANESTHÉSIE - RÉANIMATION

Par M. **Charles MORET**

*[Données à caractère personnel]*

Et Mlle **Diana KABANOVA**

*[Données à caractère personnel]*

THÈSE SOUTENUE PUBLIQUEMENT À LA FACULTÉ DE MÉDECINE DE GRENOBLE

Le 22/06/2022

DEVANT LE JURY COMPOSÉ DE :

Président du jury :

M. le Pr Pierre ALBALADEJO (directeur de thèse)

Membres :

M. le Pr Jean-Luc FAUCHERON

M. le Pr Jean-Luc BOSSON

Mme le Dr Claire CHAPUIS

*L'UFR de Médecine de Grenoble n'entend donner aucune approbation ni improbation aux opinions émises dans les thèses ; ces opinions sont considérées comme propres à leurs auteurs.*

CORPS	NOM-PRENOM	Discipline universitaire
PU-PH	ALBALADEJO Pierre	Anesthésiologie-réanimation et médecine péri-opératoire
PU-PH	ARVIEUX-BARTHELEMY Catherine	Chirurgie viscérale et digestive
PU-PH	BAILLET Athan	Rhumatologie
PU-PH	BARONE-ROCHETTE Gilles	Cardiologie
PU-PH	BAYAT Sam	Physiologie
MCF Ass.MG	BENDAMENE Farouk	Médecine Générale
PU-PH	BENHAMOU Pierre-Yves	Endocrinologie, diabète et maladies métaboliques
PU-PH	BERGER François	Biologie cellulaire
MCU-PH	BETRY Cécile	Nutrition
MCU-PH	BIDART-COUTTON Marie	Biologie cellulaire
PU-PH	BIOULAC-ROGIER Stéphanie	Pédopsychiatrie ; addictologie
PU-PH	BLAISE Sophie	Chirurgie vasculaire ; médecine vasculaire
PR Ass. Méd.	BOILLOT Bernard	Urologie
MCU-PH	BOISSET Sandrine	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	BONAZ Bruno	Gastroentérologie ; hépatologie ; addictologie
PU-PH	BONNETERRE Vincent	Médecine et santé au travail
PU-PH	BOREL Anne-Laure	Nutrition
PU-PH	BOSSON Jean-Luc	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	BOTTARI Serge	Biologie cellulaire
PR Ass.MG	BOUCHAUD Jacques	Médecine Générale
PU-PH	BOUGEROL Thierry	Psychiatrie d'adultes
PU-PH	BOUILLET Laurence	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
MCU-PH	BOUSSAT Bastien	Epidémiologie, économie de la santé et prévention
PU-PH	BOUZAT Pierre	Anesthésiologie-réanimation et médecine péri-opératoire
PU-PH émérite	BRAMBILLA Christian	Pneumologie
PU-PH émérite	BRAMBILLA Elisabeth	Anatomie et cytologie pathologiques
MCU-PH	BRENIER-PINCHART Marie Pierre	Parasitologie et mycologie
PU-PH	BRICAULT Ivan	Radiologie et imagerie médicale
PU-PH	BRICHON Pierre-Yves	Chirurgie thoracique et cardiovasculaire
MCU-PH	BRIOT Raphaël	Thérapeutique-médecine de la douleur ; Addictologie
PU-PH émérite	CAHN Jean-Yves	Hématologie
PU-PH émérite	CARPENTIER Patrick	Chirurgie vasculaire, médecine vasculaire
PR Ass.MG	CARRILLO Yannick	Médecine Générale
MCU-PH	CASPAR Yvan	Bactériologie-virologie ; hygiène hospitalière
PU-PH	CESBRON Jean-Yves	Immunologie
PU-PH	CHABARDES Stephan	Neurochirurgie
PU-PH	CHABRE Olivier	Endocrinologie, diabète et maladies métaboliques
PU-PH	CHAFFANJON Philippe	Anatomie
MCF Ass.MG	CHAMBOREDON Benoît	Médecine Générale
PU-PH	CHARLES Julie	Dermato-vénéréologie
MCF Ass.MG	CHAUVET Marion	Médecine Générale

PU-PH	CHAVANON Olivier	Chirurgie thoracique et cardiovasculaire
PU-PH	CHIQUET Christophe	Ophthalmologie
PU-PH	CHIRICA Mircea	Chirurgie viscérale et digestive
PU-PH	CINQUIN Philippe	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	CLAVARINO Giovanna	Immunologie
MCU-PH	CLIN CHERPEC Rita	Nutrition
PU-PH	COHEN Olivier	Histologie, embryologie et cytogénétique
PU-PH	COSTENTIN Charlotte	Gastroentérologie ; hépatologie ; addictologie
PU-PH	COURVOISIER Aurélien	Chirurgie infantile
PU-PH	COUTTON Charles	Génétique
PU-PH	COUTURIER Pascal	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PU-PH	CRACOWSKI Jean-Luc	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PU-PH	DEBATY Guillaume	Médecine d'Urgence
PU-PH	DEBILLON Thierry	Pédiatrie
PU-PH	DECAENS Thomas	Gastroentérologie ; hépatologie ; addictologie
PR Ass. Méd.	DEFAYE Pascal	Cardiologie
PU-PH	DEGANO Bruno	Pneumologie ; addictologie
PU-PH	DEMATTEIS Maurice	Pharmacologie fondamentale ; pharmacologie clinique ; addictologie
PU-PH émérite	DEMONGEOT Jacques	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	DERANSART Colin	Physiologie
PU-PH	DESCOTES Jean-Luc	Urologie
PU-PH	DETANTE Olivier	Neurologie
MCU-PH	DIETERICH Klaus	Génétique
MCU-PH	DOUTRELEAU Stéphane	Physiologie
PU-PH	EPAULARD Olivier	Maladies infectieuses ; Maladies tropicales
PU-PH	ESTEVE François	Biophysique et médecine nucléaire
MCU-PH	EYSSERIC Hélène	Médecine légale et droit de la santé
PU-PH émérite	FAGRET Daniel	Biophysique et médecine nucléaire
PU-PH	FAUCHERON Jean-Luc	Chirurgie viscérale et digestive
MCU-PH	FAURE Julien	Biochimie et biologie moléculaire
PU-PH	FERRETTI Gilbert	Radiologie et imagerie médicale
PU-PH	FONTAINE Éric	Nutrition
PU-PH	FRANCOIS Patrice	Epidémiologie, économie de la santé et prévention
MCU-MG	GABOREAU Yoann	Médecine Générale
PU-PH	GARBAN Frédéric	Hématologie ; Transfusion
PU-PH	GAUDIN Philippe	Rhumatologie
PU-PH	GAVAZZI Gaétan	Médecine interne ; gériatrie et biologie du vieillissement ; addictologie
PU-PH	GAY Emmanuel	Neurochirurgie
MCU-PH	GIAI Joris	Biostatistiques, informatique médicale et technologies de communication
MCU-PH	GILLOIS Pierre	Biostatistiques, informatique médicale et technologies de communication
PU-PH	GIOT Jean-Philippe	Chirurgie plastique, reconstructrice et esthétique ; Brûlologie
MCU-PH	GRAND Sylvie	Radiologie et imagerie médicale
PU-PH émérite	GRIFFET Jacques	Chirurgie infantile
PU-PH	HAINAUT Pierre	Biochimie et biologie moléculaire
PU-PH émérite	HALIMI Serge	Nutrition
PU-PH	HENNEBICQ Sylviane	Biologie et médecine du développement et de la reproduction ; gynécologie médicale
PU-PH	HOFFMANN Pascale	Gynécologie-obstétrique ; gynécologie médicale

PU-PH émérite	HOMMEL Marc	Neurologie
PU-MG	IMBERT Patrick	Médecine Générale
PU-PH émérite	JOUK Pierre-Simon	Génétique
PU-PH	KAHANE Philippe	Physiologie
MCU-PH	KASTLER Adrian	Radiologie et imagerie médicale
PU-PH	KRAINIK Alexandre	Radiologie et imagerie médicale
PU-PH	LABARERE José	Epidémiologie, économie de la santé et prévention
MCU-PH	LABLANCHE Sandrine	Endocrinologie, diabète et maladies métaboliques
PU-PH	LANDELLE Caroline	Bactériologie – virologie ; Hygiène hospitalière
PU-PH	LANTUEJOU Sylvie	Anatomie et cytologie pathologiques
PR Ass. Méd.	LARAMAS Mathieu	Cancérologie ; radiothérapie
MCU-PH	LARDY Bernard	Biochimie et biologie moléculaire
MCU-PH	LE GOUELLEC LE PISSART Audrey	Biochimie et biologie moléculaire
PU-PH	LECCIA Marie-Thérèse	Dermato-vénérologie
PR Ass.MG	LEDOUX Jean-Nicolas	Médecine Générale
PU-PH émérite	LETOUBLON Christian	Chirurgie viscérale et digestive
PU-PH	LEVY Patrick	Physiologie
PU-PH	LONG Jean-Alexandre	Urologie
MCU-PH	LUPO Julien	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	MAITRE Anne	Médecine et santé au travail
MCU-PH	MARLU Raphaël	Hématologie ; Transfusion
PR Ass. Méd.	MATHIEU Nicolas	Gastroentérologie ; hépatologie ; addictologie
MCU-PH	MAUBON Danièle	Parasitologie et mycologie
PU-PH	MAURIN Max	Bactériologie-virologie ; Hygiène hospitalière
MCU-PH	MC LEER Anne	Histologie, embryologie et cytogénétique
PR Ass. Méd	MICHY Thierry	Gynécologie-obstétrique
MCU-PH	MONDET Julie	Histologie, embryologie et cytogénétique
PU-PH	MORAND Patrice	Bactériologie-virologie ; Hygiène hospitalière
PU-PH	MOREAU-GAUDRY Alexandre	Biostatistiques, informatique médicale et technologies de communication
PU-PH	MORO Elena	Neurologie
PU-PH	MORO-SIBILOT Denis	Pneumologie ; addictologie
MCU-PH	MORTAMET Guillaume	Pédiatrie
PU-PH	MOUSSEAU Mireille	Cancérologie ; radiothérapie
PU-PH émérite	MOUTET François	Chirurgie plastique, reconstructrice et esthétique ; brûlologie
MCF Ass.MG	ODDOU Christel	Médecine Générale
PR Ass. Méd.	ORMEZZANO Olivier	Cardiologie
MCU-PH	PACLET Marie-Hélène	Biochimie et biologie moléculaire
PU-PH	PAILHE Régis	Chirurgie orthopédique et traumatologie
PU-PH	PALOMBI Olivier	Anatomie
PU-PH	PARK Sophie	Hématologie ; Transfusion
PR Ass.MG	PAUMIER-DESBRIERES Françoise	Médecine Générale
PU-PH	PAYEN DE LA GARANDERIE Jean-François	Anesthésiologie-réanimation et médecine péri-opératoire
MCU-PH	PAYSANT François	Médecine légale et droit de la santé
MCU-PH	PELLETIER Laurent	Biologie cellulaire
PU-PH	PELLOUX Hervé	Parasitologie et mycologie
PU-PH	PEPIN Jean-Louis	Physiologie
PU-PH	PERARD DUMESTRE Chantal	Immunologie

PU-PH	<b>PERENNOU Dominique</b>	<b>Médecine physique et de réadaptation</b>
PU-PH	<b>PERNOD Gilles</b>	Chirurgie vasculaire ; <b>Médecine vasculaire</b>
MCF	<b>PINSAULT Nicolas</b>	Sciences de la rééducation et de réadaptation
PU-PH	<b>PIOLAT Christian</b>	<b>Chirurgie infantile</b>
PU-PH	<b>PISON Christophe</b>	<b>Pneumologie ; Addictologie</b>
PU-PH	<b>PLANTAZ Dominique</b>	<b>Pédiatrie</b>
PU-PH	<b>POIGNARD Pascal</b>	<b>Bactériologie-virologie ; Hygiène hospitalière</b>
PU-PH émérite	<b>POLACK Benoît</b>	<b>Hématologie ; Transfusion</b>
PU-PH	<b>POLOSAN Mircea</b>	<b>Psychiatrie d'adultes ; Addictologie</b>
PU-PH émérite	<b>RAMBEAUD Jean-Jacques</b>	<b>Urologie</b>
PU-PH	<b>RAY Pierre</b>	<b>Biologie et médecine du développement et de la reproduction ; gynécologie médicale</b>
PR Ass. Méd.	<b>RECHE Fabian</b>	<b>Chirurgie viscérale et digestive</b>
MCU-PH	<b>RENDU John</b>	<b>Biochimie et biologie moléculaire</b>
MCU-PH émérite	<b>RIALLE Vincent</b>	<b>Biostatistiques, informatique médicale et technologies de communication</b>
PU-PH	<b>RIETHMULLER Didier</b>	<b>Gynécologie-obstétrique ; gynécologie médicale</b>
PU-PH	<b>RIGHINI Christian</b>	<b>Oto-rhino-laryngologie</b>
PU-PH émérite	<b>ROMANET Jean Paul</b>	<b>Ophthalmologie</b>
PU-PH	<b>ROSTAING Lionel</b>	<b>Néphrologie</b>
PU-PH	<b>ROUSTIT Matthieu</b>	<b>Pharmacologie fondamentale ; pharmacologie clinique ; addictologie</b>
MCU-PH	<b>ROUX-BUISSON Nathalie</b>	<b>Biochimie et biologie moléculaire</b>
PR Ass.MG	<b>ROYER DE VERICOURT Guillaume</b>	<b>Médecine Générale</b>
PU-PH émérite	<b>SARAGAGLIA Dominique</b>	<b>Chirurgie orthopédique et traumatologie</b>
MCU-PH	<b>SATRE Véronique</b>	<b>Génétique</b>
PU-PH	<b>SAUDOU Frédéric</b>	<b>Biologie cellulaire</b>
PU-PH	<b>SCHMERBER Sébastien</b>	<b>Oto-rhino-laryngologie</b>
PU-PH	<b>SCHWEBEL Carole</b>	<b>Médecine intensive-réanimation</b>
PU-PH	<b>SCOLAN Virginie</b>	<b>Médecine légale et droit de la santé</b>
PU-PH	<b>SEIGNEURIN Arnaud</b>	<b>Epidémiologie, économie de la santé et prévention</b>
PU-PH	<b>SPEAR Rafaëlle</b>	<b>Chirurgie vasculaire ; médecine vasculaire</b>
PU-PH émérite	<b>STAHL Jean-Paul</b>	<b>Maladies infectieuses ; Maladies tropicales</b>
PU-PH	<b>STANKE Françoise</b>	<b>Pharmacologie fondamentale ; pharmacologie clinique ; addictologie</b>
MCU-PH	<b>STASIA Marie-José</b>	<b>Biochimie et biologie moléculaire</b>
PU-PH	<b>STURM Nathalie</b>	<b>Anatomie et cytologie pathologiques</b>
PU-PH	<b>TAMISIER Renaud</b>	<b>Physiologie</b>
PU-PH	<b>TERZI Nicolas</b>	<b>Médecine intensive-réanimation</b>
PU-PH	<b>THEVENON Julien</b>	<b>Génétique</b>
PU-PH	<b>TOFFART Anne-Claire</b>	<b>Pneumologie ; Addictologie</b>
PU-PH	<b>TONETTI Jérôme</b>	<b>Chirurgie orthopédique et traumatologie</b>
PU-PH	<b>TOUSSAINT Bertrand</b>	<b>Biochimie et biologie moléculaire</b>
PU-PH	<b>VALMARY-DEGANO Séverine</b>	<b>Anatomie et cytologie pathologiques</b>
PU-PH	<b>VANZETTO Gérald</b>	<b>Cardiologie</b>
PU-PH	<b>VUILLEZ Jean-Philippe</b>	<b>Biophysique et médecine nucléaire</b>
PU-PH	<b>WEIL Georges</b>	<b>Epidémiologie, économie de la santé et prévention</b>
PU-PH	<b>ZAQUI Philippe</b>	<b>Néphrologie</b>
PU-PH émérite	<b>ZARSKI Jean-Pierre</b>	<b>Gastroentérologie ; hépatologie ; addictologie</b>

# TABLE DES MATIÈRES

<b>RESUME</b>	<b>14</b>
<b>ABSTRACT</b>	<b>16</b>
<b>INTRODUCTION</b>	<b>18</b>
<b>METHODS</b>	<b>19</b>
<b>RESULTS</b>	<b>20</b>
A.    BEFORE SURGERY	20
1. <i>Information and counseling</i>	20
2. <i>Preoperative fasting</i>	21
3. <i>Preoperative evaluation and optimization</i>	21
4. <i>Patient blood management</i>	24
5. <i>Prehabilitation</i>	25
6. <i>Other preoperative issues in colorectal surgery</i>	26
7. <i>Other preoperative issues in orthopedic surgery</i>	28
B.    DURING SURGERY	29
1. <i>Goal directed fluid therapy</i>	29
2. <i>Corticotherapy</i>	29
3. <i>Target thermic control</i>	30
4. <i>Prophylactic antibiotic treatment</i>	30
5. <i>Choice of surgical technique</i>	31
6. <i>Type of anesthesia</i>	32
7. <i>Other intraoperative issues in orthopedic surgery</i>	32
C.    AFTER SURGERY	33
1. <i>Multimodal analgesia</i>	33
2. <i>Venous thromboembolism prevention</i>	33
3. <i>Drains</i>	34
4. <i>Early mobilization and feeding</i>	35
5. <i>Other postoperative issues in colorectal surgery</i>	36
6. <i>Other postoperative issues in orthopedic surgery</i>	37
7. <i>Post discharge management</i>	37
D.    ADDITIONAL ISSUES: ERAS AND OPTIMIZED HOSPITAL STAY: TIME IS CARBON	39
<b>DISCUSSION</b>	<b>41</b>
<b>CONCLUSION</b>	<b>43</b>
<b>FIGURES</b>	<b>44</b>
<b>APPENDIX</b>	<b>46</b>
<b>LEXICON</b>	<b>54</b>
<b>BIBLIOGRAPHY</b>	<b>55</b>



## REMERCIEMENTS COMMUNS

Nous tenons d'une voix commune à remercier les membres du jury:

Monsieur le **Professeur ALBALADEJO**, pour vos idées, votre ouverture d'esprit et pour vos conseils, précieux et justes. Pour nous avoir soutenu sur ce sujet de thèse actuel et innovant. Pour vos enseignements toujours pertinents et instructifs.

Monsieur le **Professeur FAUCHERON**, pour l'honneur qu'il nous fait d'avoir accepté d'être présent ce jour. Pour votre expertise clinique et votre rigueur dans l'exercice chirurgical.

Monsieur le **Professeur BOSSON**, pour votre disponibilité et pour l'intérêt que vous portez à notre thématique de thèse.

Madame la **Docteur CHAPUIS**, pour votre motivation dans la participation à ce jury et pour votre action au sein du PAR.

## REMERCIEMENTS de la part de Diana

Je remercie **Charles Moret**, mon co-thésard et co-interne pour le travail accompli, pour sa patience et ses remarques pertinentes, son humeur toujours égale, même dans les moments de doute.

Je remercie évidemment **Christine Buisson**, qui est un pilier du PAR, et sans qui l'internat aurait été vraiment plus compliqué.

Je remercie **Vincent Perignon** pour son soutien sans aucune faille au quotidien, et merci à **Oscar** pour le soutien émotionnel.

Merci à **Margarita** et **Youri Kabanov** pour m'avoir accompagné jusqu'à la thèse, sans qui rien n'aurait été possible.

Je remercie mes co internes de la promotion AR 2018 : ma colocataire **Marjolaine** pour son enthousiasme et ses actions coup de poing, **Florian Ruping** qui a toujours le mot pour rire, **Théophile** pour son sens de la solidarité et de l'initiative, **Pauline** merci pour ta gentillesse et ton empathie, merci à **Stéphane** pour sa festivité, merci à **Mariano** pour son calme en toute circonstance, merci à **Benjamin** pour son énergie et sa fiabilité, et merci à **Antoine** et **Victor** que j'aurai aimé mieux connaître.

Je remercie mes collègues de la RPC qui m'ont fait découvrir la réanimation : mes co internes **Paula, Junélie, Zoé, Marvin, Rémi, Cloé, Camille** et **Delphine**, les équipes d'infirmiers, AS, ASH, les médecins qui ont été là pour m'accompagner lorsque je savais tout juste ce qu'était un pousse-seringue. Merci à **Sarah** et **Quentin**, internes seniorisés d'avoir été exemple à suivre. Merci à **Cédric Préaubert** et **Manon Gabin** pour ces gardes fort instructives lors de votre seniorisation.

Je remercie les médecins et IADE de la maternité qui m'ont appris à poser mes premières péridurales. Merci au **Dr Ana Rogé** qui m'a sensibilisé à la communication thérapeutique, merci à Julien **Ondel** et **Julie Blachère**, qui m'ont fait endormir ma première patiente.

Merci aux équipes de l'hôpital sud : les MAR et les IADE grâce à qui j'ai pu acquérir des compétences pointues en ALR, et qui ont su créer une ambiance particulièrement agréable et bienveillante.

Merci à **Roch Mader**, rencontré à l'hôpital Sud, et **Emmanuelle Orset**, avec qui les aventures se sont poursuivies en dehors de l'hôpital, et qui seront je l'espère mes amis et collègues pendant encore longtemps.

Un grand merci aux équipes d'anesthésie pédiatrique qui m'ont enseigné la rigueur, la sécurité et m'ont fait découvrir le plaisir et la joie de travailler avec des enfants. Merci à **Chantal, Kelly, Giuseppe, Jean-Noel** et tous les autres pour vos enseignements. Une pensée particulière pour **Jeanne** qui a fait preuve de patience, et m'a fait persévérer et progresser en anesthésie.

Je tiens à remercier les équipes d'anesthésie d'orthopédie et de chirurgie de la main pour m'avoir accompagné vers l'autonomisation et la prise de confiance, merci d'avoir continué à me former pendant ce stage de début de covid. Merci aux MAR, merci **Marie-Pierre, Yves, Daniel** pour votre enthousiasme et votre bienveillance.

Je remercie les équipes d'anesthésie et de chirurgie du CH de Sallanches qui m'ont fait découvrir l'organisation des hôpitaux périphériques, dans un cadre incroyable et des équipes soudées et conviviales. Merci à **Bernard** d'avoir été là pour me faire découvrir l'histoire de l'alpinisme et du secours en montagne, ainsi que de m'avoir aidé à traverser cette période de Covid avec plus d'empathie et de tolérance. Merci à **Sophie**, pour son professionnalisme, son humour, ses connaissances linguistiques et pour tout ce temps incroyable passé en Haute Savoie. Merci à **Séverine, Camille, Sandrine, Marie-Pierre, Claire, Isabelle**, ainsi qu'à mes collègues et amies de la salle de réveil. Merci à **Stéphanie, Aurélien, Lucas, Gianluca, Céline** et tous les autres que j'ai eu la chance de rencontrer pendant ce semestre.

Je remercie chaleureusement l'équipe de la Réanimation CVT, qui malgré la période difficile de covid, a su garder des équipes motivées, agréables, pleines d'optimisme et de bonne humeur. Merci aux médecins réanimateurs qui ont été des exemples tant d'un point de vue humain que médical, et qui ont su être à l'écoute. Merci **Michel, Daniel, Géraldine, Lucie, Amélie, Romain, Laura, Emmanuelle, Myriam, Damien, Sylvaine, Timothé**, ainsi qu'à toutes les équipes d'infirmières, aide soignantes et ASH. Merci à mes co-internes pour ce semestre enrichissant, et merci **Adrien** pour ce semestre à tes côtés.

Je tiens à remercier la Réanimation de Chambéry pour ce stage estival en compagnie de co-internes à qui je tiens énormément, **Marine, Florian, Léa, Gauthier** et **Nicolas Segond**. Merci à tous les médecins de ce service, pour m'avoir beaucoup appris, merci à Michel pour ses précieux conseils, merci **Clémence** et **Catherine** pour mes premières gardes séniorisées, merci **Margot, Vincent, Julien et Julien, Stéphane, David** et **David, Jean-Marc** et tous autres. Un merci particulier pour toutes les infirmières et aide soignantes qui ont été là pendant ces journées et ces nuits, merci **Vanessa, Laetitia, Romane, Angélique, Pierre Louis, Delphine, Cécile, Laure, David, Marianne, Angélique, Estelle**, et les nombreux autres qui seraient trop longs à nommer ici. Merci à **Charlotte** et **Anita** à qui je pense très souvent.

Merci aux équipes du BU Déchocage, merci pour ces 3 mois intenses à vos côtés. Merci à mes co-internes, **Florian, Alexandre, Oussama et Anaïs** ; merci aux infirmières anesthésistes, merci à **Pauline, Charlotte, Alexandra, Alexia, Julien, Aline, Amandine, Baptiste** et tous les

autres que je ne peux pas nommer ici. Merci aux MAR, merci aux chirurgiens et IBODE qui font tourner ce bloc exigeant, jour et nuit.

Merci à la clinique Belledonne pour m'avoir accueilli pour ce stage, trop court à mon goût, merci de m'avoir fait progresser et permis de consolider mes techniques d'Anesthésie. Merci **Christine, Isabelle, Katharina, Pierre, Philippe, Julie, Emmanuel, Vincent, Stéphane**, merci à tous ces formidables IADES et bien sûr merci à tous ces chirurgiens, avec qui il a été très agréable de travailler. Merci **Florian**, pour m'avoir guidé pendant l'internat et pour m'avoir épaulé pendant les moments heureux et difficiles. Merci **Julie, Estelle, Valérian, Isabelle, Pierre et Océane** pour toutes ces belles journées passées à vos côtés.

Merci à **Laurie** et **Olivier**, mes chers amis, merci à **Franck** mon entraîneur et ami.

Merci à tous mes amis sportifs que j'ai eu la chance de rencontrer en arrivant à Grenoble, merci à **Julie** et **Edouard, Anne-Lyse** et **Simon, Ania** et **Alexis**.

Merci à mes amis bisontins, merci à mon club d'athlétisme le CMI, merci à **Cécilia** et **Ibrahim**.

Merci à **Julien**, qui m'a conseillé tout au long de mes études, du lycée jusqu'à la thèse.

Merci à **Matthieu** et **Matthieu**, merci à mes amis d'Arches, mon lieu sûr d'adoption.

## REMERCIEMENTS de la part de Charles

Aux médecins et aux personnels paramédicaux qui ont participé à ma formation,

A la **RNC**, pour leur accueil, pour leur bienveillance après ces premières gardes de chef, et pour cette équipe Buvette toujours plus disponible lorsqu'il s'agit de se réunir !

Au **bloc de Chambéry et sa maternité**, une très belle expérience de la vie en périphérie et une équipe bienveillante avec qui c'était un plaisir d'évoluer et de progresser.

Au **bloc DIGIDUNE**, pour mon premier semestre d'anesthésie qui n'aura fait que confirmer que ce choix d'internat était le bon.

Au **bloc de pédiatrie**, pour la découverte qu'a été celle de la pédiatrie. Et un remerciement particulier à **Kelly** pour sa relecture attentive et ses conseils bienveillants dans l'élaboration de cette thèse.

Aux **blocs d'orthopédie Nord et Sud**, qui m'auront aidé à développer un réel intérêt pour l'anesthésie loco-régionale et pour la qualité de leur exercice de l'anesthésie au quotidien.

A **l'URCVT**, pour leur soutien et gestion d'équipe dans les périodes de Covid et pour leur pédagogie au quotidien.

A la **Réanimation Médicale**, pour son enseignement et son accompagnement bienveillant durant mon premier semestre d'internat.

A l'ensemble du **PAR**, pour la qualité de son enseignement universitaire, à **Christine** pour sa réactivité et sa bienveillance auprès des internes. Aux **infirmiers de réanimation** et aux **IADES**, sans qui ces années d'apprentissage n'auraient pas été aussi enrichissantes.

A ma famille,

A ma **maman**, pour son soutien inconditionnel, dans la vie personnelle et durant mes études.

A mon **papa**, qui aurait été fier de l'aboutissement de ces longues années d'études.

A **Marlène** et **Nicolas**, car la famille compte, même à distance.

A **Pierre**, pour ces parties de pétanques et ces discussions toujours intéressantes du quotidien ; à **Camille** et **Clément**, quelle bonne surprise ce fut de vous rencontrer.

A **Léa**, on se suit depuis le début de ces études et nous en voyons finalement le bout. Avoir passé ces années ensemble restera une richesse, et que cela dure pour longtemps.

A **Catherine** et **Éric**, pour le plaisir que représentent ces tablées du dimanche.

A **Michel**, pour ces sorties en randonnées l'hiver, ces tours de skis nautiques, et ces coups de mains bricolage.

A **Valentin**, maintenant que Paris est ton fief, nous attendons les invitations au Ritz.

A mes amis,

Aux **LGC**, sans conteste la meilleure bande d'amis, des bons vivants à l'évidence, et qui savent rire de chaque instant ! On se change très vite les idées en leur compagnie, et ce sont finalement ces moments-là qui rendent heureux.

A l'**Animalerie**, que de découvertes dans cette team de VSS ! Que ce soit dans le Beaujo, dans le Sud-Ouest ou jusqu'au Maroc, votre convivialité rend chaque événement unique !

A la **CovidFreeBand**, la crème des Grelous d'adoptions sans qui ces années d'internats n'auraient pas été les mêmes !

A la **RBB**, un semestre à vos côtés aura suffi pour que l'on rende réguliers ces arrosages de RMM revisités.

Aux **Lyonnais de l'externat**, même éparpillés dans toute la France, les limites de la fête sont repoussées dès que l'on se retrouve, que rien ne change !

A la promo de **DESAR**, avec qui on aura partagé chaque joie de ces années au bloc et en réa, chaque midi à l'internat, et tant de moments précieux. Et à **Diana** en particulier, pour ta bonne humeur quotidienne. Mener cette thèse à deux n'était pas une évidence pour nous, mais c'est finalement avec fierté que l'on peut se féliciter d'en être venus à bout !

**Marie** et **Ori**, peut-être les plus belles trouvailles de Pologne qui auront rendu si riches cet Erasmus ! C'est peut-être grâce à vous que j'ai poursuivi ma route chez vous à Grenoble...

Au **personnel de l'internat**, pour leur accueil inégalable depuis tant d'années, pour leur patience les lendemains de soirées difficiles, pour leur sens de la fête qui ne faiblira donc jamais !

Et bien sûr à ceux qui ont toujours été présents, qu'ils viennent de Bourg-en-Bresse, Saint-Maurice de Gourdans, du Jura, de la Réunion ou de la Motte-de-Galaure. Votre présence à chaque moment important de ma vie a participé à ce que je suis aujourd'hui, et je vous en remercie.

### “ Au moment d’être admis(e) à exercer la médecine, je promets et je jure d’être fidèle aux lois de l’honneur et de la probité.

Mon premier souci sera de rétablir, de préserver ou de promouvoir la santé dans tous ses éléments, physiques et mentaux, individuels et sociaux.

Je respecterai toutes les personnes, leur autonomie et leur volonté, sans aucune discrimination selon leur état ou leurs convictions. J'interviendrai pour les protéger si elles sont affaiblies, vulnérables ou menacées dans leur intégrité ou leur dignité. Même sous la contrainte, je ne ferai pas usage de mes connaissances contre les lois de l'humanité.

J'informerai les patients des décisions envisagées, de leurs raisons et de leurs conséquences.

Je ne tromperai jamais leur confiance et n'exploiterai pas le pouvoir hérité des circonstances pour forcer les consciences.

Je donnerai mes soins à l'indigent et à quiconque me les demandera. Je ne me laisserai pas influencer par la soif du gain ou la recherche de la gloire.

Admis(e) dans l'intimité des personnes, je tairai les secrets qui me seront confiés. Reçu(e) à l'intérieur des maisons, je respecterai les secrets des foyers et ma conduite ne servira pas à corrompre les mœurs.

Je ferai tout pour soulager les souffrances. Je ne prolongerai pas abusivement les agonies. Je ne provoquerai jamais la mort délibérément.

Je préserverai l'indépendance nécessaire à l'accomplissement de ma mission. Je n'entreprendrai rien qui dépasse mes compétences. Je les entretiendrai et les perfectionnerai pour assurer au mieux les services qui me seront demandés.

J'apporterai mon aide à mes confrères ainsi qu'à leurs familles dans l'adversité.

Que les hommes et mes confrères m'accordent leur estime si je suis fidèle à mes promesses; que je sois déshonoré(e) et méprisé(e) si j'y manque. ”



## RESUME

### *Objectif*

Déterminer l'empreinte carbone de la déviation à un protocole de RAAC après une chirurgie orthopédique lourde du membre inférieur type PTH/PTG ou après une chirurgie colorectale.

### *Méthode*

Notre revue narrative a été construite à partir d'articles référencés dans la base PubMed. Nos recherches ont été basées sur l'impact environnemental d'un protocole de RAAC dans le cadre de la chirurgie colorectale et d'arthroplastie de hanche ou de genou. Chaque point des recommandations françaises et internationales a été étudié. Pour nos recherches, nous avons utilisé les termes "empreinte carbone", "durabilité", "coût énergétique", "empreinte environnementale", "évaluation du cycle de vie" ET un mot clef de chaque sous partie des recommandations RAAC.

### *Résultats*

Durant la période péri-opératoire, les écarts à un protocole de RAAC, dans le cadre d'une chirurgie colorectale ou d'une arthroplastie lourde du membre inférieur, sont très coûteux en équivalent carbone. Le surcoût est estimé à environ 2500 Kg CO<sub>2</sub>/patient pour une chirurgie colorectale. La part la plus importante se produit durant la période peropératoire. Le surcoût est estimé à environ 1400 Kg CO<sub>2</sub>/patient dans le cadre d'une chirurgie orthopédique majeure. La part la plus importante se produit durant la période postopératoire. Ceci est un argument supplémentaire à l'application d'un protocole RAAC lorsque celui-ci est recommandé. Il existe des leviers supplémentaires, en dehors des protocoles de RAAC, pour réduire notre empreinte carbone dans notre pratique de l'anesthésie. Celle-ci devra probablement s'adapter à l'urgence du réchauffement climatique.



### *Conclusion*

Le non-respect des recommandations RAAC est très coûteux en termes de carbone. Notre étude montre un argument supplémentaire de la nécessité de mettre en place des parcours RAAC pour le patient, qui devrait être considéré comme la norme éco responsable d'une prise en charge chirurgicale. Des considérations plus globales et organisationnelles sont également à mettre en place pour réduire les émissions de CO2 dans le secteur de la santé.

## ABSTRACT

### *Objective*

The aim of this study is to evaluate the carbon footprint of deviations from ERAS pathway applied to a joint replacement surgery and to a colorectal surgery.

### *Design, settings*

This narrative review was designed to collect articles related to the environmental impact of deviations from the ERAS protocol in colorectal and joint replacement surgery. We decided to deal with each point of interest included in recent international and French recommendations. We based our search on the data base PubMed by including relevant references. Our search was oriented on available articles with the terms “carbon footprint”, “sustainability”, “ecology”, “energy cost”, “environmental footprint”, “life cycle assessment” or “climate” AND a keyword from each subpart of the ERAS recommendation.

### *Results*

The non-compliance to the ERAS protocol is costly in terms of carbon footprint in almost all the steps in perioperative time. The balance between benefits and costs in terms of the carbon footprint is strongly positive. The total cost for a colorectal surgery (from the admission to the hospital leave) is about 2 500 KgCO<sub>2</sub>e/patient, the main part during the intraoperative period. In a joint replacement surgery, the total cost is about 1 400 KgCO<sub>2</sub>e/patient, the main part during the postoperative period.

## *Conclusion*

Regarding ecological balance, our study shows an additional argument to follow the current ERAS guidelines for the colorectal and joint replacement surgery. Additional actions beyond the ERAS pathway are strongly efficient on carbon savings.

## INTRODUCTION

After the Brundtland Commission of 1987, the concept of sustainable development implicates public institutions, and, in particular, the health care system. It represents between 3% to 8% of the annual CO<sub>2</sub> production for European countries [1, 2], about 2.7% for China [3] and up to 9.8 % for the USA [2]. French authorities recommend that health care institutions should meet criteria regarding the environment and waste recycling. Since 2010, private and public institutions have to publish their annual carbon emissions [4].

Climate change is one of the major issues of modern times. Anthropogenic pollution is slowly becoming part of our daily considerations, and is challenging our professional habits. Burrell even proposed to extend the concept of the medical ethic “first do not harm” to the environment [5] since it impacts our patients. We may find ourselves in the paradox of providing care with a system that is itself harmful for health [6].

In this context, we aimed to explore the role of an Enhanced Recovery After Surgery (ERAS) pathway in the context of eco sustainability. ERAS refers to evidence-based multidisciplinary care pathways for a surgical specialty [7, 8, 9, 10] that aim to reduce surgical stress response, optimize physiologic function, and facilitate recovery.. The significant clinical benefits for the patient and the cost-effectiveness of ERAS are now well established [8, 9, 11, 12]. We focus our analysis on the environmental aspect with the aim of this article being to evaluate the cumulative carbon impact of deviations from the ERAS pathway applied to colorectal surgery and major joint replacements.

## METHODS

This narrative literature review was designed to collect articles related to the environmental impact of deviations from ERAS protocols in colorectal and major joint replacement surgery.

Recent international and French recommendations [7, 8, 9, 10] served as reference documents for the PubMed literature search performed. There was no limit in the publication date, since the ecological cost of healthcare is a relatively recent concern.

Our research was based on available articles including the terms “carbon footprint”, “sustainability”, “ecology”, “energy cost”, “environmental footprint”, “life cycle assessment” or “climate” AND a keyword from each subpart of the ERAS recommendations, for example “chronic disease optimization” or “target thermic control”. In the case of data pertaining to a keyword that was not specific to ERAS, the selection of articles was narrowed by adding a third inter-related term like “anesthesia”, or “operating room”.

The focus was on lower-limb joint replacement surgery and colorectal surgery in adults. When the publications available concerned other types of surgery, it is mentioned in the text.

For each item developed in ERAS recommendations, direct data of carbon cost were searched. If not available, carbon cost was approximated indirectly.

## **RESULTS**

Results are presented for the major orthopedic surgery and colorectal surgery during the pre operative, intraoperative, and postoperative periods.

### **A. Before Surgery**

#### **1. Information and counseling**

Communication has a certain cost, related not only to transport and human resources necessary for consultation, but also to internet use.

An important part of the ERAS pathway is the quality of patient information and education, and furthermore, patient adherence to the protocol. One option is to promote targeted information by text messages or email (about 0.3 g of CO<sub>2</sub> per email [13]), or even by individualized calls to patients at considerable risk of cancellation. Such patients could be identified during the anesthesiologist's appointment [14]. However, it remains difficult to assess the carbon impact of the quality of patient information and education.

## **2. Preoperative Fasting**

Current preoperative fasting guidelines are not associated with increased anesthesia risk or any modification of length-of-stay (LOS) after surgery but do provide more patient comfort [7]; The carbon balance of this measure is neutral.

## **3. Preop evaluation and optimization**

As a general principle, prevention of any postoperative complications by thorough and evidence-based preoperative evaluation will decrease the costs (financial and environmental) related to a surgical procedure. The ERAS Society Guidelines strongly recommend optimizing medical conditions prior to surgery. However, the cost of invasive investigations is often significant. For example, interventional radiology or coronarography produces 243 000 g CO<sub>2</sub>e/procedure and 8.28 kg of waste [15]. Each kilogram is equal to 3000 g of CO<sub>2</sub>, thus 24 840 g of CO<sub>2</sub> secondary to waste must be added to the global cost (including energy, medical devices and human resources) [16].

Diabetes is one of the chronic medical diseases that needs to be diagnosed and treated. The principal perioperative complication of chronic hyperglycemia is susceptibility to infections. The risk is multiplied by 5.29, if the preoperative HbA<sub>1c</sub> is higher than 7.8% [32, 33, 34]. For each 1% increase in HbA<sub>1c</sub> above 7%, the risk of infectious complications increases by 40% [35]. Every 1% increase in HbA<sub>1c</sub> above normal adds 6 480 g CO<sub>2</sub> per patient, considering the risk for patients without risk factors of 0.72% [35] (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

In terms of substance use and addiction, tobacco cessation is a major challenge because of its role in perioperative morbidity and mortality. The most effective interventions are intensive interventions using a nicotine substitute such as varenicline that start at least 4 weeks before surgery [17, 18, 19, 20]. Stopping cigarette smoking could help avoid 2 millions of complications in the USA per year.

The proportion of surgical patients who smoke is estimated to be 25.9% [21]. Smokers have a two-fold risk of surgical site complications. The risk of pulmonary complications is multiplied by a factor up to 4 [22]. The overall risk of complications for active smokers is between 31% and 45%, and LOS is increased by up to 3 days [21].

According to the French National Authority for Health, the odds ratio (OR) for complications does not return to zero after stopping tobacco. Interventions for smoking cessation implemented 6 to 8 weeks before hip or knee arthroplasty can however, significantly reduce up to 83% [23] the complication rate.

Stopping smoking represents a reduction of carbon costs of 140 063 g of CO<sub>2</sub> due to lower risk surgical complications.

The carbon cost of implementing various tobacco cessation services can be integrated into the above calculation. This ranges from 8.1 to 16.3 kg CO<sub>2</sub>e/patient depending on whether it is by text messages or individual counseling (Figure 1 and 2, detailed in Appendix 2, 3 and 4). If one considers the long term savings in carbon emission of a person who stops smoking, their carbon footprint will be reduced by a factor of more than 1:1000 [24].

Of note, the cradle-to-grave life cycle of a cigarette costs 14 g of CO<sub>2</sub> [25]. This may be a further argument to motivate ecologically-minded smokers.



Excessive alcohol consumption can be identified by the preoperative assessment. A simple questionnaire like AUDIT (Alcohol Use Disorder Test) can be useful. It can predict the perioperative risk of delirium and a risk of decompensation of an asymptomatic cirrhosis or cardiac dysfunction. This latter condition can recover within 4 weeks of alcohol withdrawal [26]. Up to 15% of patients in France have alcohol related problems and 5% will develop a delirium tremens during a hospital stay [27]. These patients are 50% more likely to develop postoperative complications, which rises to 200% if they are heavy drinkers [26]. However, alcohol cessation programs do not appear to decrease LOS, thus LOS is not a factor that can be taken into account when estimating carbon cost [28].

A delirium can occur 24 to 48 hours after stopping or reducing alcohol intake and can be prevented by benzodiazepines, correct hydration and analgesia. A short LOS is less likely to be complicated by a delirium. Once symptoms start, the LOS is likely to be prolonged and can even require an ICU admission. Correct management entails considerable human resources and availability of consumables (oxygen, IV hydration, specific medicaments). The mean carbon cost of an ICU stay is between 178 000 to 624 000 g of CO<sub>2</sub> per day [29]. The quantity of Propofol needed for general anesthesia (GA) is increased by 30% in chronic alcoholic patients [30]. Its carbon footprint is lower compared to anesthetic gasses, nonetheless the cradle-to-grave carbon cost of Propofol is about 10 g of CO<sub>2</sub> for 20mL [31] and it exerts a significant impact on aquatic life. Propofol is not biodegradable in water, thus possessing a high potential for bioaccumulation. It is toxic to aquatic organisms which may result in long-term adverse effects in aquatic environments. Propofol has a Persistence Bioavailability Toxicity (PBT) index of 6 (9 being the most toxic) [32].

#### **4. Patient blood management**

PBM consists in combining a number of strategies to avoid transfusion of blood products, including increasing preoperative hemoglobin, reducing blood loss during surgery and respecting current guidelines about restrictive transfusion according to lower hemoglobin (Hb) thresholds.

Anemia is a frequent state in elderly patients and can be responsible for worse outcomes after surgery. Iron administration for a month before a surgery can correct 79% of anemias, raises hemoglobin levels by 1 g/L, reduces by 20-22% the risk of transfusion and the LOS by 24 hours [43, 45]. Patients who benefit from a PBM program are 27% more likely to be discharged directly to their domicile (rather than needing an interim stay in a rehabilitation facility) However the cost of a PBM program is not neutral: a physician visit can cost 66 kg of CO<sub>2</sub> [42], and up to 3 out-patient admissions may be necessary for the intravenous iron administration and EPO injection. This can entail a relatively high carbon cost: 198 kg of CO<sub>2</sub> (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

Major orthopedic surgery carries a significant hemorrhagic risk : between 25% to 32% of patients will often require transfusion [46, 47].

Some studies [39, 40] show that 20% to 25% of transfusions are still undertaken above the current recommended Hb thresholds. Lowering transfusion thresholds can save 250kg of CO<sub>2</sub> by patient by reducing LOS and saves 325g of CO<sub>2</sub> after each major orthopedic surgery [37]. A recent study that quantified the carbon footprint of transfusions showed that 3000 tons of CO<sub>2</sub> are wasted each year in the UK by transfusing patients above a Hb of 7-8 g/L. A single blood transfusion was responsible for the production of 6,5kg of CO<sub>2</sub> [36].

Transfusions are linked to a longer LOS [40, 41], so a restrictive strategy could shorten by 2 days the hospital stay. Additionally, 5.2% of new hospitalizations (75 kg CO<sub>2</sub>e per ambulance emergency response, 125 kg of CO<sub>2</sub> per day) could be avoided [41] in this study by applying a restrictive transfusion strategy in spinal surgery. Thus, by respecting the transfusion thresholds guidelines, a practitioner can save 10kg of CO<sub>2</sub> of LOS per patient receiving IV Iron [42] (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

For colorectal surgery, we consider an average transfusion rate of 50% [60]. The impact of preoperative management of anemia and of the respect to the transfusion threshold is about 387 026 gCO<sub>2</sub>e/patient (Figure 1 and 2, detailed in Appendix 2, 3 and 4). We decided to extrapolate the findings in spinal surgery [41] about the risk of new hospitalization of 5,2% to the colorectal surgery.

The use of tranexamic acid is now recommended to reduce transfusion requirements during and after joint replacement, which will impact on LOS in this particular surgical context. Since antifibrinolytic drug administration results in one-third less transfusion and shortening of LOS by up to 0,67 LOS [10, 37], failure to give Tranexamic acid accounts for approximately 84 179 g CO<sub>2</sub>e/patient (Figure 1 and 2, detailed in Appendix 2, 3 and 4). The precise carbon impact for tranexamic acid manufacture is not available, but can be estimated at 340 g CO<sub>2</sub>e/g per drug dose administration during surgery [31, 38].

## **5. Prehabilitation**

The potential benefits of increasing a patient's capacity to cope with surgical stress by following a prehabilitation program is appealing. In reality, there appears to be little proof of

benefits, especially in terms of LOS. Some studies [44, 48] involving patients with denutrition and sarcopenia undergoing major abdominal surgery have reported a shortened LOS (up to 2 days) with implementation of a preoperative nutrition program. It proved especially effective in diminishing pulmonary morbidity [49]. These programs were not beneficial for surgical specialties other than major abdominal and thoracic surgery, and recommendations are assigned a low grade of evidence. One study found similar results between intensive bike and aerobic activities at home on pulmonary complications [50]. Translated into carbon gain, these interventions represent a maximum of 250 kg CO<sub>2</sub> per malnourished patient (the carbon footprint of immunonutrition and protein supplementation remains unknown).

Only a few patients could benefit from these costly programs [51]. Proposing specific prehabilitation protocols to all other ERAS patient can be considered as an additional carbon cost without obvious clinical benefit. A physical reeducation program in a specialized center, costly in both electricity and transport, could be replaced by home exercises. If the patient travels 10 km twice a week for a physiotherapy session using a private car or taxi, this will equal 19.2kg of CO<sub>2</sub> without a specific benefit for the patient [52] (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

## **6. Other preoperative issues in colorectal surgery**

According to recent recommendations in colorectal surgery [7], two nutritional interventions can each reduce LOS by 24 hours: carbohydrate loading for ASA 1-2 patients and immunonutrients for carcinologic surgery (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

A considerable proportion of patients suffer from anxiety before a surgery (11 to 80%). All the ERAS societies agree that anxiolytic medication is not an effective way to prevent preoperative anxiety. It is also associated with certain side effects, including delayed time to extubation and cognitive recovery [53]. In terms of calculating carbon costs, there is no benefit pharmacological treatment of such anxiety, which it will simply generate more carbon because of medication itself and the longer time to extubation (4 min of extra time in the operating room is equivalent to roughly 160g of CO<sub>2</sub>) (Appendix 1). The delayed cognitive recovery after a benzodiazepine (e.g. Lorazepam) is difficult to convert into carbon, but if the patient cannot return home after day-case surgery, the subsequent overnight stay will increase carbon costs.

A non-pharmacology approach, such as hypnosis, is an innovative way to reduce anxiety and pain [54, 55], which has shown good success rates for appropriate patients. Patients tend to be less anxious after a 30-minutes hypnosis session, and report less pain when a painful procedure is performed under hypnosis. Nevertheless, even a brief hypnosis session costs resources. Furthermore, hypnosis is not a large-scale solution, but rather an option that can be proposed to patients with high anxiety levels.

In terms of carbon expenditure, hypnosis can reduce the length of a painful procedure [55], but on the other hand, needs 30 minutes of work from a hypnotherapist, so it is unclear if the final carbon balance is worthwhile.

## **7. Other preoperative issues in orthopedic surgery**

There is no perfect score to determine which patients will continue to experience pain and difficulty walking after major orthopedic surgery involving the spine, hip, or knee. If the predicted surgical outcome of a patient is poor consideration should be given to favoring conservative (non-interventional) therapy. A decision not to perform surgery in this context could save a lot of carbon.

Besides the benefit of ERAS for the patient, it is recognized to be useful in reinforcing teamwork. Since most recommendations of ERAS Society are about better organization and anticipation, each team member is a crucial element for success. As health workers are usually well-informed about climate change, adherence to ERAS can be strengthened by the evidence for carbon saving inherent to these protocols

Routine infectious screening before TJA surgery is not recommended. No specific data about such laboratory investigations could be retrieved, but some studies report about 50 to 100 g/CO<sub>2</sub>e per laboratory test, which is mainly due to the use of reagents [58] (Appendix 1).

## **B. During surgery**

### **1. Goal directed fluid therapy**

The carbon impact of GDFT leads to saving up to 250 000 g CO<sub>2</sub>e/patient with LOS shortened by 2 days in colorectal surgery [7] (Figure 1 and 2, detailed in Appendix 2, 3 and 4). No such effect is observed in orthopedic surgery [8, 10]. The carbon footprint of sophisticated hemodynamic monitoring was not available in the current literature, and its practice will also depend on professional habits.

### **2. Corticotherapy**

Use of dexamethasone for nausea and vomiting prophylaxis, leads to shortening LOS by 24 hours in colorectal surgery. The same anti-emetic administered to orthopedic surgery patients results in improved analgesia and less nausea but does not influence LOS. No data was found regarding indirect effects of nausea, for example the wastes in each unconsumed meal tray. The carbon impact could be major, as almost 50% of orthopedic patients in 1 study consumed less than one quarter of their first post operative meal [57]: the incidence of nausea was 42%. An 8 mg ampoule of dexamethasone could be approximated to 2.7 g CO<sub>2</sub>e [31]. Failure to administrate dexamethasone during colorectal surgery costs 125 000 g CO<sub>2</sub>/patient (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

### **3. Target thermic control**

Failure to maintain normothermia perioperatively increases the risk of SSI in colorectal surgery by 13%/degree of hypothermia, increases LOS by 1-2.6 day [58, 59] and delays return of intestinal transit. In orthopedic surgery, there is a lower bleeding but no change in LOS. The most efficient way to maintain normothermia is forced air warming, so such devices (e.g. Bair Hugger) must be integrated into the carbon costs. Considering that 1kWh equals 74 g of CO<sub>2</sub>, using the warming device set at 38° for 1 hour costs 1 225 g of CO<sub>2</sub> [61].

The estimation of blood loss for a patient with mild hypothermia is about 54mL. Approximately 50% of patients undergoing colorectal surgery will require transfusion [60] which corresponds to 650 g CO<sub>2</sub>e/patient [36]. Combined with the increased LOS associated with hypothermia, the carbon cost of non respect of appropriate temperature control is about 330 300 g CO<sub>2</sub>e/patient (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

### **4. Prophylactic antibiotic treatment**

Antibioprophylaxis reduces the risk of surgical site infection (SSI) by up to 50%, as consistently stated in current guidelines of ERAS [7, 10]. In case of lack of, or inappropriately-timed administration of antibiotics [36], there is a major risk of complications notably need for antibiotic treatment, a prolonged LOS and possible reoperation. We used actual risk of an SSI [36], the rate of unanticipated further surgery during the initial admission and the additional LOS incurred [62, 63, 67] have been taken into account for our



calculations. The total amount of carbon is 60 335 (colorectal surgery) and 194 469 g CO<sub>2</sub>/patient (TJR) each time antibiotic prophylaxis did not conform to standards. In terms of absolute risk, the rate of non-compliance with prophylactic antibiotics did not influence the carbon emission calculations (51.2% in digestive surgery and 60.8% in orthopedics surgery respectively) (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

## **5. Choice of surgical technique**

In colorectal surgery, there are known advantages to using minimally invasive surgery. The main benefits concern the recovery for the patient, with the impact in terms of LOS being accentuated when an ERAS program is implemented. After a laparoscopy, the LOS is shorter by 2 days compared to a laparotomy, and shorter by 5 days if the ERAS program is followed [7]. The cost of the total CO<sub>2</sub>e emission necessary for an hour-long laparoscopy [64] is reported to be about 100 000 g CO<sub>2</sub>e. In addition, the extra cost in terms of energy and wastes of laparoscopic techniques [65] is certain - about 6 500 g CO<sub>2</sub>. The carbon cost gap by not using laparoscopy could be as high as 518 500 g CO<sub>2</sub>e/patient (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

There is no clinical effect of the type of incision for TJA in terms of complication or LOS. The fixed cost for joint prosthetic surgery was not included in our final calculations (an estimated 190 000 g CO<sub>2</sub>e/patient) [66].

## **6. Type of anesthesia**

A “standard anesthetic protocol” is proposed [9] without clear recommendation for either anesthetic gasses or total intravenous anesthesia (TIVA). In order to avoid wasting anesthetic drugs without any adverse effects for the patient, a simple change in clinical practice could achieve a great deal. Using a lower fresh gas flow for halogenated anesthetics or TIVA during GA, a lower oxygen flow during regional anesthesia (RA) and avoiding desflurane and nitrous oxide are some of the eco-responsible strategies to implement. For TJA, these steps could lighten the carbon expenditure by 20-25% [38]. Between 2000-3000 g CO<sub>2</sub>e/patient can be saved depending on the modality of anesthesia (Figure 1 and 2, detailed in Appendix 2, 3 and 4). The carbon cost of a GA is estimated at 12 kg CO<sub>2</sub>e and 10 kg CO<sub>2</sub>e for a RA using European energy sources.

## **7. Other intraoperative issues in orthopedic surgery**

Application of a tourniquet in TKA is not recommended [10]. Tourniquet use is associated with contradictory results for the post-operative pain, carries a major risk of distal deep venous thrombosis (DVT) and shows no significant influence on transfusion requirements or LOS. This last point was not found in a recent review [68], this is the reason why we did not include any impact in terms of carbon related to the use of tourniquet.

## **C. After surgery**

### **1. Multimodal analgesia**

Beside the modality of anesthesia itself, a strategy of multimodal analgesia is an important limiting factor LOS after surgery [7, 8, 9, 10]. For colorectal surgery, up to 88 750 g CO<sub>2</sub>e can be saved per patient using IV lidocaine for laparoscopy or thoracic epidural analgesia for a laparotomy in addition to perioperative multimodal analgesia. This figure increases up to 125 000 g CO<sub>2</sub>e saved by a multimodal analgesia in TJA (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

### **2. Venous thromboembolism prevention**

A number of surgical operations are identified as high risk for DVT, including major orthopedic surgery and cancer surgery. Strict DVT prophylaxis is mandatory in these cases. Non-pharmacological prophylaxis consists of wearing compression stockings during and after surgery, and if the patient's condition permits, early mobilization, which can start as early as the PACU. These are virtually carbon free measures, which makes them very attractive (the carbon footprint of compression stockings remains unknown).

For some types of surgeries, patients will also require pharmacological prophylaxis, for example, injectable low molecular weight heparin (LMWH) or non-fractionated heparin in carcinologic surgery. Without prophylaxis, colorectal surgery has a 1.2% incidence of venous thromboembolic disease [7, 69]. For major abdominal cancer surgery, the incidence can rise to 35-45%, with a 3% of cases experiencing fatal pulmonary embolism [70].

Major orthopedic surgery also necessitates thromboprophylaxis which will reduce the risk of DVT by 50% [10]. Three options are possible: injectable heparin, oral prophylaxis using new oral anticoagulants (NOA) or aspirin alone. Despite widespread availability and evidence of efficacy for NOA, some clinicians are still reluctant to use this oral medication due to fear of hemorrhagic complications. Studies have clearly documented the safety profile of these NOA [69, 71, 72, 73]. Besides the patient comfort afforded by an oral medication compared to daily injections, using the oral route is relatively a green option. Heparin injections often need to be administered by a nurse, and furthermore, regular monitoring of platelet count is obligatory. The carbon footprint of a patient receiving injectable heparin after TJA (35 days for hip prosthesis, and 14 days for knee prosthesis) is as follows : 20km roundtrip by car for the nursing visit plus 2 blood tests per week equals 85.16 kgCO<sub>2</sub> for the hip and 34.06 kg CO<sub>2</sub> for the knee (Figure 1 and 2, detailed in Appendix 2, 3 and 4). Some types of surgeries (e.g. knee ligamentoplasty) for low risk patients do not need DVT prophylaxis, or can be managed by aspirin alone. This can help save 24.5 kg of CO<sub>2</sub> per patient.

### **3. Drains**

Intraoperative gastric decompression for colorectal surgery with a nasogastric tube does not significantly impact LOS significantly. The impact of this intervention on postoperative ileus and bronchopulmonary complications lacks sufficient precision to be converted into carbon cost.

There is a risk of catheter associated urinary tract infection (CAUTI) with a urinary probe left in-situ more than 24 hours after colon surgery. The basal risk of CAUTI is about 5% [74, 75].

This complication entails an increase in LOS of around 11 days according to recent literature [76]. Between 17 to 70% of CAUTIs can be prevented by removal or avoidance of a urinary catheter [74]. In total, the carbon cost of CAUTI related to unjustified prolonged urinary catheterization may reach 47 438 g CO<sub>2</sub>e/patient.

Orthopedic surgery does not require drains in the postoperative course: inappropriate drainage is estimated to cost 125 000 g CO<sub>2</sub>e/patient (Figure 1 and 2, detailed in Appendix 2, 3 and 4). There is insufficient data about the carbon footprint of the avoided catheter or urinary probes.

#### **4. Early mobilization and feeding**

One of the key points of ERAS is to maintain the patient as close as possible to a normal physiological state. Early mobilization, reduction in bed-rest and early oral refeeding are all strongly promoted to avoid deleterious effects such as muscle loss and insulin resistance. All ERAS guidelines [7, 10] recommend standing, or even better walking on the day of surgery, which can reduce LOS by 1 to 2 days, equivalent to 125-250 kg of CO<sub>2</sub>.

Early intensive physiotherapy can allow patients to return home 2 to 3.5 days sooner after TJA. The carbon cost of healthcare workers such as physiotherapists is directly linked to the carbon footprint of their mode of transport. If the physiotherapist treats multiple patients in the same day at the hospital, this is more eco-sustainable than if the patients visit the physiotherapist separately. This translates to 437.5 kg of CO<sub>2</sub> savings because of the work patterns of nurses and physiotherapists.

Physical exercises can be started in the PACU. This high-intensity care-zone offers reassurance for patients by allowing them to re-start oral fluids and mobilize in their bed in total security. The PACU should play a specific role in improving the patient's immediate postoperative care as needed; but should be avoided for those who don't. A shortened PACU stay can be achieved thanks to a better prevention of PONV (each vomiting patient stays at least 20min more in the PACU, consuming plastic and energy) [77], or by avoiding hypnotics when using loco regional anesthesia. The energetic cost of a PACU is hard to determine because of a large variety of different types and sizes of PACU.

Early feeding in colorectal surgery can be linked to a shorter LOS (0.89 day, or 111kg of CO<sub>2</sub>) (Figure 1 and 2, detailed in Appendix 2, 3 and 4).

## **5. Other postoperative issues in colorectal surgery**

One interesting measure could be allowing patients to chew gum as soon as they arrive in the PACU, helping to restore a bowel transit sooner, but the impact on LOS is not certain [7]. Education of patients after colorectal surgery about how to deal with a stoma can be done by stomatherapists. Besides helping patients to cope with this, it helps shorten the hospital stay by 2.5 days [78]. We will not take into account this in our calculation because of the lack of information about the carbon cost of this human resource.

## **6. Other postoperative issues in orthopedic surgery**

Cryotherapy reduces pain, but does not significantly reduce surgical bleeding and does not impact on LOS [10]. Some devices use compressive cryotherapy (e.g. Game ready) but are consuming a lot of energy, by needing ice to be filled with. Probably these devices are more carbon costing than efficient for the patient, but precise carbon costs are difficult to obtain.

## **7. Post discharge**

With the development of same day surgery, surgical and anesthesia teams need to use solid scores in their practice to avoid complications due to patients who would have gone home too early. Besides, useless time spent in the hospital while waiting before heading home can be shortened by using these scores. A 12 item score can allow a patient to leave the PACU without waiting a minimum time thanks to the Aldrete score [79, 80], which helps saving about 24min per patient in the PACU. The carbon cost of PACU is not known, but there is a clear gain of time and this can allow more patients to have surgery. The final carbon price would be divided by more patients, and so it would be lower. Same opportunity with the Chung score with 9 items which allow the patient to leave the hospital after surgery. This is a good way to shorten the LOS without spending any carbon and just completing a physical examination.

How to determine which patient can go home, and who needs a rehabilitation center?

Some functional surgeries benefit patients who can cope with the changes and efforts to give to recover. Another partly subjective assessment has to be done by the surgeon before

performing the surgery. If the patient appears to be skilled to do his rehabilitation at home, it can help save a lot of carbon, by not using a high electricity consuming health center. But if the patient is not well surrounded by friends or family, and cannot find physiotherapists and nurses around his home, the praticien should notice this before planning the surgery, because an unplanned need for a rehab center will delay the rehab and will cost some carbon with useless hospitalization days. There is a need for stronger scores to find out the needs of the patients for rehab after surgery.

While the patient is back home, he can call back to the hospital if he needs to ask a question or if he has an issue like pain or ileus. Sometimes, a nurse is in charge of calling the patients to make sure they make it through the postoperative period. Some studies propose a post discharge text message instead of a telephone call: the results are good with less than 10% of non-response and a high satisfaction.

These calls usually last for 6 min with a rate of response of 87% and cost 342g of CO<sub>2</sub> [81]. A text message is also less consuming carbon, with a cost of 0.014g CO<sub>2</sub>, and it divides by 4 the need of being called, so it costs 85.5g of CO<sub>2</sub> per day [13] (Figure 1 and 2, detailed in Appendix 2, 3 and 4).



## **D. Additional issues : ERAS and optimized hospital stay: time is carbon**

Beyond all recommendations about ERAS pathway, one should consider a global vision of the healthcare system. During our research, we found some clues to reduce and prevent waste of carbon by saving time at all steps of a patient's hospital stay.

First of all, a simple prevention of patient cancellation could help reducing carbon costs by just calling the patient the day before the surgery [82, 14, 83, 84] and by collecting some information with an entry checklist by a nurse : cancellations are responsible for 7% of time of OR wasted because of canceled cases [85]. Considering one minute of OR costs 40g of CO<sub>2</sub>, and that a day lasts 600min, 7% of that corresponds to 42min wasted and 1,680g CO<sub>2</sub> per day (Appendix 1). A D-1 day call could help dividing by 2 or even 4 this waste of time.

A large point to develop is the use of tele counseling. Indeed, the scope 3 of the greenhouse gas emissions characterized by the costs for goods, services and travels, account for a wide part of the total emissions, from 12% [86] to 92% [87], depending on countries. Tele counseling could help to divide by ten the CO<sub>2</sub> emission based on European standards of car emissions, with a carbon footprint for a 15min appointment of 450 to 2,100 gCO<sub>2</sub>e [88, 89].

Lean management is an interesting approach of optimization OR time [90, 91, 92]. By using some Lean programs like Sigma method, by reducing late starts by 5 min could save 200g CO<sub>2</sub> per patient, and up to 640g with a better OR organization, reducing the whole turn over time by 16min (Appendix 1).

In the same way of calculating carbon by considering that time is carbon, time spent in the PACU gains to be optimized. By using discharge criteria, time in PACU can be shortened by 24%. As we didn't find the carbon footprint of PACU, we cannot calculate these costs. The optimization of the OR schedule so that PACU is not overloaded can lead to 7% of time available for surgery [91], which can be converted into a 42 min and 1680g CO<sub>2</sub> (Appendix 1).

To go further in our topic, more actions could be used as levers to lower carbon emissions of the healthcare system. The widespread isolation of buildings [42], improved gas scavenging system with halogens [93], generalized water sparing scrubs techniques [94] or sorting domestic and clinical waste in OR [16].

## DISCUSSION

Even if the attention for carbon footprint and greener solution within the healthcare system is growing, there is a need for more studies to assess carbon costs of all steps of the patient pathway. The literature is heterogeneous, as previously described [96]. One of the weaknesses of our study are the unknown costs and the approximations of the calculations performed. One classic example of this conundrum is the missing life cycle assessment for pharmaceuticals due to proprietary nature of drug synthesis, thus only limited data is available [95]. Missing data also proved to be a problem in evaluating energy consumption for monitoring device, involved in volemia optimization for example. It is also difficult to include the total repercussions of an event: in particular, the professional and social impacts are hard to translate in terms of carbon and financial costs [97].

Our calculations mainly assume that a patient consumes no more than 125kg of CO<sub>2</sub> at home per day, and that a day at hospitalization is producing more CO<sub>2</sub> than a personal dwelling.

It is hard to approximate the distance traveled by a nurse or a physiotherapist, especially in countries with geographical heterogeneity like France or those where long distances may be traveled. To simplify our calculations, the distance to provide care to a patient was approximately 20km round trip. The fact that we chose to include the carbon cost of distance traveled for nursing home visits for LMWH injections could also be criticized, because both LMWH and alternative oral prophylaxis match current recommendations. We included this carbon cost to indicate that an alternative process, namely oral medications, exists.

Furthermore, some professional practices during a GA are very carbon costly, and are mentioned as “standard anesthetic protocol” [9]. One of our objectives would be to define a standard protocol with ecological concerns.

The type of energy source is another factor that can modify carbon costs by a factor from 1 to 8, limiting the external validity of our results. An important factor with a profound impact is the extraction of carbon dioxide for which open access is rarely available from companies. Secondly, there are possible important variations in carbon footprint depending on energy sources available in a particular country. Better transparency is needed about the sources of energy for all healthcare facilities.

Finally, some carbon balances were not quantified in the existing literature, for example, the carbon costs of time spent in the recovery room or staff overtime accrued in managing patient anxiety. We also exclude data with divergent clinical results such as tourniquet usage in orthopedic surgery.

## CONCLUSION


The non-compliance to the ERAS protocol is costly in almost all the steps in perioperative time. The balance in terms of carbon footprint is strongly positive. Indeed, the total cost for a colorectal surgery is about 2,500,000 gCO<sub>2</sub>e/patient, the major part involved is during the intraoperative time. In joint replacement surgery, the total cost is about 1,400,000 gCO<sub>2</sub>e/patient, the major part involved is during the postoperative time. Results are summarized in Figure 3.

Regarding the ecological balance, our study shows an additional argument to respect the current ERAS guidelines for colorectal and joint replacement surgery. Additional actions beyond ERAS pathway are strongly efficient on carbon savings, and the future of our practices in anesthesia will have to face the global warming emergency.



All these points must be part of a general dynamic, sometimes hard to implement in the hospital where the so-called “environmental numbness” [98] could limit carbon transition. It can correspond to the temporal justification of the medical staff being too busy to pay attention to short term concerns, the presupposed expensiveness of green practices or even the “moral offset” in which working in healthcare is already enough. Transition to a new model of care in which human and environment health is both promoted and rewarded must be considered [99].

# FIGURES

Figure 1: Carbon costs of perioperative actions based on ERAS recommendations



**Figure 2: Sharing out of peri operative carbon costs**



## APPENDIX

### Appendix 1 – Details in calculations

- **1min in operating room :**

Calculations from 3 hospitals in USA, UK and Canada

1min in OR =  $500,000\text{g CO}_2\text{e} / 365\text{d} / 24\text{h} / 60\text{min} / 22\text{ OR} = 40\text{ g CO}_2\text{e}/\text{min}$

Considering countries with carbon energy: 363 g CO<sub>2</sub>e/min

We chose countries with the least carbon energy (Canada, Vancouver) to meet with France consumption

Not included : wastes count

*Sources: McNeil Lancet 2017*

- **Carbon cost per bed day**

125 kg CO<sub>2</sub>E /bed-day and 76 kg CO<sub>2</sub>e /outpatient appointment for acute care

Estimation in NHS in England

*Source: Tennison, 2021, Lancet*



**Appendix 2 - Formulas for carbon footprint calculations** (time in minutes, carbon costs in grams of CO<sub>2</sub>e, length of stay in days, amount in grams, distance in kilometers)

Formula A	increase in length of stay x average carbon cost for additional bed-day in hospital
Formula B	wasted time in operating room x average carbon cost for additional minutes in operating room
Formula C	variation in complication rate x absolute risk for general population x average additional length of hospital stay per complication x average carbon cost for additional bed-day in hospital
Formula D	proportion of TJR surgeries transfused x variation in rate of transfusion x average carbon cost per transfusion
Formula D'	proportion of general surgeries transfused x variation in rate of transfusion x average carbon cost per transfusion
Formula E	absolute over risk of unnecessary blood test x average carbon cost per blood test
Formula F	amount of wasted involved x carbon cost per kilograms of wastes incinerated
Formula G	days of treatment x distance traveled by nurses x carbon cost per kilometer by car + number of blood test x carbon cost per CBC

Formula	reoperative rate x carbon cost per surgery (1 hour long coelioscopy or TJR
H	surgery) + 1 hour long anesthesia

**Appendix 3 - Details in calculations for each ERAS wrong turn costs (results in gCO<sub>2</sub>e/patient)**

ERAS points	Formula	Calculation	Results
Tobacco cessation	C	$0,83 \times 0,45 \times 3 \times 125\ 000$	140 063
Diabetes balance	C	$0,4 \times 0.0072 \times 18 \times 125\ 000$	6 480
Patient blood management <ul style="list-style-type: none"> <li>• Iron supply</li> <li>• Tranexamic acid</li> <li>• Restrictive transfusion strategy</li> </ul>	D A  A D  A A D	$0,25 \times 0,2 \times 6\ 500$ $1 \times 125\ 000$  $0,67 \times 125\ 000$ $0,2 \times 0,33 \times 6\ 500$  $0,052 \times 200\ 000$ $2 \times 125\ 000$ $0,25 \times 0,2 \times 6\ 500$	470 229 (orthopedic surgery)
Patient blood management <ul style="list-style-type: none"> <li>• Iron supply</li> </ul>	D' A	$0,25 \times 0,5 \times 6\ 500$ $1 \times 125\ 000$	387 026 (colorectal surgery)

<ul style="list-style-type: none"> <li>Restrictive transfusion strategy</li> </ul>	A A D'	0,052 x 200 000 2 x 125 000 0,25 x 0,5 x 6 500	
Prehab	A	2 x 125 000	250 000 (colorectal surgery)
Carbohydrates and immunonutrients	A	2 x 125 000	250 000 (colorectal surgery)
Routine infectious blood sample	none		50 - 100 (orthopedic surgery)
Goal directed therapy	A	2 x 125 000	250 000 (colorectal surgery)
Corticotherapy	A	1 x 125 000	125 000 (colorectal surgery)
TTC	C A D'	0,13 x 0,0159 x 18 x 125 000 2,6 x 125 000 0.2 x 0,5 x 6 500	330 300 (colorectal surgery)  
Prophylactic antibiotic treatment	C H C	0,5 x 0,0157 x 18 x 125 000 0,381 x (100 000 + 12 000) 0,5 x 0,0135 x 21 x 125 000	60 335 (colorectal surgery) 

	H	0,875 x (190 000 + 12 000)	194 469 (orthopedic surgery)
Choice in surgery technic: laparoscopy	A	5 x 125 000	625 000 (colorectal surgery)
Balanced anesthesia	none		2000 - 3000
Multimodal analgesia	A	0,7 x 125 000	88 750 (colorectal surgery)
	A	1 x 125 000	125 000 (orthopedic surgery)
Venous thromboembolism prevention	G	35 x 10 x 2 x 120 + 10 x 116	85 160 (orthopedic surgery - THR)
	G	14 x 10 x 2 x 120 + 4 x 116	34 064 (orthopedic surgery - TKR)
Drains	C	0,69 x 0,05 x 11 x 125 000	47 438 (colorectal surgery)
	A	1 x 125 000	125 000 (orthopedic surgery)
Early mobilization	A	2 x 125 000	250 000 (orthopedic surgery and colorectal surgery)

Physiotherapy	A	3,5 x 125 000	437 500 (orthopedic surgery)
Early feeding	A	0,89 x 125 000	111 250 (colorectal surgery)

**Appendix 4 - Details in calculations for costs involved in each ERAS pathway subpart (in gCO<sub>2</sub>e/patient)**

Tobacco cessation services	8 100 to 16 100
Tranexamic acid (gram)	340 (orthopedic surgery)
PBM: Physician visit for iron perfusion and EPO injection	198 000 (orthopedic surgery)
Prehab: physiotherapist transport	19 200 (colorectal surgery)
Laparoscopy	100 000 (per hour) (colorectal surgery)
Wastes and energy consumption difference (Laparoscopy 29 200 vs Laparotomy 22 700)	6 500 (colorectal surgery)
ABSW, active body surface warming (Bair Hugger)	1 225 (per hour) (colorectal surgery)
Post discharge call	86 to 342

## LEXICON

- ERAS Enhanced recovery after surgery
- SSI Surgical site infection
- LOS Length of stay
- TJA Total joint arthroplasty



THÈSE SOUTENUE PAR : Charles MORET Et Diana KABANOVA

TITRE :

LA RAAC COMME NORME ÉCO-RESPONSABLE : CHAQUE ÉTAPE COÛTE DU CARBONE

#### CONCLUSION :

##### Objectif

Déterminer l'empreinte carbone de la déviation à un protocole de RAAC après une chirurgie orthopédique lourde du membre inférieur type PTH/PTG ou après une chirurgie colorectale.

##### Méthode

Notre revue narrative a été construite à partir d'articles référencés dans la base PubMed. Nos recherches ont été basées sur l'impact environnemental d'un protocole de RAAC dans le cadre de la chirurgie colorectale et d'arthroplastie de hanche ou de genou. Chaque point des recommandations françaises et internationales a été étudié. Pour nos recherches, nous avons utilisé les termes "empreinte carbone", "durabilité", "coût énergétique", "empreinte environnementale", "évaluation du cycle de vie" ET un mot clef de chaque sous partie des recommandations RAAC.

##### Résultats

Durant la période péri-opératoire, les écarts à un protocole de RAAC, dans le cadre d'une chirurgie colorectale ou d'une arthroplastie lourde du membre inférieur, sont très coûteux en équivalent carbone. Le surcoût est estimé à environ 2500 KgCO<sub>2</sub>/patient pour une chirurgie colorectale. La part la plus importante se produit durant la période peropératoire. Le surcoût est estimé à environ 1400 KgCO<sub>2</sub>/patient dans le cadre d'une chirurgie orthopédique majeure. La part la plus importante se produit durant la période postopératoire. Ceci est un argument supplémentaire à l'application d'un protocole RAAC lorsque celui-ci est recommandé. Il existe des leviers supplémentaires, en dehors des protocoles de RAAC, pour réduire notre empreinte carbone dans notre pratique de l'anesthésie. Celle-ci devra probablement s'adapter à l'urgence du réchauffement climatique.

##### Conclusion

Le non-respect des recommandations RAAC est très coûteux en termes de carbone. Notre étude montre un argument supplémentaire de la nécessité de mettre en place des parcours RAAC pour le patient, qui devrait être considéré comme la norme éco responsable d'une prise en charge chirurgicale. Des considérations plus globales et organisationnelles sont également à mettre en place pour réduire les émissions de CO<sub>2</sub> dans le secteur de la santé.

VU ET PERMIS D'IMPRIMER

Grenoble, le :

LE DOYEN DE L'UFR DE MÉDECINE

Le Doyen de l'UFR de Médecine

Pr. Patrice MORAND

Pr. Patrice MORAND

LE PRÉSIDENT DU JURY



Pr. Pierre ALBALADEJO

## BIBLIOGRAPHY

- [1] The Shift Project, Décarboner la Santé pour soigner durablement, 2021
- [2] Pichler P-P, Jaccard IS, Weisz U, Weisz H. International comparison of health care carbon footprints. *Environ Res Lett* 2019;14:064004
- [3] Wu R. The carbon footprint of the Chinese health-care system: an environmentally extended input–output and structural path analysis study. *The Lancet Planetary Health* 2019;3:e413–9
- [4] [https://www.legifrance.gouv.fr/codes/article\\_lc/LEGIARTI000039369665](https://www.legifrance.gouv.fr/codes/article_lc/LEGIARTI000039369665)
- [5] Burrell R. H., *Primum non nocere*. *Anaesth Intensive Care*, 2018
- [6] Muñoz A. Reducing Health Care’s Carbon Footprint—The Power of Nursing 2012;60:4
- [7] Alfonsi P, Slim K, Chauvin M, Mariani P, Faucheron J-L, Fletcher D. Réhabilitation rapide après une chirurgie colorectale programmée. *Annales Françaises d’Anesthésie et de Réanimation* 2014;33:370–84
- [8] Wainwright TW, Gill M, McDonald DA, Middleton RG, Reed M, Sahota O, et al. Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS) Society recommendations. *Acta Orthopaedica* 2020;91:3–19
- [9] Gustafsson UO, Scott MJ, Hubner M, Nygren J, Demartines N, Francis N, et al. Guidelines for Perioperative Care in Elective Colorectal Surgery: Enhanced Recovery After Surgery (ERAS®) Society Recommendations: 2018. *World J Surg* 2019;43:659–95

- [10] Delaunay L, Jenny JY, Albi-Feldzer A, Alfonsi P, Bloc S, Cittanova ML, et al. En association avec la Société Française de Chirurgie Orthopédique et Traumatologique (SOFOT)
- [11] Scott MJ, Baldini G, Fearon KCH, Feldheiser A, Feldman LS, Gan TJ, et al. Enhanced Recovery After Surgery ( ERAS ) for gastrointestinal surgery, part 1: pathophysiological considerations. *Acta Anaesthesiol Scand* 2015;59:1212–31
- [12] McGain F, Muret J, Lawson C, Sherman JD. Environmental sustainability in anaesthesia and critical care. *British Journal of Anaesthesia* 2020;125:680–92
- [13] Berners-Lee M. “How Bad are Bananas?: The Carbon Footprint of Everything“. Greystone Books; 2011
- [14] Chan CYW. Outpatient Pre-operative Assessment in Joint Replacement Surgery 2008;63:4
- [15] Anthony Luis B. Chua, The Environmental Impact of Interventional Radiology: An Evaluation of Greenhouse Gas Emissions from an Academic Interventional Radiology Practice. *J Vasc Interv Radiol* 2021; 32:907–915
- [16] Southorn T, Norrish A, Gardner K, Baxandall R. Reducing the Carbon Footprint of the Operating Theatre: A Multicentre Quality Improvement Report. *Journal of Perioperative Practice* 2013;23:144–6
- [17] Sørensen LT. Wound Healing and Infection in Surgery: The Pathophysiological Impact of Smoking, Smoking Cessation, and Nicotine Replacement Therapy A Systematic Review. *Annals of Surgery* 2012;255:1069–79

- [18] Grønkjær M, Eliassen M, Skov-Ettrup LS, Tolstrup JS, Christiansen AH, Mikkelsen SS, et al. Preoperative Smoking Status and Postoperative Complications: A Systematic Review and Meta-analysis. *Annals of Surgery* 2014;259:52–71
- [19] Mills E, Eyawo O, Lockhart I, Kelly S, Wu P, Ebbert JO. Smoking Cessation Reduces Postoperative Complications: A Systematic Review and Meta-analysis. *The American Journal of Medicine* 2011;124:144-154.e8
- [20] Thomsen T, Villebro N, Møller AM. Interventions for preoperative smoking cessation. *Cochrane Database of Systematic Reviews* 2014
- [21] Dautzenberg B, Masquelet AJ. Le défi de la prise en charge du tabagisme péri-opératoire, *BEH* n° 21-22/2006
- [22] Conférence d'experts Tabagisme-perioperatoire. *Annales Françaises d'Anesthésie et de Réanimation* 25 (2006) 479–481
- [23] Møller AM, Villebro N, Pedersen T, Tønnesen H. Effect of preoperative smoking intervention on postoperative complications: a randomized clinical trial. *The Lancet* 2002;359:114–7
- [24] Smith AJB, Tennison I, Roberts I, Cairns J, Free C. The carbon footprint of behavioral support services for smoking cessation. *Tob Control* 2013;22:302–7
- [25] Zafeiridou M, Hopkinson NS, Voulvoulis N. Cigarette Smoking: An Assessment of Tobacco's Global Environmental Footprint Across Its Entire Supply Chain. *Environ Sci Technol* 2018;52:8087–94
- [26] Kennedy ND, Winter DC. Impact of alcohol & smoking on the surgical management of gastrointestinal patients. *Best Practice & Research Clinical Gastroenterology* 2017;31:589–95

- [27] Mantz J., Gestion des syndromes de sevrage en réanimation, MAPAR 2011
- [28] Oppedal K, Moller AM, Pedersen B et al. Preoperative alcohol cessation prior to elective surgery. Cochrane Database Syst Rev 2012
- [29] McGain F, Burnham JP, Lau R, Aye L, Kollef MH, McAlister S. The carbon footprint of treating patients with septic shock in the intensive care unit 2019:18.
- [30] Fassoulaki A, Farinotti R, Servin F, Desmots JM. Chronic Alcoholism Increases the Induction Dose of Propofol in Humans. Anesth Analg 1993;77:553-6
- [31] Parvatker AG, Tunceroglu H, Sherman JD, Coish P, Anastas P, Zimmerman JB, et al. Cradle-to-Gate Greenhouse Gas Emissions for Twenty Anesthetic Active Pharmaceutical Ingredients Based on Process Scale-Up and Process Design Calculations. ACS Sustainable Chem Eng 2019;7:6580–91
- [32] Sherman JD, McGain F, Lem M, Mortimer F, Jonas WB, MacNeill AJ. Net zero healthcare: a call for clinician action. BMJ 2021:n1323
- [33] Stryker LS, Abdel MP, Morrey ME, Morrow MM, Kor DJ, Morrey BF. Elevated Postoperative Blood Glucose and Preoperative Hemoglobin A1C Are Associated with Increased Wound Complications Following Total Joint Arthroplasty. Journal of Bone and Joint Surgery 2013;95:808–14.
- [34] Prothèse de hanche ou de genou: diagnostic et prise en charge de l'infection dans le mois suivant l'implantation, HAS, 2014
- [35] Gestion du patient diabétique en périopératoire. Fiches pratiques. Anesthésie & Réanimation 2017;3:263–300

- [36] Surveillance des infections du site opératoire dans les établissements de santé français. Mission Spicmi, septembre 2020. Données 2018 du réseau ISO Raisin
- [37] Hibbs SP, Thomas S, Murphy MF. Convergence of zero carbon healthcare with patient blood management. *BMJ* 2021:n3112
- [38] McGain, Carbon Footprint of General, Regional, and Combined Anesthesia for Total Knee Replacements. *Anesthesiology*, 2021
- [39] Goodnough LT, Maggio P, Hadhazy E, Shieh L, Hernandez-Boussard T, Khari P, et al. Restrictive blood transfusion practices are associated with improved patient outcomes: Reduced transfusions with improved patient outcomes. *Transfusion* 2014;54:2753–9
- [40] Goodnough LT, Shieh L, Hadhazy E, Cheng N, Khari P, Maggio P. Improved blood utilization using real-time clinical decision support: Best practice alerts decrease blood use. *Transfusion* 2014;54:1358–65
- [41] Elsamadicy AA, Adogwa O, Vuong VD, Mehta AI, Vasquez RA, Cheng J, et al. Association of Intraoperative Blood Transfusions on Postoperative Complications, 30-Day Readmission Rates, and 1-Year Patient-Reported Outcomes. *Spine* 2017;42:610–5
- [42] Tennison I, Roschnik S, Ashby B, Boyd R, Hamilton I, Oreszczyn T, et al. Health care's response to climate change: a carbon footprint assessment of the NHS in England. *The Lancet Planetary Health* 2021;5:e84–92
- [43] Franchini M, Marano G, Veropalumbo E, Masiello F, Pati I, Candura F, et al. Patient blood management: a revolutionary approach to transfusion medicine. *Blood Transfusion* 2019.

- [44] Lambert Joel E et al. The Impact of Prehabilitation on Patient Outcomes in Hepatobiliary, Colorectal, and Upper Gastrointestinal Cancer Surgery. *Annals of Surgery* 2021; 274:1-70-77
- [45] Van Remoortel H, Laermans J, Avau B, Bekkering G, Georgsen J, Manzini PM, et al. Effectiveness of Iron Supplementation With or Without Erythropoiesis-Stimulating Agents on Red Blood Cell Utilization in Patients With Preoperative Anaemia Undergoing Elective Surgery: A Systematic Review and Meta-Analysis. *Transfusion Medicine Reviews* 2021;35:103–24.
- [46] Bruce W. Practical recommendations for patient blood management and the reduction of perioperative transfusion in joint replacement surgery *ANZ J Surg* 2013;83: 222–229
- [47] Qian F. Variation of Blood Transfusion in Patients Undergoing Major Noncardiac Surgery. *Annals of Surgery* 2013;257:2
- [48] Gillis C, Buhler K, Bresee L, Carli F, Gramlich L, Culos-Reed N, et al. Effects of Nutritional Prehabilitation, With and Without Exercise, on Outcomes of Patients Who Undergo Colorectal Surgery: A Systematic Review and Meta-analysis. *Gastroenterology* 2018;155:391-410.e4.
- [49] Hughes MJ, Hackney RJ, Lamb PJ, Wigmore SJ, Christopher Deans DA, Skipworth RJE. Prehabilitation Before Major Abdominal Surgery: A Systematic Review and Meta-analysis. *World J Surg* 2019;43:1661–8
- [50] Carli F, Charlebois P, Stein B, Feldman L, Zavorsky G, Kim DJ, et al. Randomized clinical trial of prehabilitation in colorectal surgery. *British Journal of Surgery* 2010;97:1187–97

- [51] Dort JC, Farwell DG, Findlay M, Huber GF, Kerr P, Shea-Budgell MA, et al. Optimal Perioperative Care in Major Head and Neck Cancer Surgery With Free Flap Reconstruction: A Consensus Review and Recommendations From the Enhanced Recovery After Surgery Society. *JAMA Otolaryngol Head Neck Surg* 2017;143:292
- [52] Ecoscore n.d. <https://ecoscore.be/en/home>
- [53] Maurice-Szamburski A, Auquier P, Viarre-Oreal V, Cuvillon P, Carles M, Ripart J, et al. Effect of Sedative Premedication on Patient Experience After General Anesthesia: A Randomized Clinical Trial. *JAMA* 2015;313:916
- [54] Saadat et al. Hypnosis Reduces Preoperative Anxiety in Adult Patients, *Anesth Analg* 2006;102:1394–6
- [55] Elvira V Lang, et al. Adjunctive non-pharmacological analgesia for invasive medical procedures: a randomized trial, *Lancet*, 2020
- [56] McAlister S, Barratt AL, Bell KJ, McGain F. The carbon footprint of pathology testing. *Medical Journal of Australia* 2020;212:377–82
- [57] Karola Traviss et al. Rethinking postoperative diets for short-stay orthopedics surgery patients, *Journal of the American Dietetic Association*, 1997
- [58] Kurz et al. Perioperative normothermia to reduce the incidence of surgical wound infection and Shorten hospitalization, *NEJM*, 1996
- [59] Andrew V. Scott, Compliance with Surgical Care Improvement Project for Body Temperature Management (SCIP Inf-10) Is Associated with Improved Clinical Outcomes. *Anesthesiology*, 2015



- [60] Benoist. La transfusion périopératoire en chirurgie colorectale, *Annales de chirurgies*, 130 (2005) 365–37
- [61] [https://www.bairhugger.com/3M/fr\\_CA/bair-hugger-ca/products/~/Module-de-réchauffement-Bair-Hugger-MC-3M-MC-77500/?N=8707795+8707798+8710661+8710865+8711017+8711716+8717793+8744795+8753191+3291780535&rt=rud](https://www.bairhugger.com/3M/fr_CA/bair-hugger-ca/products/~/Module-de-réchauffement-Bair-Hugger-MC-3M-MC-77500/?N=8707795+8707798+8710661+8710865+8711017+8711716+8717793+8744795+8753191+3291780535&rt=rud).
- [62] Badia JM, Casey AL, Petrosillo N, Hudson PM, Mitchell SA, Crosby C. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *Journal of Hospital Infection* 2017;96:1–15
- [63] Lamarsalle L, Hunt B, Schauf M, Szwarzensztein K, Valentine WJ. Evaluating the clinical and economic burden of healthcare-associated infections during hospitalization for surgery in France. *Epidemiol Infect* 2013;141:2473–82
- [64] Power NE, Silberstein JL, Ghoneim TP, Guillonneau B, Touijer KA. Environmental Impact of Minimally Invasive Surgery in the United States: An Estimate of the Carbon Dioxide Footprint. *Journal of Endourology* 2012;26:1639–44
- [65] Woods et al. Carbon footprint of robotically-assisted laparoscopy, laparoscopy and laparotomy: a comparison, *Int J Med Robotics Comput Assist Surg* 2015; 11: 406–412
- [66] Delaie. Impact écologique de la réalisation d'une prothèse totale de genou : analyse du cycle de vie de la matière première jusqu'aux implants définitifs, 2021
- [67] IQSS 2021 - ISO PTG : Rapport résultats nationaux détaillés 2021
- [68] Arthur JR, Spangehl MJ. Tourniquet Use in Total Knee Arthroplasty. *J Knee Surg* 2019;32:719–29

- [69] Samama C-M, Gafsou B, Jeandel T, Laporte S, Steib A, Marret E, et al. Prévention de la maladie thromboembolique veineuse postopératoire. Actualisation 2011. Texte court. *Annales Françaises d'Anesthésie et de Réanimation* 2011;30:947–51
- [70] Barret PA. *Société française d'anesthésie et de réanimation* 2005:25
- [71] Van der Veen L, Segers M, van Raay JJ, Gerritsma-Bleeker CL, Brouwer RW, Veeger NJ, et al. Bleeding complications of thromboprophylaxis with dabigatran, nadroparin or rivaroxaban for 6 weeks after total knee arthroplasty surgery: a randomised pilot study. *BMJ Open* 2021;11:e040336
- [72] Yu Z, Shan P, Yang X, Lou X. Comparison of efficiency and safety of rivaroxaban, apixaban and enoxaparin for thromboprophylaxis after arthroplastic surgery: a meta-analysis. *Bioscience Reports* 2018;38:BSR20180423
- [73] Eriksson BI, Huisman MV, Muehlhofer E. Rivaroxaban versus Enoxaparin for Thromboprophylaxis after Hip Arthroplasty. *N Engl J Med* 2008:11.
- [74] Rebmann T, Greene LR. Preventing catheter-associated urinary tract infections: An executive summary of the Association for Professionals in Infection Control and Epidemiology, Inc, Elimination Guide. *American Journal of Infection Control* 2010;38:644–6.
- [75] Kin C, Rhoads KF, Jalali M, Shelton AA, Welton ML. Predictors of Postoperative Urinary Retention After Colorectal Surgery. *Diseases of the Colon & Rectum* 2013;56:738–46.
- [76] Farsi. Risk Factors and Outcomes of Postoperative Catheter-Associated Urinary Tract Infection in Colorectal Surgery Patients: A Retrospective Cohort Study, 2012
- [77] Tong J. Gan, Consensus Guidelines for the Management of Postoperative Nausea and Vomiting, 2014

- [78] Forsmo et al. Pre- and postoperative stoma education and guidance within an enhanced recovery after surgery (ERAS) programme reduces length of hospital stay in colorectal surgery, 2016
- [79] Phillips NM, Street M, Kent B, Haesler E, Cadeddu M. Post-anaesthetic discharge scoring criteria: key findings from a systematic review. *International Journal of Evidence-Based Healthcare* 2013;11:275–84
- [80] Brown I, Jellish WS, Kleinman B, Fluder E, Sawicki K, Katsaros J, et al. Use of postanesthesia discharge criteria to reduce discharge delays for inpatients in the postanesthesia care unit. *Journal of Clinical Anesthesia* 2008;20:175–9
- [81] Leconte D, Beloeil H, Dreano T, Ecoffey C. Post Ambulatory Discharge Follow-up Using Automated Text Messaging. *J Med Syst* 2019;43:217
- [82] Al Talalwah N, McIltrout KH. Cancellation of Surgeries: Integrative Review. *Journal of PeriAnesthesia Nursing* 2019;34:86–96
- [83] Singhal R, Warburton T, Charalambous C. Reducing Same Day Cancellations Due to Patient Related Factors in Elective Orthopaedic Surgery: Experience of a Center in the UK. *Journal of Perioperative Practice* 2014;24:70–4
- [84] Kaddoum et al. Causes of cancellations on the day of surgery, 2016
- [85] Keller et al. Causes of elective surgery cancellation, 2014
- [86] MacNeill AJ, Lillywhite R, Brown CJ. The impact of surgery on global climate: a carbon footprinting study of operating theaters in three health systems. *The Lancet Planetary Health* 2017;1:e381–8

- [87] Devroedt C. Trophée BEGES, 2018
- [88] Holmner et al. Carbon Footprint of Telemedicine Solutions, 2014
- [89] Beloeil H, Albaladejo P. Initiatives to broaden safety concerns in anesthetic practice: The green operating room. *Best Practice & Research Clinical Anaesthesiology* 2021;35:83–91
- [90] Cima et al. Use of Lean and Six Sigma Methodology to Improve Operating Room Efficiency in a High Volume Tertiary-Care Academic Medical Center, 2011
- [91] Levine et Dunn. Optimizing Operating Room Scheduling, 2015
- [92] Fong AJ, Smith M, Langerman A. Efficiency improvement in the operating room. *Journal of Surgical Research* 2016;204:371–83
- [93] Barwise et al. An Initial Evaluation of a Novel Anesthetic Scavenging Interface, 2011
- [94] Water use in the surgical scrub: surgeons can reduce their environmental footprint. *ANZ Journal of Surgery* 2009;79:319–20
- [95] McAlister et al. The Environmental footprint of morphine a life cycle, 2016
- [96] Slim, Kareem; Selvy, Marie; Albaladejo, Pierre. Enhanced recovery programs and carbon footprint. *Anaesthesia Critical Care & Pain Medicine* 2020;39(5):665–666
- [97] Alfonso JL, Pereperez SB, Canoves JM, Martinez MM, Martinez IM, Martin-Moreno JM. Are we really seeing the total costs of surgical site infections? A Spanish study. *Wound Repair Regen* 2007;15:474–81
- [98] McGain et Naylor. Environmental sustainability in hospitals, 2014
- [99] Moynihan. The greening of medicine, 2012

# Le serment d'Hippocrate

Texte revu par l'Ordre des médecins en 2012

“ **Au moment d'être admis(e) à exercer la médecine, je promets et je jure d'être fidèle aux lois de l'honneur et de la probité.**

Mon premier souci sera de rétablir, de préserver ou de promouvoir la santé dans tous ses éléments, physiques et mentaux, individuels et sociaux.

Je respecterai toutes les personnes, leur autonomie et leur volonté, sans aucune discrimination selon leur état ou leurs convictions. J'interviendrai pour les protéger si elles sont affaiblies, vulnérables ou menacées dans leur intégrité ou leur dignité. Même sous la contrainte, je ne ferai pas usage de mes connaissances contre les lois de l'humanité.

J'informerai les patients des décisions envisagées, de leurs raisons et de leurs conséquences.

Je ne tromperai jamais leur confiance et n'exploiterai pas le pouvoir hérité des circonstances pour forcer les consciences.

Je donnerai mes soins à l'indigent et à quiconque me les demandera. Je ne me laisserai pas influencer par la soif du gain ou la recherche de la gloire.

Admis(e) dans l'intimité des personnes, je tairai les secrets qui me seront confiés. Reçu(e) à l'intérieur des maisons, je respecterai les secrets des foyers et ma conduite ne servira pas à corrompre les mœurs.

Je ferai tout pour soulager les souffrances. Je ne prolongerai pas abusivement les agonies. Je ne provoquerai jamais la mort délibérément.

Je préserverai l'indépendance nécessaire à l'accomplissement de ma mission. Je n'entreprendrai rien qui dépasse mes compétences. Je les entretiendrai et les perfectionnerai pour assurer au mieux les services qui me seront demandés.

J'apporterai mon aide à mes confrères ainsi qu'à leurs familles dans l'adversité. Que les hommes et mes confrères m'accordent leur estime si je suis fidèle à mes promesses ; que je sois déshonoré(e) et méprisé(e) si j'y manque.

”